



# Modelling is key to forecast how climate change will impact ecosystems

- ▶ **Credible** model projections are critical for **natural resource managers**



Need a thorough model **evaluation**

# Modelling is key to forecast how climate change will impact ecosystems


- ▶ **Credible** model projections are critical for **natural resource managers**




Need a thorough model **evaluation**

- ▶ One approach: compare model predictions to observations from **previous time periods** (hindcasting)


## Decrease of model performance in the past




Future climatic conditions are expected to be challenging



# Future climatic conditions are expected to be challenging



# Future climatic conditions are expected to be challenging



## Urgency for action?

---

- ▶ Concerns about statistical model transferability into future climatic conditions
- ▶ Process-based models could provide more robust projections?



## Urgency for action?


- ▶ **Concerns about statistical model** transferability into future climatic conditions
- ▶ **Process-based** models could provide **more robust** projections?

 Yet to be verified !

# What conveys model robustness and transferability?

---

Correlative models  
(SDM)




- ▶ **statistical** relationships
- ▶ calibrated with current **distribution data**

# What conveys model robustness and transferability?

Correlative models  
(SDM)

Process-based models  
(PBM)



► **statistical** relationships

► **causal** relationships


► calibrated with current  
**distribution data**

► calibrated with  
**ecophysiological data**

# What conveys model robustness and transferability?

Correlative models  
(SDM)

Process-based models  
(PBM)



► **statistical** relationships



► **causal** relationships


► calibrated with current  
**distribution data**

► calibrated with  
**ecophysiological data**

# What conveys model robustness and transferability?

Correlative models  
(SDM)

Process-based models  
(PBM)



► statistical relationships



► causal relationships

► calibrated with current  
distribution data




► calibrated with  
ecophysiological data

# What conveys model robustness and transferability?

Correlative models  
(SDM)

Process-based models  
(PBM)



**Fitted** process-based models

► calibrated with current  
distribution data

► **causal** relationships

# Multi-model comparison across the Holocene

---


- ▶ **5** European tree species (*F. sylvatica*, *A. alba*, *Q. ilex*, *Q. robur*, *Q. petraea*)
- ▶ **8** models:
  - 4 correlative models (RF, BRT, GLM, GAM)
  - 2 fitted process-based models
  - 2 process-based models (PHENOFIT, CASTANEA)

# Multi-model comparison across the Holocene

---

- ▶ 5 European tree species (*F. sylvatica*, *A. alba*, *Q. ilex*, *Q. robur*, *Q. petraea*)
- ▶ 8 models:
  - 4 correlative models (RF, BRT, GLM, GAM)
  - 2 fitted process-based models
  - 2 process-based models (**PHENOFIT**, CASTANEA)

model based  
on phenology!






# Multi-model comparison across the Holocene

---

► 5 European tree species (*F. sylvatica*, *A. alba*, *Q. ilex*, *Q. robur*, *Q. petraea*)

► 8 models:    4 correlative models (RF, BRT, GLM, GAM)  
                  2 fitted process-based models  
                  2 process-based models (**PHENOFIT**, CASTANEA)


model based  
on **phenology!**




► past climate data from **HadCM3B** simulations (+ daily weather generation)

► model performance evaluated against **fossil pollen data**


# Example of deciduous *Quercus* paleosimulations




Decrease of model performance when moving further into the past




## Decrease of model performance when moving further into the past



# Process-based models are more performant/transferable?



# Process-based models are more performant/transferable




## Mechanisms, not calibration method, convey model robustness

- ▶ Expert and fitted **PBMs less affected** by the increase in climatic dissimilarity
- ▶ Fitted PBMs **similar** to expert PBMs

## Mechanisms, not calibration method, convey model robustness


- ▶ Expert and fitted **PBM**s **less affected** by the increase in climatic dissimilarity
- ▶ Fitted **PBM**s **similar** to expert **PBM**s



**Biological mechanisms** are necessary to improve the robustness of models  
(more important than the calibration method?)



## A real advantage to increase projections reliability in the upcoming decades?



## On our way to scale-up process-based models

---

- ▶ Fitted PBMs bring together the strengths from statistical and mechanistic approaches
- ▶ Inverse calibration: an opportunity to improve parameter estimates?

## On our way to scale-up process-based models

- ▶ Fitted PBMs **bring together the strengths** from statistical and mechanistic approaches
- ▶ Inverse calibration: an opportunity to **improve parameter estimates?**



→ **Spread the use** of process-based approaches?