

Numerical Analysis PhD Exam, May 2017, Part 1

Do 4 (four) problems.

1. This problem has two parts.

- (a) Given $A \in \mathbb{C}^{m \times n}$ with $m \geq n$, show that A^*A is nonsingular if and only if A has full rank.
- (b) If $u, v \in \mathbb{C}^m$ and $A = uv^*$, show that $\|A\|_2 = \|u\|_2\|v\|_2$.

2. Assume $S \in \mathbb{C}^{m \times m}$ is skew-Hermitian, so $S^* = -S$.

- (a) Show that the eigenvalues of S are pure imaginary.
- (b) Show that $I - S$ is nonsingular.
- (c) Show that the matrix $Q = (I - S)^{-1}(I + S)$ is unitary.

3. If $A \in \mathbb{R}^{m \times n}$ with $m \geq n$, $\text{rank}(A) = n$ and $b \in \mathbb{R}^m$.

- (a) Define the least squares solution to $Ax = b$.
- (b) Derive the normal equations for the least squares problem.
- (c) Prove that the unique solution to the least squares problem is $(A^T A)^{-1} A^T b$.
- (d) Describe how to solve the least squares problem using the SVD decomposition of A .

4. This problem has two parts.

- (a) Prove that every square matrix A has a Schur factorization.
- (b) If A is normal (so $A^*A = AA^*$) show that the triangular matrix in its Schur factorization is diagonal.

5. Assume $A \in \mathbb{C}^{m \times m}$.

- (a) If A has a collection of m linearly independent eigenvectors, show that A is diagonalizable.
- (b) When A is diagonalizable, prove the Cayley–Hamilton theorem for A , i.e. A satisfies its own characteristic polynomial.