

Probability PhD Exam, September 1997

1. Let X and Y be independent and uniform on $[0, 1]$. Find
 - a) the distribution of $X + Y$.
 - b) the conditional density of X given $X + Y = z$.
2. Let $X_0, X_1, \dots, X_n, \dots$ be i.i.d. with mean m . Let N be Poisson with mean λ , independent of the X 's. Define

$$Y = X_0 + \dots + X_N.$$

Prove that Y is integrable. Find $E(Y)$.

3. State precisely (all for i.i.d.):
 - a) the strong law of large numbers and prove it.
 - b) the central limit theorem.
 - c) the law of the iterated logarithm.
4. Define a martingale relative to a filtration $\{\mathcal{F}_n\}$, $n = 0, 1, 2, \dots$. Define a stopping time relative to this filtration. State the optional sampling theorem. State and prove the martingale convergence theorem.
5. What is an infinitely divisible distribution on $\mathbb{R}^1$? Prove that a non-trivial infinitely divisible distribution cannot have compact support.
6. Define a standard Brownian Motion W starting at 0. Define new processes W_1, W_2, W_3 by:

$$W_1(t) = cW\left(\frac{t}{c^2}\right), \quad t \geq 0, \quad c \text{ real } \neq 0,$$

$$W_2(t) = tW\left(\frac{1}{t}\right), \quad t > 0, \quad W_2(0) = 0,$$

$$W_3(t) = \begin{cases} W(1) - W(1-t), & 0 \leq t \leq 1, \\ W(t), & \text{otherwise.} \end{cases}$$

Show that W_1, W_2, W_3 are the standard Brownian motions.

7. Let $\varphi(t) = \int e^{itx} \mu(dx)$ be a characteristic function where μ is a probability measure on $\mathbb{R}^1$.
 - a) Suppose $|\varphi(t)| = 1$ for some $t \neq 0$. Then prove that unless $|\varphi(t)| \equiv 1$, there is a smallest $t_0 \neq 0$ and a d such that μ is concentrated on the set

$$\left\{d + \frac{2\pi j}{t_0}\right\}, \quad j = 0, \pm 1, \pm 2, \dots$$

- b) Using a) prove that $|\cos t|$ is not a characteristic function. Hint: $\cos t$ and $\cos^2 t$ are characteristic functions.

8. Let $S_n = (U_n, V_n)$ denote the position after n steps of a random walk on $\mathbb{Z}^2$, starting from $(0, 0)$. Suppose that

$$U_{n+1} = U_n \pm 1, \quad V_{n+1} = V_n \pm 1,$$

where the signs are picked by two independent tosses of a fair coin, independently at each step. Define

$$p_n = P(S_n = (0, 0)),$$

$$\tau_n = \inf\{m > \tau_{n-1} : S_m = (0, 0)\}.$$

An easy fact is $P(\tau_n < \infty) = P(\tau_1 < \infty)^n$.

- (1) Prove that for any random walk $S_n \in \mathbb{Z}^2$ the following are equivalent:

- (i) $P(\tau_n < \infty) = 1$.
- (ii) $P(S_n = (0, 0) \text{ i.o.}) = 1$.
- (iii) $\sum_{n=1}^{\infty} P(S_n = (0, 0)) = \infty$.

- (2) Find a formula for p_n , $n > 0$.

- (3) Use (1) and Stirling's formula

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^S \quad (S \rightarrow 0 \text{ as } n \rightarrow \infty)$$

to show that the diagonal random walk in $\mathbb{Z}^2$ is recurrent.

9. Let $(P_t)_{t>0}$ be a Markov semigroup. Assume $P_t(x, dy) = p_t(x, y) dy$ with $p_t(x, y) = p_t(y, x)$ and $p_t(x, y) \leq M(t)$, where $M(t)$ is a constant depending on t . If μ is a finite signed measure, show that

$$\int p_t(x, y) \mu(dx) \mu(dy) \geq 0.$$

10. An urn at $t = 0$ contains R_0 (≥ 1) red balls and B_0 (≥ 1) black balls, and “random sampling with double replacement” is called out as follows:

- (i) Choose a ball at random (meaning that each ball in the urn has the same probability to be drawn), and note its color;
- (ii) Replace this ball in the urn together with an extra ball of the same color;
- (iii) Go to (i).

Let M_t be the proportion in red balls in the urn after t such operations, so that

$$M_0 = \frac{R_0}{R_0 + B_0}.$$

Show that, with respect to a suitable filtration (which you should specify), the sequence $M_0, M_1, \dots, M_t, \dots$ is a martingale. Prove that M_t converges as $t \rightarrow +\infty$ almost surely to a random variable M_∞ , $0 \leq M_\infty \leq 1$. Show that

$$M_0 = EM_1 = EM_2 = \dots = EM_t = EM_\infty,$$

and that

$$M_0^2 \leq E[M_1^2] \leq \dots \leq E[M_t^2] \leq E[M_\infty^2],$$

and that

$$P(M_\infty = M_0) < 1.$$

NOTE. You may quote without proof any martingale theorems you need, but should state them carefully.

11. Let X_n be independent and all distributed according to standard normal distribution. Find constants C_n such that

$$\limsup \frac{X_n}{C_n} = 1 \quad \text{a.s.}$$

Hint.

$$\int_a^\infty e^{-b^2/2} db \leq \frac{1}{a} e^{-a^2/2},$$

and let $c_n^2 = 2 \log(n \log n)$.