

Analysis PhD Exam, May 2015

Do six of eight. Write solutions in a neat and logical fashion, giving complete reasons for all steps.

1. Let X and Y be topological spaces. Prove, if $f : X \rightarrow Y$ is continuous, then f is $\mathcal{B}_X$ - $\mathcal{B}_Y$ measurable (that is, measurable with respect to the Borel σ -algebras on X and Y respectively).
2. (Short Answer.) Do two of three.

- (i) Give an example, if possible, of a closed subset F of $\mathbb{R}$ of positive measure which contains no nontrivial open interval.
- (ii) Give an example, if possible, which shows that the σ -finite hypothesis is needed in Tonelli's Theorem.
- (iii) Determine the limit of the sequence,

$$\int_1^\infty \frac{\sin(nx)}{x^n} dx.$$

3. Let $(X, \mathcal{M}, \mu)$ be a measure space and $(f_n)_{n=1}^\infty$ be a sequence from $L^1(\mu)$. Prove, if there is a $g \in L^1(\mu)$ such that $g(x) \geq |f_n(x)|$ for all n and $x \in X$, then

$$\limsup_{\delta \rightarrow 0} \int_n \int_{\{|f_n| < \delta\}} |f_n| d\mu = 0.$$

4. Prove, if $E \subset \mathbb{R}$ is a Lebesgue measurable set of positive measure, then $E - E = \{x - y : x, y \in E\}$ contains a nontrivial open interval.
5. Let $(X, \mathcal{M}, \mu)$ be a measure space and $1 \leq p < q < \infty$. Prove, $L^q(\mu) \subset L^p(\mu)$ if and only if there is a (real) constant K such that for each $E \in \mathcal{M}$ either $\mu(E) \leq K$ or $\mu(E) = \infty$.
6. Prove, if X is a Banach space and M is a finite dimensional subspace of X , then there is a closed subspace N of X such that $M \cap N = \{0\}$ and $M + N = X$.
7. Suppose $\|\cdot\|$ and $\|\cdot\|_*$ are both norms on a vector space X and $\|x\| \leq \|x\|_*$ for all $x \in X$. Show, if both $(X, \|\cdot\|)$ and $(X, \|\cdot\|_*)$ are Banach spaces, then these norms are equivalent.
8. (Short Answer. Do two of three)
 - (a) Explain why c_{00} can not be a Banach space.
 - (b) Explain why there is no bounded one-one onto linear map from c_0 to ℓ^∞ .
 - (c) Explain what is meant by the Fourier transform of a function $f \in L^2(\mathbb{R})$.