

Analysis PhD Exam, May 2011

Answer SIX questions. Write solutions in a neat and logical fashion, giving complete reasons for all steps and stating carefully any substantial theorems used.

1. This problem has two parts.
 - (i) State the Hahn–Banach theorem.
 - (ii) Show that there exists an isometric linear map from each separable normed space into ℓ_∞ .
2. Prove that the spaces c (of all convergent scalar sequences) and c_0 (of all null scalar sequences) are not isometrically isomorphic when equipped with the sup norm.
3. State the Closed Graph Theorem and the Banach Isomorphism Theorem; deduce one of these from the other.
4. Let the sequence $(T_n)_{n=1}^\infty \subset L(X, Y)$ be bounded in operator norm and assume Y complete. Prove that

$$Z = \{z \in X : (T_n z)_{n=1}^\infty \text{ converges}\}$$

is a closed subspace of X .

5. Let X and Y be closed subspaces of a Hilbert space. Prove that if $X \perp Y$, then $X + Y$ is closed.
6. Let $(\Omega, \mathcal{F}, \mu)$ be a measure space. Define $\overline{\mathcal{F}}$ to comprise all those $A \subseteq \Omega$ for which there exist $L, U \in \mathcal{F}$ such that $L \subseteq A \subseteq U$ and $\mu(U \setminus L) = 0$ and then define $\overline{\mu}(A)$ to be the common value $\mu(L) = \mu(U)$. Show that $(\Omega, \overline{\mathcal{F}}, \overline{\mu})$ is a measure space that is complete in the sense that each subset of a null set is null.
7. Let $(\Omega, \mathcal{F}, \mu)$ be an arbitrary measure space; let $p, q, r > 1$ satisfy $\frac{1}{p} + \frac{1}{q} = \frac{1}{r}$. Prove that if $f \in L_p(\mu)$ and $g \in L_q(\mu)$ then $fg \in L_r(\mu)$ with

$$\|fg\|_r \leq \|f\|_p \|g\|_q.$$

8. Let $1 \leq p < q < \infty$. Show that $L_p(\mu) \not\subseteq L_q(\mu)$ iff $(\Omega, \mathcal{F}, \mu)$ contains measurable sets of arbitrarily small positive measure.