

# 2+1 Gravity as a Gauge Theory

Vicente V. Figueira


II Agorá Meeting — IFUSP

October 23, 2025

# General Relativity

DFMA  
IFUSP

$$S_{\text{EH}} = \frac{1}{2\kappa} \int_M d^D x \sqrt{|g|} g^{ab} R_{ca}{}^c{}_b$$



# General Relativity

DFMA  
IFUSP

$$S_{\text{EH}} = \frac{1}{2\kappa} \int_M d^D x \sqrt{|g|} g^{ab} R_{ca}{}^c{}_b$$

- Non-polynomial  $\rightarrow$  Hard to quantize.

# General Relativity

DFMA  
IFUSP

$$S_{\text{EH}} = \frac{1}{2\kappa} \int_M d^D x \sqrt{|g|} g^{ab} R_{ca}{}^c{}_b$$

- Non-polynomial  $\rightarrow$  Hard to quantize.
- Dimensionful coupling  $\rightarrow$  Not perturbatively renormalizable.


# General Relativity

DFMA  
IFUSP

$$S_{\text{EH}} = \frac{1}{2\kappa} \int_M d^D x \sqrt{|g|} g^{ab} R_{ca}{}^c{}_b$$

- Non-polynomial  $\rightarrow$  Hard to quantize.
- Dimensionful coupling  $\rightarrow$  Not perturbatively renormalizable.
- Has diffeomorphism redundancies:  $\phi_* \mathbf{g} \cong \mathbf{g}$ .

# General Relativity


$$S_{\text{EH}} = \frac{1}{2\kappa} \int_M d^D x \sqrt{|g|} g^{ab} R_{ca}{}^c{}_b$$

- Non-polynomial  $\rightarrow$  Hard to quantize.
- Dimensionful coupling  $\rightarrow$  Not perturbatively renormalizable.
- Has diffeomorphism redundancies:  $\phi_* \mathbf{g} \cong \mathbf{g}$ .


We know how to quantize a class of theories with redundancies:

# Yang-Mills

$$S_{\text{YM}} = \frac{1}{g^2} \int_M \text{Tr} [\mathbf{F} \wedge \star \mathbf{F}], \quad \mathbf{F} = d\mathbf{A} + [\mathbf{A} \wedge \mathbf{A}]$$

- Polynomial.


# Yang-Mills


$$S_{\text{YM}} = \frac{1}{g^2} \int_M \text{Tr} [\mathbf{F} \wedge \star \mathbf{F}], \quad \mathbf{F} = d\mathbf{A} + [\mathbf{A} \wedge \mathbf{A}]$$

- Polynomial.
- Dimensionless coupling  $\rightarrow$  Renormalizable.




# Yang-Mills



$$S_{\text{YM}} = \frac{1}{g^2} \int_M \text{Tr} [\mathbf{F} \wedge \star \mathbf{F}], \quad \mathbf{F} = d\mathbf{A} + [\mathbf{A} \wedge \mathbf{A}]$$

- Polynomial.
- Dimensionless coupling  $\rightarrow$  Renormalizable.
- Has local Lie group-valued redundancies,  $\mathbf{A} \cong U(\mathbf{A} + d)U^{-1}$ .

# Yang-Mills



$$S_{\text{YM}} = \frac{1}{g^2} \int_M \text{Tr} [\mathbf{F} \wedge \star \mathbf{F}], \quad \mathbf{F} = d\mathbf{A} + [\mathbf{A} \wedge \mathbf{A}]$$

- Polynomial.
- Dimensionless coupling  $\rightarrow$  Renormalizable.
- Has local Lie group-valued redundancies,  $\mathbf{A} \cong U(\mathbf{A} + d)U^{-1}$ .

It's possible to formulate GR as YM theory?

# Why $D = 2 + 1$ ?

DFMA  
IFUSP


- No local propagating modes:

$$\text{d.o.f.} = \frac{1}{2}D(D-3)$$


Simpler but non-trivial toy model.

# Recover of polynomiality

- Change of variables:  $\mathbf{g} \rightarrow \begin{cases} \mathbf{e}^\mu, & \text{Vielbein} \\ \omega_{\alpha\beta}, & \text{Spin connection} \end{cases}$



# Recover of polynomiality

- 
- Change of variables:  $\mathbf{g} \rightarrow \begin{cases} \mathbf{e}^\mu, & \text{Vielbein} \\ \omega_{\alpha\beta}, & \text{Spin connection} \end{cases}$
  - Relax torsionless condition  $\rightarrow \mathbf{e}^\mu$  and  $\omega_{\alpha\beta}$  are independent.

# Recover of polynomiality

- Change of variables:  $\mathbf{g} \rightarrow \begin{cases} \mathbf{e}^\mu, & \text{Vielbein} \\ \omega_{\alpha\beta}, & \text{Spin connection} \end{cases}$
- Relax torsionless condition  $\rightarrow \mathbf{e}^\mu$  and  $\omega_{\alpha\beta}$  are independent.

$$S_{\text{EH}} = \frac{1}{2\kappa} \int_M \star \mathbf{R}^{\alpha\beta} \wedge \mathbf{e}_\alpha \wedge \mathbf{e}_\beta \stackrel{D=3}{=} \frac{1}{2\kappa} \epsilon_{\alpha\beta\mu} \int_M \mathbf{R}^{\alpha\beta} \wedge \mathbf{e}^\mu$$

# Recover of polynomiality

- Change of variables:  $\mathbf{g} \rightarrow \begin{cases} \mathbf{e}^\mu, & \text{Vielbein} \\ \omega_{\alpha\beta}, & \text{Spin connection} \end{cases}$
- Relax torsionless condition  $\rightarrow \mathbf{e}^\mu$  and  $\omega_{\alpha\beta}$  are independent.

$$S_{\text{EH}} = \frac{1}{2\kappa} \int_M \star \mathbf{R}^{\alpha\beta} \wedge \mathbf{e}_\alpha \wedge \mathbf{e}_\beta \stackrel{D=3}{=} \frac{1}{2\kappa} \epsilon_{\alpha\beta\mu} \int_M \mathbf{R}^{\alpha\beta} \wedge \mathbf{e}^\mu$$

Can we write the integrand as a trace?

# Recover of dimensionless coupling constant

- For  $\text{iso}(2,1)$ :

$$\text{Tr}[J_{\alpha\beta}P_{\mu}] = \frac{\lambda}{\kappa}\epsilon_{\alpha\beta\mu}, \quad \text{Tr}[P_{\nu}P_{\mu}] = 0, \quad \text{Tr}[J_{\alpha\beta}J_{\mu\nu}] = 0$$



# Recover of dimensionless coupling constant

- For  $\mathfrak{iso}(2,1)$ :

$$\mathrm{Tr}[J_{\alpha\beta}P_{\mu}] = \frac{\lambda}{\kappa}\epsilon_{\alpha\beta\mu}, \quad \mathrm{Tr}[P_{\nu}P_{\mu}] = 0, \quad \mathrm{Tr}[J_{\alpha\beta}J_{\mu\nu}] = 0$$

- Dressing the fields with the algebra,

$$S_{\mathrm{EH}} = \frac{1}{\lambda} \int_M \mathrm{Tr}[\mathbf{R} \wedge \mathbf{e}], \quad \mathbf{R} = \frac{1}{2} \mathbf{R}^{\alpha\beta} J_{\alpha\beta}, \quad \mathbf{e} = \mathbf{e}^{\mu} P_{\mu}$$

# Recover of dimensionless coupling constant

- For  $\mathfrak{iso}(2,1)$ :

$$\mathrm{Tr}[J_{\alpha\beta}P_\mu] = \frac{\lambda}{\kappa}\epsilon_{\alpha\beta\mu}, \quad \mathrm{Tr}[P_\nu P_\mu] = 0, \quad \mathrm{Tr}[J_{\alpha\beta}J_{\mu\nu}] = 0$$


- Dressing the fields with the algebra,

$$S_{\mathrm{EH}} = \frac{1}{\lambda} \int_M \mathrm{Tr}[\mathbf{R} \wedge \mathbf{e}], \quad \mathbf{R} = \frac{1}{2} \mathbf{R}^{\alpha\beta} J_{\alpha\beta}, \quad \mathbf{e} = \mathbf{e}^\mu P_\mu$$

Not standard YM form...

# Chern-Simons

- d.o.f. mismatch: 0 v.s.  $2N_c$



# Chern-Simons

- d.o.f. mismatch: 0 v.s.  $2N_c$
- Available gauge theory with 0 d.o.f.: Chern-Simons!

$$S_{\text{CS}} = \frac{1}{2\lambda} \int_M \text{Tr} \left[ \mathbf{A} \wedge \left( d\mathbf{A} + \frac{1}{3} [\mathbf{A} \wedge \mathbf{A}] \right) \right]$$

# Chern-Simons

- d.o.f. mismatch: 0 v.s.  $2N_c$
- Available gauge theory with 0 d.o.f.: Chern-Simons!

$$S_{\text{CS}} = \frac{1}{2\lambda} \int_M \text{Tr} \left[ \mathbf{A} \wedge \left( d\mathbf{A} + \frac{1}{3} [\mathbf{A} \wedge \mathbf{A}] \right) \right]$$

- Tentative: CS theory of  $\text{ISO}(2,1)$ ,  $\mathbf{A} = \boldsymbol{\omega} + \mathbf{e}$ ,

# Chern-Simons

- d.o.f. mismatch: 0 v.s.  $2N_c$
- Available gauge theory with 0 d.o.f.: Chern-Simons!

$$S_{\text{CS}} = \frac{1}{2\lambda} \int_M \text{Tr} \left[ \mathbf{A} \wedge \left( d\mathbf{A} + \frac{1}{3} [\mathbf{A} \frown \mathbf{A}] \right) \right]$$

- Tentative: CS theory of  $\text{ISO}(2,1)$ ,  $\mathbf{A} = \boldsymbol{\omega} + \mathbf{e}$ ,


$$S_{\text{CS}} = \frac{1}{\lambda} \int_M \left\langle \mathbf{e} \frown \left( d\boldsymbol{\omega} + \frac{1}{2} [\boldsymbol{\omega} \frown \boldsymbol{\omega}] \right) \right\rangle = S_{\text{EH}}$$

Match!

# Conclusions

DFMA  
IFUSP

- $2+1$  dimensional gravity is a CS gauge theory of  $\text{ISO}(2,1)$ .




# Conclusions

DFMA  
IFUSP

- $2 + 1$  dimensional gravity is a CS gauge theory of  $\text{ISO}(2,1)$ .
- Opens path to quantization via well known CS methods.



# Conclusions

- 
- $2 + 1$  dimensional gravity is a CS gauge theory of  $\text{ISO}(2,1)$ .
  - Opens path to quantization via well known CS methods.
  - Inclusion of cosmological constant change gauge group,  
$$\begin{cases} \Lambda > 0 : & \text{SO}(3,1), \text{ (dS)} \\ \Lambda < 0 : & \text{SO}(2,2), \text{ (AdS)} \end{cases}$$

# Conclusions

- 2 + 1 dimensional gravity is a CS gauge theory of  $\text{ISO}(2,1)$ .
- Opens path to quantization via well known CS methods.
- Inclusion of cosmological constant change gauge group,  
$$\begin{cases} \Lambda > 0 : & \text{SO}(3,1), \text{ (dS)} \\ \Lambda < 0 : & \text{SO}(2,2), \text{ (AdS)} \end{cases}$$
- Offers a bridge between GR and usual gauge theory intuition.


# Conclusions

- $2 + 1$  dimensional gravity is a CS gauge theory of  $\text{ISO}(2,1)$ .
- Opens path to quantization via well known CS methods.
- Inclusion of cosmological constant change gauge group,  
$$\begin{cases} \Lambda > 0 : & \text{SO}(3,1), \text{ (dS)} \\ \Lambda < 0 : & \text{SO}(2,2), \text{ (AdS)} \end{cases}$$
- Offers a bridge between GR and usual gauge theory intuition.
- Pose some suggestions about  $D = 3 + 1$ .


# DFMA

IFUSP

Thank You!



# References

- 
- [1] Edward Witten. “(2+1)-Dimensional Gravity as an Exactly Soluble System”. In: *Nucl. Phys. B* 311 (1988), p. 46. DOI: 10.1016/0550-3213(88)90143-5.
- [2] Edward Witten. *Three-Dimensional Gravity Revisited*. 2007. arXiv: 0706.3359 [hep-th]. URL: <https://arxiv.org/abs/0706.3359>.