

# Homework III

Vicente V. Figueira — NUSP 11809301

June 2, 2025

## Contents

<b>Problem 1</b>	<b>1</b>
1.A) . . . . .	1
1.B) . . . . .	2
1.C) . . . . .	2
1.D) . . . . .	2
<b>Problem 2</b>	<b>3</b>
2.A) . . . . .	3
2.B) . . . . .	3
2.C) . . . . .	3
2.D) . . . . .	3
<b>Problem 3</b>	<b>4</b>
3.A) . . . . .	4
3.B) . . . . .	4
3.C) . . . . .	4
3.D) . . . . .	4
3.E) . . . . .	4
3.F) . . . . .	4
<b>A Faddeev-Popov Gauge Fixing</b>	<b>5</b>
<b>B BRST Quantization</b>	<b>10</b>
<b>C Closure of BRST transformations</b>	<b>15</b>

## Problem 1

### 1.A)

For an operator  $\mathcal{O}$  to be BRST closed, it means  $[Q_{\text{BRST}}, \mathcal{O}] = 0$ , just remembering the usual BRST transformations<sup>1</sup>,

$$\begin{aligned}[Q_{\text{BRST}}, X^\mu] &= (c\partial + \tilde{c}\bar{\partial})X^\mu \\ [Q_{\text{BRST}}, b] &= T^X + T^g\end{aligned}$$

---

<sup>1</sup>We're using a graded commutator notation, that is, it's to be interpreted as either a commutator or an anti-commutator depending on the statistic of what is inside.

$$[Q_{\text{BRST}}, c] = c\partial c$$

with,

$$T^X = \frac{1}{\alpha'} : \partial X^\mu \partial X_\mu :, \quad T^g =: c\partial b : -2 : b\partial c :$$

so,

$$\begin{aligned} [Q_{\text{BRST}}, V^a] &= \lambda^a \epsilon_\mu [Q_{\text{BRST}}, c\partial X^\mu \exp(ik \cdot X)] \\ [Q_{\text{BRST}}, V^a] &= \lambda^a \epsilon_\mu [Q_{\text{BRST}}, c] \partial X^\mu \exp(ik \cdot X) - \lambda^a \epsilon_\mu c [Q_{\text{BRST}}, \partial X^\mu \exp(ik \cdot X)] \\ [Q_{\text{BRST}}, V^a] &= \lambda^a \epsilon_\mu c \partial c \partial X^\mu \exp(ik \cdot X) \\ &\quad - \lambda^a \epsilon_\mu c \partial X^\mu [Q_{\text{BRST}}, \exp(ik \cdot X)] - \lambda^a \epsilon_\mu c [Q_{\text{BRST}}, \partial X^\mu] \exp(ik \cdot X) \\ [Q_{\text{BRST}}, V^a] &= \lambda^a \epsilon_\mu c \partial c \partial X^\mu \exp(ik \cdot X) \\ &\quad - \lambda^a \epsilon_\mu c \partial X^\mu [Q_{\text{BRST}}, \exp(ik \cdot X)] - \lambda^a \epsilon_\mu c \partial [Q_{\text{BRST}}, X^\mu] \exp(ik \cdot X) \\ [Q_{\text{BRST}}, V^a] &= \lambda^a \epsilon_\mu c \partial c \partial X^\mu \exp(ik \cdot X) \\ &\quad - \lambda^a \epsilon_\mu c \partial X^\mu [Q_{\text{BRST}}, \exp(ik \cdot X)] - \lambda^a \epsilon_\mu c \partial (c\partial + \tilde{c}\bar{\partial}) X^\mu \exp(ik \cdot X) \\ [Q_{\text{BRST}}, V^a] &= \lambda^a \epsilon_\mu c \partial c \partial X^\mu \exp(ik \cdot X) \\ &\quad - \lambda^a \epsilon_\mu c \partial X^\mu [Q_{\text{BRST}}, \exp(ik \cdot X)] - \lambda^a \epsilon_\mu c \partial (c\partial + \tilde{c}\bar{\partial}) X^\mu \exp(ik \cdot X) \\ &\quad - \lambda^a \epsilon_\mu c c \partial - \lambda^a \epsilon_\mu c \partial (\tilde{c}\bar{\partial}) X^\mu \exp(ik \cdot X) \end{aligned}$$

**1.B)**

**1.C)**

**1.D)**

## Problem 2

2.A)

2.B)

2.C)

2.D)

### Problem 3

3.A)

3.B)

3.C)

3.D)

3.E)

3.F)

## A Faddeev-Popov Gauge Fixing

Our Action functional is,

$$S_X + \lambda\chi = \frac{1}{4\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} \partial_a X^\mu \partial_b X_\mu + \frac{\lambda}{4\pi} \int_M d^2\sigma \sqrt{h} R + \frac{\lambda}{2\pi} \int_{\partial M} ds K \quad (\text{A.1})$$

we would like to define the quantum theory by means of the path integral, that is, we expect that,

$$Z \stackrel{?}{=} \int \mathcal{D}X \mathcal{D}h \exp(-S_X[X, h] - \lambda\chi) \quad (\text{A.2})$$

should give a well defined theory, but, already from A.2 there're several problems that arise, one of them is: *What should be interpreted from the path integral itself? We haven't defined any manifold to our metric  $h$  and scalar fields  $X$  to live in, also, even if we had defined such, the path integral relies on explicit coordinate points,  $\mathcal{D}h = \prod_\sigma dh_{ab}(\sigma)$ , which are highly dependent on charts.*

This is a valid claim, our way to avoid it is to *define*  $\mathcal{D}h$  to mean: *Sum over all **allowed** two dimensional Riemannian manifolds, and all possible metric structures in these.* Here, **allowed** requires a prescription, which manifolds are or aren't allowed impacts the obtained string theory. Happily, every two dimensional manifold has a definite value for the Euler Characteristic  $\chi$ , hence, we can sort them out by it,

$$\begin{aligned} Z &\stackrel{?}{=} \sum_{\{M\}_{\text{Met}(M)}} \int \mathcal{D}h \int \mathcal{D}X \exp(-S_X[X, h] - \lambda\chi) \\ Z &\stackrel{?}{=} \sum_{\{\chi\}} \exp(-\lambda\chi) \sum_{\{M_\chi\}_{\text{Met}(M_\chi)}} \int \mathcal{D}h \int \mathcal{D}X \exp(-S_X[X, h]) \end{aligned} \quad (\text{A.3})$$

Where  $M$  is to be understood as a two dimensional Riemannian manifold and  $M_\chi$  is one with Euler Characteristic  $\chi$ ,  $\text{Met}(M_\chi)$  is the space of all metrics which can be assigned to  $M_\chi$ , we have written  $\sum_{\{M_\chi\}}$  in the special case of there being more than one manifold with same Euler Characteristic<sup>2</sup>, also, the functional integral over  $X$  should be read as integrating over all maps from  $M_\chi$  to  $\mathbb{R}^{1,D-1}$ . While this is better defined than before, i.e. not coordinate dependent, we still have a few problems, first, it's known that A.1 has a Gauge Group of  $\text{Diff}(M) \times \text{Weyl}(M)$ , but, in our second try of a definition of the path integral, we're integrating the metrics over  $\text{Met}(M_\chi)$ , it's clear that may happen of two elements of  $\text{Met}(M_\chi)$  be equivalent under a  $\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$  transformation, to put in more clear terms, we're worried if exists  $h', h \in \text{Met}(M_\chi)$  such,

$$h'_{ab}(\sigma'(\sigma)) = \exp(2\omega(\sigma)) \frac{\partial \sigma^c}{\partial \sigma'^a} \frac{\partial \sigma^d}{\partial \sigma'^b} h_{cd}(\sigma)$$

the existence of those kinds of elements is troublesome, as  $\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$  is a infinite dimensional group of redundancies, this means we're over-counting physical configurations by a infinite amount. The solution is to look for an equivalence class of metrics under this Gauge Group action,

$$\mathcal{M}_\chi = \text{Met}(M_\chi) / \text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$$

---

<sup>2</sup>As we're interested only in Differentiable Manifolds, more than manifold should read: More than one equivalence class of Differentiable Manifolds.

the equivalence class is to be understood as<sup>3</sup>,

$$h' \sim h \Leftrightarrow h'_{ab}(\sigma'(\sigma)) = \exp(2\omega(\sigma)) \frac{\partial \sigma^c}{\partial \sigma'^a} \frac{\partial \sigma^d}{\partial \sigma'^b} h_{cd}(\sigma)$$

that is, two metrics are the same representative of an element of  $\mathcal{M}_\chi$  iff they differ by a composition of a Diffeomorphism and Weyl transformation. We'll denote a given composition of a Diffeomorphism followed by a Weyl transformation by  $\zeta$ ,

$$h' = \zeta \circ h$$

Notice that the set of equivalence class of metrics, or, the set of inequivalent  $\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$  metrics  $\mathcal{M}_\chi$  is highly dependent on the topology of  $M_\chi$ , for example, for  $M_\chi \cong \mathbb{R}^2 \cong \mathbb{C}$ , it's trivial, there is just one point in the set  $\mathcal{M}_\chi$ , in other words, every metric is equivalent, which isn't true for more complex topologies.

Thus, it's possible for us to set up a well defined version of the path integral, just replace  $\text{Met}(M_\chi)$  by  $\mathcal{M}_\chi$ ,

$$Z = \sum_{\{x\}} \exp(-\lambda\chi) \sum_{\{M_\chi\}} \int_{\mathcal{M}_\chi} \mathcal{D}h \int \mathcal{D}X \exp(-S_X[X, h]) \quad (\text{A.4})$$

where the integration is to be understood as by choosing for each equivalence class in  $\mathcal{M}_\chi$  a representative element in  $\text{Met}(M_g)$ . While this is a satisfactory definition for the path integral, it feels a little clunky, we rather have an path integral over all the possible metrics — in the sense defined before —, well, this is achievable. First, for each equivalence class of  $\mathcal{M}_\chi$  elect one representative element of  $\text{Met}(M_\chi)$ , we'll denote these elements as  $\hat{h}(\mathbf{t})$  — here  $\mathbf{t}$  is a parametrization of the correspondent equivalence class in  $\mathcal{M}_\chi$ , we haven't proved here, and won't, but  $\mathcal{M}_\chi$  is a finite  $N$  dimensional manifold, hence,  $\mathbf{t}$  is a  $N$ -tuple of real numbers —, by construction, these representatives are inequivalent under  $\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$ , hence,

$$\zeta_1 \circ \hat{h}(\mathbf{t}_1) = \zeta_2 \circ \hat{h}(\mathbf{t}_2) \Leftrightarrow \mathbf{t}_1 = \mathbf{t}_2 \text{ and } \zeta_1 = \zeta_2$$

so that every element in  $\text{Met}(M_g)$  can be written as a unique<sup>4</sup> composition of a given  $\zeta$  into a given  $\hat{h}(\mathbf{t})$ . Now, we rewrite the pictorial integral over  $\mathcal{M}_\chi$  is a more formal way, using the parametrization we just described,

$$\begin{aligned} Z &= \sum_{\{x\}} \exp(-\lambda\chi) \sum_{\{M_\chi\}} \int_{\mathcal{M}_\chi} d^N \mathbf{t} \int \mathcal{D}X \exp\left(-S_X[X, \hat{h}(\mathbf{t})]\right) \\ Z &= \sum_{\{x\}} \exp(-\lambda\chi) \sum_{\{M_\chi\}} \int_{\mathcal{M}_\chi} d^N \mathbf{t} \int_{\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)} \mathcal{D}\zeta \delta(\zeta) \int \mathcal{D}X \exp\left(-S_X[X, \hat{h}(\mathbf{t})]\right) \end{aligned}$$

in the last line we introduced a one by integrating<sup>5</sup> over the delta functional, as this integral picks only  $\zeta = 0$ , what should be understood as  $\zeta = \text{id}$  in the group, we can deform a little the

<sup>3</sup>In all charts.

<sup>4</sup>The uniqueness or not depends on a few factors, here we'll always, unless specified otherwise, interpret  $\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$  as the group *generated by* all possible compositions of Diffeomorphisms and Weyl transformations, but a element of it,  $\zeta$ , is not to be interpreted as a unique composition of Diffeomorphism and Weyl factors, as there might be some Diffeomorphism which are equivalent to Weyl transformations, what is indeed true is that every element  $\zeta$  of the Gauge Group is a unique combination of an element of  $\text{Diff}(M_\chi)/\text{Weyl}(M_\chi)$  and an element of  $\text{Weyl}(M_\chi)$ .

<sup>5</sup>Again, following the same remarks made before, the integral over  $\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$  should not be interpreted as integrating over the whole of  $\text{Diff}(M_\chi)$  and after integrating over the whole  $\text{Weyl}(M_\chi)$ , this would for sure be an over-counting, but rather should be interpreted as integrating over the whole group *generated by* compositions of  $\text{Diff}(M_\chi)$  and  $\text{Weyl}(M_\chi)$ , which is equivalent of integrating over the whole  $\text{Diff}(M_\chi)/\text{Weyl}(M_\chi)$ , and after integrating over the whole  $\text{Weyl}(M_\chi)$ .

integration to,

$$Z = \sum_{\{\chi\}} \exp(-\lambda\chi) \sum_{\{M_\chi\} \mathcal{M}_\chi} \int d^N \mathbf{t} \int_{\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)} \mathcal{D}\zeta \delta(\zeta) \int \mathcal{D}X \exp\left(-S_X\left[X, \zeta \circ \hat{h}(\mathbf{t})\right]\right) \quad (\text{A.5})$$

This is almost in the form that we would like, notice that we're integrating over the set of representative of the inequivalent metrics,  $d^N \mathbf{t}$ , and also over the whole group  $\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)$ ,  $\mathcal{D}\zeta$ , by construction, **every** metric in  $\text{Met}(M_\chi)$  can be written uniquely<sup>6</sup> as,

$$h = \zeta_{\mathbf{t}} \circ \hat{h}(\mathbf{t})$$

in other words, to integrate over  $d^N \mathbf{t} \mathcal{D}\zeta$  is to integrate over all metrics of the form  $\zeta \circ \hat{h}(\mathbf{t})$ , which is to integrate over all metrics  $h = \zeta \circ \hat{h}(\mathbf{t})$  in  $\text{Met}(M_\chi)$ ! We cannot yet make this change, due to the presence of an explicit dependence in  $\zeta$  at the functional delta. We'll eliminate it by means of a change of variable of the functional delta, notice that,

$$\delta\left(\hat{h}(\mathbf{t}) - \zeta \circ \hat{h}(\mathbf{t})\right)$$

picks up just the contribution of  $\zeta = 0$ , so it's a good candidate for a change of variables,

$$\delta\left(\hat{h}(\mathbf{t}) - \zeta \circ \hat{h}(\mathbf{t})\right) = \delta(\zeta) \left| \text{Det} \left[ \frac{\delta}{\delta\zeta} \left( \hat{h}(\mathbf{t}) - \zeta \circ \hat{h}(\mathbf{t}) \right) \right]_{\zeta=0} \right|^{-1}$$

let's compute step by step the right-hand side of this equation, as we're only interested in the solution of  $\zeta = 0$ , what matters is just the connected component to the identity of the Gauge Group, this is parametrized by a function  $\omega$  related to the Weyl transformation, and a vector field  $\xi$  related to the connected component to the identity of the Diffeomorphisms — there is an additional requirement of  $\xi$  not generating any transformation which can be undone by a Weyl transformation —, also, for ease of our manipulation, we'll write the expression inside the delta with respect to  $h$  instead of  $\hat{h}(\mathbf{t})$ <sup>7</sup>, that is,

$$\delta\left(\hat{h}(\mathbf{t}) - \zeta \circ \hat{h}(\mathbf{t})\right) = \delta(\zeta^{-1} \circ h - h) = \delta(\zeta^{-1}) \left| \text{Det} \left[ \frac{\delta}{\delta\zeta^{-1}} (\zeta^{-1} \circ h - h) \right]_{\zeta^{-1}=0} \right|^{-1}$$

one might worry about the  $\zeta^{-1}$  instead of the  $\zeta$ , but, the integration measure  $\mathcal{D}\zeta$  is formally a Haar measure in the Group, that means it's a group invariant measure, in other words,  $\mathcal{D}\zeta^{-1} = \mathcal{D}\zeta$ , so that we can forget about the inverse, now,

$$\begin{aligned} [\zeta \circ h]_{ab} &= [h]_{ab} + 2\omega[h]_{ab} + [\mathcal{L}_\xi h]_{ab} + \mathcal{O}(\omega^2, \xi^2, \omega\xi) \\ [\zeta \circ h]_{ab} &= [h]_{ab} + 2\omega[h]_{ab} + 2\nabla_{(a}\xi_{b)} + \mathcal{O}(\omega^2, \xi^2, \omega\xi) \end{aligned}$$

of course  $\nabla$  here is with respect to the  $h$  metric,

$$\begin{aligned} [\zeta \circ h]_{ab} - [h]_{ab} &= 2\omega[h]_{ab} + 2\nabla_{(a}\xi_{b)} + \mathcal{O}(\omega^2, \xi^2, \omega\xi) \\ \frac{\delta}{\delta\zeta}([\zeta \circ h]_{ab} - [h]_{ab}) &= \text{????} \end{aligned}$$

---

<sup>6</sup>With the remarks made before.

<sup>7</sup>We would have to carry out the  $\mathbf{t}$  dependence in  $h$  also, but, soon it will disappear as matter of uniting the integrals  $d^N \mathbf{t} \mathcal{D}\zeta$  so we won't keep track of it anymore.

The  $\zeta$  derivative actually has two parts, the derivative with respect to  $\omega$  and the other with respect to  $\xi$ , let's do one by one,

$$\left. \frac{\delta}{\delta\omega(\sigma')} ([\zeta \circ h]_{ab} - [h]_{ab})(\sigma) \right|_{\zeta=0} = 2\delta^{(2)}(\sigma - \sigma')h_{ab}(\sigma)$$

and for the  $\xi$ ,

$$\left. \frac{\delta}{\delta\xi^c(\sigma')} ([\zeta \circ h]_{ab} - [h]_{ab})(\sigma) \right|_{\zeta=0} = 2h_{c(b}\nabla_a)\delta^{(2)}(\sigma - \sigma')$$

Thus,

$$\begin{aligned} \left. \frac{\delta}{\delta\zeta} ([\zeta \circ h]_{ab} - [h]_{ab}) \right|_{\zeta=0} &= 2\delta^{(2)}(\sigma - \sigma')h_{ab}(\sigma) + 2h_{c(b}\nabla_a)\delta^{(2)}(\sigma - \sigma') \\ \text{Det} \left[ \left. \frac{\delta}{\delta\zeta} ([\zeta \circ h]_{ab} - [h]_{ab}) \right|_{\zeta=0} \right] &= \text{Det} [2\delta^{(2)}(\sigma - \sigma')h_{ab}(\sigma) + 2h_{c(b}\nabla_a)\delta^{(2)}(\sigma - \sigma')] \end{aligned}$$

the determinant can be computed by means of path integral of Grassmannian variables,

$$\begin{aligned} \text{Det} [2\delta^{(2)}(\sigma - \sigma')h_{ab} + 2h_{c(b}\nabla_a)\delta^{(2)}(\sigma - \sigma')] &= \int \mathcal{D}b\mathcal{D}c\mathcal{D}d \exp \left( -\frac{1}{2\pi} \int d^2\sigma \sqrt{h}b^{ab} [h_{ab}d + h_{c(b}\nabla_a)c^c] \right) \\ \text{Det} \left[ \left. \frac{\delta}{\delta\zeta} ([\zeta \circ h]_{ab} - [h]_{ab}) \right|_{\zeta=0} \right] &= \int \mathcal{D}b\mathcal{D}c\mathcal{D}d \exp (-S_{\text{gh}}[b, c, d, h]) \end{aligned}$$

Substituting all of this back into our path integral,

$$\begin{aligned} Z &= \sum_{\{\chi\}} \exp(-\lambda\chi) \sum_{\{M_\chi\} \mathcal{M}_\chi} \int d^N \mathbf{t} \int_{\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)} \mathcal{D}\zeta \delta(\zeta) \int \mathcal{D}X \exp \left( -S_X[X, \zeta \circ \hat{h}(\mathbf{t})] \right) \\ Z &= \sum_{\{\chi\}} \exp(-\lambda\chi) \sum_{\{M_\chi\} \mathcal{M}_\chi} \int d^N \mathbf{t} \int_{\text{Diff}(M_\chi) \times \text{Weyl}(M_\chi)} \mathcal{D}\zeta \delta(\zeta^{-1} \circ h - h) \int \mathcal{D}X \mathcal{D}b\mathcal{D}c\mathcal{D}d \exp(-S_X - S_{\text{gh}}) \\ Z &= \sum_{\{\chi\}} \exp(-\lambda\chi) \sum_{\{M_\chi\} \text{Met}(M_\chi)} \int \mathcal{D}h \delta(\hat{h} - h) \int \mathcal{D}X \mathcal{D}b\mathcal{D}c\mathcal{D}d \exp(-S_X[X, h] - S_{\text{gh}}[b, c, d, h]) \\ Z &= \int \mathcal{D}h \mathcal{D}X \mathcal{D}b\mathcal{D}c\mathcal{D}d \delta(\hat{h} - h) \exp(-S_X[X, h] - S_{\text{gh}}[b, c, d, h] - \lambda\chi) \end{aligned}$$

where  $\hat{h}$  is a family of choices of representatives of the equivalence classes of the Gauge equivalent metrics, of course this choice is dependent on the equivalence class  $h$  lies in, so, in a certain sense we have  $\hat{h} = \hat{h}[h]$ ,

$$Z = \int \mathcal{D}h \mathcal{D}X \mathcal{D}b\mathcal{D}c\mathcal{D}d \delta(\hat{h}[h] - h) \exp(-S_X[X, h] - S_{\text{gh}}[b, c, d, h] - \lambda\chi)$$

we express the delta functional in terms of a path integral,

$$\begin{aligned} Z &= \int \mathcal{D}h \mathcal{D}X \mathcal{D}b\mathcal{D}c\mathcal{D}d \mathcal{D}B \exp \left( \frac{i}{4\pi} \int d^2\sigma \sqrt{h} B^{ab} (\hat{h}_{ab}[h] - h_{ab}) \right) \exp(-S_X - S_{\text{gh}} - \lambda\chi) \\ Z &= \int \mathcal{D}h \mathcal{D}X \mathcal{D}b\mathcal{D}c\mathcal{D}d \mathcal{D}B \exp(-S_X[X, h] - S_{\text{gh}}[b, c, d, h] - S_{\text{gf}}[B, h] - \lambda\chi) \end{aligned}$$



where we lastly defined the Gauge Fixing Action. This is the final expression for our path integral with the identifications,

$$S_X[X, h] + \lambda\chi = \frac{1}{4\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} \partial_a X^\mu \partial_b X_\mu + \frac{\lambda}{4\pi} \int_M d^2\sigma \sqrt{h} R + \frac{\lambda}{2\pi} \int_{\partial M} ds K \quad (\text{A.6a})$$

$$S_{\text{gh}}[b, c, d, h] = \frac{1}{2\pi} \int_M d^2\sigma \sqrt{h} b^{ab} [h_{ab} d + \nabla_a c_b] \quad (\text{A.6b})$$

$$S_{\text{gf}}[B, h] = -\frac{i}{4\pi} \int_M d^2\sigma \sqrt{h} B^{ab} (\hat{h}_{ab}[h] - h_{ab}) \quad (\text{A.6c})$$

## B BRST Quantization

Following the action principle derived from the Faddeev-Popov Gauge Fixing A.6, we can describe it's BRST symmetry by the transformations of the *matter fields* under Gauge, we know the following,

$$\begin{aligned} X^\mu(\sigma) &\rightarrow X'^\mu(\sigma'(\sigma)) = X^\mu(\sigma) \\ h_{ab}(\sigma) &\rightarrow h'_{ab}(\sigma'(\sigma)) = \exp(2\omega(\sigma)) \frac{\partial \sigma^c}{\partial \sigma'^a} \frac{\partial \sigma^d}{\partial \sigma'^b} h_{cd}(\sigma) \end{aligned}$$

which have an *infinitesimal* form,

$$\begin{aligned} \delta X^\mu &= \mathcal{L}_\xi X^\mu = \xi^a \partial_a X^\mu \\ \delta h_{ab} &= 2\omega h_{ab} + \mathcal{L}_\xi h_{ab} = 2\omega h_{ab} + \nabla_a \xi_b + \nabla_b \xi_a = 2\omega h_{ab} + \xi^c \partial_c h_{ab} + h_{ac} \partial_b \xi^c + h_{bc} \partial_a \xi^c \end{aligned}$$

the BRST transformation can be obtained from these by the substitution inferred from the Faddeev-Popov gauge fix, that is,  $\xi^a \rightarrow i\epsilon c^a$  and  $\omega \rightarrow i\epsilon d$ , where  $\epsilon$  is a Grassmannian parametrization of the BRST transformation,

$$\delta_{\text{BRST}} X^\mu = i\epsilon c^a \partial_a X^\mu \quad (\text{B.1a})$$

$$\delta_{\text{BRST}} h_{ab} = 2i\epsilon (dh_{ab} + \nabla_{(a} c_{b)}) = 2i\epsilon dh_{ab} + i\epsilon (c^c \partial_c h_{ab} + h_{ac} \partial_b c^c + h_{bc} \partial_a c^c) \quad (\text{B.1b})$$

while the BRST transformation of the non-matter fields can be obtained by the BRST procedure, which are<sup>8</sup>,

$$\delta_{\text{BRST}} (\sqrt{h} B_{ab}) = 0 \quad (\text{B.1c})$$

$$\delta_{\text{BRST}} (\sqrt{h} b_{ab}) = -\epsilon \sqrt{h} B_{ab} \quad (\text{B.1d})$$

$$\delta_{\text{BRST}} d = i\epsilon c^a \partial_a d \quad (\text{B.1e})$$

$$\delta_{\text{BRST}} c^a = i\epsilon c^b \nabla_b c^a = i\epsilon c^b \partial_b c^a \quad (\text{B.1f})$$

the proof of nilpotency of these transformations is done in Appendix C. For the BRST quantization to be realized we need two things,

$$\begin{aligned} \delta_{\text{BRST}} (S_X + S_{\text{gh}} + S_{\text{gf}}) &= 0 \\ i\epsilon (S_{\text{gh}} + S_{\text{gf}}) &= \delta_{\text{BRST}} \mathcal{O} \end{aligned}$$

for  $\mathcal{O}$  some composition of fields, we'll check both of these, and for this it'll be necessary to know the BRST transformations of a few more fields,

- $h^{ab}$

$$\begin{aligned} 0 &= \delta_{\text{BRST}} \delta_a^c \\ 0 &= \delta_{\text{BRST}} (h_{ab} h^{bc}) \\ 0 &= h_{ab} \delta_{\text{BRST}} h^{bc} + h^{bc} \delta_{\text{BRST}} h_{ab} \\ h_{ab} \delta_{\text{BRST}} h^{bc} &= -2i\epsilon h^{bc} (dh_{ab} + \nabla_{(a} c_{b)}) \\ h^{da} h_{ab} \delta_{\text{BRST}} h^{bc} &= -2i\epsilon h^{da} h^{bc} (dh_{ab} + \nabla_{(a} c_{b)}) \\ \delta_b^d \delta_{\text{BRST}} h^{bc} &= -2i\epsilon (dh^{dc} + \nabla^{(d} c^{c)}) \\ \delta_{\text{BRST}} h^{dc} &= -2i\epsilon (dh^{dc} + \nabla^{(d} c^{c)}) \end{aligned} \quad (\text{B.2a})$$

---

<sup>8</sup>The first two equations might look a bit odd, but they are in fact a consequence of normalization of  $\delta(f(\sigma)) \propto \int \mathcal{D}B \exp \left( i \int d^2\sigma \sqrt{h} B(\sigma) f(\sigma) \right)$ , instead of choosing  $\delta(f(\sigma)) \propto \int \mathcal{D}B \exp \left( i \int d^2\sigma B(\sigma) f(\sigma) \right)$ .

- $\sqrt{h} = \sqrt{\text{Det}[h_{ab}]}$

$$\begin{aligned}
\delta_{\text{BRST}}\sqrt{h} &= \frac{1}{2} \frac{1}{\sqrt{h}} \delta_{\text{BRST}}(\text{Det}[h_{ab}]) \\
\delta_{\text{BRST}}\sqrt{h} &= \frac{1}{2} \frac{1}{\sqrt{h}} \delta_{\text{BRST}}(\exp(\ln(\text{Det}[h_{ab}]))) \\
\delta_{\text{BRST}}\sqrt{h} &= \frac{1}{2} \frac{1}{\sqrt{h}} \delta_{\text{BRST}}(\exp(\text{Tr}[\ln(h_{ab})])) \\
\delta_{\text{BRST}}\sqrt{h} &= \frac{1}{2} \frac{1}{\sqrt{h}} (\exp(\text{Tr}[\ln(h_{ab})])) \delta_{\text{BRST}}(\text{Tr}[\ln(h_{ab})]) \\
\delta_{\text{BRST}}\sqrt{h} &= \frac{1}{2} \frac{1}{\sqrt{h}} h \text{Tr}[\delta_{\text{BRST}}(\ln(h_{ab}))] \\
\delta_{\text{BRST}}\sqrt{h} &= \frac{1}{2} \sqrt{h} \text{Tr}[h^{ca} \delta_{\text{BRST}} h_{ab}] \\
\delta_{\text{BRST}}\sqrt{h} &= \frac{1}{2} \sqrt{h} h^{ba} \delta_{\text{BRST}} h_{ab} \\
\delta_{\text{BRST}}\sqrt{h} &= i\epsilon \sqrt{h} h^{ba} (dh_{ab} + \nabla_{(a} c_{b)}) \\
\delta_{\text{BRST}}\sqrt{h} &= i\epsilon \sqrt{h} (2d + \nabla_a c^a)
\end{aligned} \tag{B.2b}$$

- $\sqrt{h} h^{ab}$

$$\begin{aligned}
\delta_{\text{BRST}}(\sqrt{h} h^{ab}) &= \delta_{\text{BRST}}(\sqrt{h}) h^{ab} + \sqrt{h} \delta_{\text{BRST}}(h^{ab}) \\
\delta_{\text{BRST}}(\sqrt{h} h^{ab}) &= i\epsilon \sqrt{h} (2d + \nabla_c c^c) h^{ab} - 2i\epsilon \sqrt{h} (dh^{ab} + \nabla^{(a} c^{b)}) \\
\delta_{\text{BRST}}(\sqrt{h} h^{ab}) &= 2i\epsilon \sqrt{h} \left( \frac{1}{2} h^{ab} \nabla_c c^c - \nabla^{(a} c^{b)} \right)
\end{aligned} \tag{B.2c}$$

So now we can proceed to compute the two conditions, we'll start by the second one, luckily, the BRST procedure already provides a natural candidate for  $\mathcal{O}$ ,

$$\begin{aligned}
\delta_{\text{BRST}} \left( -\frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} (\hat{h}^{ab} - h^{ab}) \right) &= -\frac{1}{4\pi} \int d^2\sigma \delta_{\text{BRST}}(\sqrt{h} b_{ab}) (\hat{h}^{ab} - h^{ab}) \\
&\quad - \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} \delta_{\text{BRST}} \hat{h}^{ab} \\
&\quad + \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} \delta_{\text{BRST}}(h^{ab}) \\
\delta_{\text{BRST}} \left( -\frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} (\hat{h}^{ab} - h^{ab}) \right) &= \epsilon \frac{1}{4\pi} \int d^2\sigma \sqrt{h} B_{ab} (\hat{h}^{ab} - h^{ab}) \\
&\quad + i\epsilon \frac{1}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} (dh^{ab} + \nabla^a c^b) \\
\delta_{\text{BRST}} \left( -\frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} (\hat{h}^{ab} - h^{ab}) \right) &= i\epsilon S_{\text{gf}} + i\epsilon S_{\text{gh}} \\
\delta_{\text{BRST}} \left( -\frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} (\hat{h}^{ab} - h^{ab}) \right) &= i\epsilon (S_{\text{gf}} + S_{\text{gh}})
\end{aligned}$$

so what remains to be *checked* is the first condition, actually we have to *impose* it, let's do it part by part, starting with,

$$\delta_{\text{BRST}} S_X = \frac{1}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} \partial_a X^\mu \partial_b \delta_{\text{BRST}} X_\mu + \frac{1}{4\pi\alpha'} \int_M d^2\sigma \delta_{\text{BRST}}(\sqrt{h} h^{ab}) \partial_a X^\mu \partial_b X_\mu$$

$$\begin{aligned}
\delta_{\text{BRST}} S_X &= \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} \partial_a X^\mu \partial_b [c^c \partial_c X_\mu] + \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} \left( \frac{1}{2} h^{ab} \nabla_c c^c - \nabla^a c^b \right) \partial_a X^\mu \partial_b X_\mu \\
\delta_{\text{BRST}} S_X &= \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} \partial_a X^\mu ((\nabla_b c^c) \partial_c X_\mu + c^c \nabla_b \nabla_c X_\mu) \\
&\quad + \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} \left( \frac{1}{2} h^{ab} \nabla_c c^c - \nabla^a c^b \right) \partial_a X^\mu \partial_b X_\mu \\
\delta_{\text{BRST}} S_X &= \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} \partial_a X^\mu \partial_b X_\mu \nabla^a c^b + \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} c^c \partial_a X^\mu \nabla_c \nabla_b X_\mu \\
&\quad + \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} \left( \frac{1}{2} h^{ab} \nabla_c c^c - \nabla^a c^b \right) \partial_a X^\mu \partial_b X_\mu \\
\delta_{\text{BRST}} S_X &= \frac{i\epsilon}{2\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} c^c \partial_a X^\mu \nabla_c \partial_b X_\mu + \frac{i\epsilon}{4\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} \nabla_c c^c \partial_a X^\mu \partial_b X_\mu \\
\delta_{\text{BRST}} S_X &= \frac{i\epsilon}{4\pi\alpha'} \int_M d^2\sigma \sqrt{h} c^c \nabla_c (h^{ab} \partial_a X^\mu \partial_b X_\mu) + \frac{i\epsilon}{4\pi\alpha'} \int_M d^2\sigma \sqrt{h} h^{ab} \nabla_c c^c \partial_a X^\mu \partial_b X_\mu \\
\delta_{\text{BRST}} S_X &= \frac{i\epsilon}{4\pi\alpha'} \int_M d^2\sigma \sqrt{h} \nabla_c (c^c h^{ab} \partial_a X^\mu \partial_b X_\mu) \\
\delta_{\text{BRST}} S_X &= \frac{i\epsilon}{4\pi\alpha'} \int_M d^2\sigma \partial_c \left( \sqrt{h} c^c h^{ab} \partial_a X^\mu \partial_b X_\mu \right) \tag{B.3}
\end{aligned}$$

now,

$$\begin{aligned}
\delta_{\text{BRST}} S_{\text{gh}} &= \delta_{\text{BRST}} \left( \frac{1}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} [h^{ab} d + \nabla^a c^b] \right) \\
\delta_{\text{BRST}} S_{\text{gh}} &= \delta_{\text{BRST}} \left( \frac{1}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} \left[ h^{ab} d + \frac{1}{2} (-c^c \partial_c h^{ab} + h^{ac} \partial_c c^b + h^{bc} \partial_c c^a) \right] \right) \\
\delta_{\text{BRST}} S_{\text{gh}} &= \frac{1}{2\pi} \int d^2\sigma \delta_{\text{BRST}} \left( \sqrt{h} b_{ab} \right) [h^{ab} d + \nabla^a c^b] \\
&\quad + \frac{1}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} [\delta_{\text{BRST}}(h^{ab}) d + h^{ab} \delta_{\text{BRST}}(d)] \\
&\quad - \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [\delta_{\text{BRST}}(c^c) \partial_c h^{ab} + c^c \partial_c \delta_{\text{BRST}}(h^{ab})] \\
&\quad + \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [\delta_{\text{BRST}}(h^{ac}) \partial_c c^b + h^{ac} \partial_c \delta_{\text{BRST}}(c^b)] \\
&\quad + \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [\delta_{\text{BRST}}(h^{bc}) \partial_c c^a + h^{bc} \partial_c \delta_{\text{BRST}}(c^a)] \\
\delta_{\text{BRST}} S_{\text{gh}} &= -\frac{\epsilon}{2\pi} \int d^2\sigma \sqrt{h} B_{ab} [h^{ab} d + \nabla^a c^b] \\
&\quad + \frac{1}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} [-2i\epsilon (dh^{ab} + \nabla^a c^b) d + h^{ab} i\epsilon c^d \partial_d d] \\
&\quad - \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [i\epsilon c^d \partial_d c^c \partial_c h^{ab} + 2i\epsilon c^c \partial_c (dh^{ab} + \nabla^a c^b)] \\
&\quad + \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [-2i\epsilon (dh^{ac} + \nabla^a c^c) \partial_c c^b + i\epsilon h^{ac} \partial_c (c^d \partial_d c^b)]
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [-2i\epsilon (dh^{bc} + \nabla^{(b} c^{c)}) \partial_c c^a + i\epsilon h^{bc} \partial_c (c^d \partial_d c^a)] \\
\delta_{\text{BRST}} S_{\text{gh}} &= -\frac{i}{2\pi} \int d^2\sigma \sqrt{h} B_{ab} (-i\epsilon) [h^{ab} d + \nabla^a c^b] \\
& + \frac{i\epsilon}{\pi} \int d^2\sigma \sqrt{h} b_{ab} \nabla^{(a} c^{b)} d \\
& + \frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [c^d \partial_d c^c \partial_c h^{ab} + 2c^c (d\partial_c h^{ab} + \partial_c \nabla^{(a} c^{b)})] \\
& + \frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [2(dh^{ac} + \nabla^{(a} c^{c)}) \partial_c c^b - h^{ac} \partial_c (c^d \partial_d c^b)] \\
& + \frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [2(dh^{bc} + \nabla^{(b} c^{c)}) \partial_c c^a - h^{bc} \partial_c (c^d \partial_d c^a)] \\
\delta_{\text{BRST}} S_{\text{gh}} &= \frac{i}{2\pi} \int d^2\sigma \sqrt{h} B_{ab} \delta_{\text{BRST}} (\hat{h}^{ab} - h^{ab}) \\
& + \frac{i\epsilon}{\pi} \int d^2\sigma \sqrt{h} b_{ab} \nabla^{(a} c^{b)} d \\
& + \frac{i\epsilon}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} d [-c^c \partial_c h^{ab} + h^{ac} \partial_c c^b + h^{bc} \partial_c c^a] \\
& + \frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [c^d \partial_d c^c \partial_c h^{ab} - h^{bc} \partial_c (c^d \partial_d c^a) - h^{ac} \partial_c (c^d \partial_d c^b)] \\
& + \frac{i\epsilon}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} [\nabla^{(a} c^{c)} \partial_c c^b + c^c \partial_c \nabla^{(a} c^{b)} + \nabla^{(b} c^{c)} \partial_c c^a] \\
\delta_{\text{BRST}} S_{\text{gh}} &= -\delta_{\text{BRST}} S_{\text{gf}} \\
& + \frac{i\epsilon}{\pi} \int d^2\sigma \sqrt{h} b_{ab} \nabla^{(a} c^{b)} d \\
& + \frac{i\epsilon}{\pi} \int d^2\sigma \sqrt{h} b_{ab} d \nabla^{(a} c^{b)} \\
& - \frac{i\epsilon}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} \nabla^{[a} (c^d \nabla_d c^{b]}) \\
& + \frac{i\epsilon}{2\pi} \int d^2\sigma \sqrt{h} b_{ab} [\nabla^{(a} c^{c)} \nabla_c c^b + c^c \nabla_c \nabla^{(a} c^{b)} + \nabla^{(b} c^{c)} \nabla_c c^a] \\
\delta_{\text{BRST}} (S_{\text{gh}} + S_{\text{gf}}) &= -\frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [\nabla^a c^d \nabla_d c^b + \nabla^b c^d \nabla_d c^a + c^d \nabla^a \nabla_d c^b + c^d \nabla^b \nabla_d c^a] \\
& + \frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [\nabla^a c^c \nabla_c c^b + \nabla^c c^a \nabla_c c^b + \nabla^b c^c \nabla_c c^a + \nabla^c c^b \nabla_c c^a] \\
& + \frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [c^c \nabla_c \nabla^a c^b + c^c \nabla_c \nabla^b c^a] \\
\delta_{\text{BRST}} (S_{\text{gh}} + S_{\text{gf}}) &= -\frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [c^c \nabla^a \nabla_c c^b + c^c \nabla^b \nabla_c c^a] \\
& + \frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} [c^c \nabla_c \nabla^a c^b + c^c \nabla_c \nabla^b c^a] \\
\delta_{\text{BRST}} (S_{\text{gh}} + S_{\text{gf}}) &= -\frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} c^c [R_d{}^{ba}{}_c + R_d{}^{ab}{}_c] c^d \\
\delta_{\text{BRST}} (S_{\text{gh}} + S_{\text{gf}}) &= -\frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} c^c c^d [R_d{}^{ba}{}_c + R_d{}^{ab}{}_c] \\
\delta_{\text{BRST}} (S_{\text{gh}} + S_{\text{gf}}) &= -\frac{i\epsilon}{4\pi} \int d^2\sigma \sqrt{h} b_{ab} c^c c^d [R_d{}^{ba}{}_c - R_c{}^{ba}{}_d + R_d{}^{ab}{}_c - R_c{}^{ab}{}_d]
\end{aligned}$$

$$\begin{aligned}
\delta_{\text{BRST}}(S_{\text{gh}} + S_{\text{gf}}) &= -\frac{\text{i}\epsilon}{4\pi} \int \text{d}^2\sigma \sqrt{\hbar} b_{ab} \dot{c}^c \dot{c}^d [R_{cd}^a{}^b - R_{dc}^a{}^b + R_d{}^{ab}{}_c - R_c{}^{ab}{}_d] \\
\delta_{\text{BRST}}(S_{\text{gh}} + S_{\text{gf}}) &= -\frac{\text{i}\epsilon}{4\pi} \int \text{d}^2\sigma \sqrt{\hbar} b_{ab} \dot{c}^c \dot{c}^d [R_c{}^{ab}{}_d - R_d{}^{ab}{}_c + R_d{}^{ab}{}_c - R_c{}^{ab}{}_d] \\
\delta_{\text{BRST}}(S_{\text{gh}} + S_{\text{gf}}) &= 0
\end{aligned}$$

so that the imposition  $\delta_{\text{BRST}}(S_X + S_{\text{gh}} + S_{\text{gf}}) = 0$ , is equivalent to,

## C Closure of BRST transformations

Nilpotency of BRST,

$$\begin{aligned}
\delta_1 X^\mu &= i\epsilon_1 c^a \partial_a X^\mu \\
\delta_2 \delta_1 X^\mu &= i\epsilon_1 \delta_2(c^a) \partial_a X^\mu + i\epsilon_1 c^a \partial_a (\delta_2(X^\mu)) \\
\delta_2 \delta_1 X^\mu &= i\epsilon_1 (i\epsilon_2) c^c \nabla_c(c^a) \partial_a X^\mu + i\epsilon_1 c^a \partial_a ((i\epsilon_2 c^b \partial_b) X^\mu) \\
\delta_2 \delta_1 X^\mu &= -\epsilon_1 \epsilon_2 c^c \nabla_c(c^a) \partial_a X^\mu + \epsilon_1 \epsilon_2 c^a \partial_a (c^b \partial_b X^\mu) \\
\delta_2 \delta_1 X^\mu &= -\epsilon_1 \epsilon_2 c^c \nabla_c(c^a) \nabla_a X^\mu + \epsilon_1 \epsilon_2 c^a \nabla_a (c^b \nabla_b X^\mu) \\
\delta_2 \delta_1 X^\mu &= -\epsilon_1 \epsilon_2 c^c \nabla_c(c^a) \nabla_a X^\mu + \epsilon_1 \epsilon_2 c^a \nabla_a (c^b) \nabla_b X^\mu + \epsilon_1 \epsilon_2 c^a c^b \nabla_a \nabla_b X^\mu \\
\delta_2 \delta_1 X^\mu &= \epsilon_1 \epsilon_2 c^a c^b \nabla_a \nabla_b X^\mu, \quad \text{using the ghost statistics} \\
\delta_2 \delta_1 X^\mu &= \epsilon_1 \epsilon_2 c^a c^b \nabla_{[a} \nabla_{b]} X^\mu, \quad \text{as } X^\mu \text{ is a world-sheet scalar: } \nabla_a \nabla_b X^\mu = \nabla_b \nabla_a X^\mu \\
\delta_2 \delta_1 X^\mu &= 0
\end{aligned}$$

$$\begin{aligned}
\delta_1 h_{ab} &= 2i\epsilon_1 dh_{ab} + i\epsilon_1 (c^c \partial_c h_{ab} + h_{ac} \partial_b c^c + h_{bc} \partial_a c^c) \\
\delta_2 \delta_1 h_{ab} &= 2i\epsilon_1 \delta_2(d) h_{ab} + 2i\epsilon_1 d\delta_2(h_{ab}) \\
&\quad + i\epsilon_1 (\delta_2(c^c) \partial_c h_{ab} + c^c \partial_c \delta_2(h_{ab})) \\
&\quad + i\epsilon_1 (\delta_2(h_{ac}) \partial_b c^c + h_{ac} \partial_b \delta_2(c^c)) \\
&\quad + i\epsilon_1 (\delta_2(h_{bc}) \partial_a c^c + h_{bc} \partial_a \delta_2(c^c)) \\
\delta_2 \delta_1 h_{ab} &= 2i\epsilon_1 i\epsilon_2 h_{ab} c^c \partial_c d + 2i\epsilon_1 d 2i\epsilon_2 (dh_{ab} + \nabla_{(a} c_{b)}) \\
&\quad + i\epsilon_1 (i\epsilon_2 c^d \partial_d c^c \partial_c h_{ab} + c^c \partial_c (2i\epsilon_2 (dh_{ab} + \nabla_{(a} c_{b)}))) \\
&\quad + i\epsilon_1 (2i\epsilon_2 (dh_{ac} + \nabla_{(a} c_{c)}) \partial_b c^c + h_{ac} \partial_b (i\epsilon_2 c^d \partial_d c^c)) \\
&\quad + i\epsilon_1 (2i\epsilon_2 (dh_{bc} + \nabla_{(b} c_{c)}) \partial_a c^c + h_{bc} \partial_a (i\epsilon_2 c^d \partial_d c^c)) \\
\delta_2 \delta_1 h_{ab} &= -2\epsilon_1 \epsilon_2 h_{ab} c^c \partial_c d + 4\epsilon_1 \epsilon_2 d \nabla_{(a} c_{b)} \\
&\quad - \epsilon_1 \epsilon_2 c^d \partial_d c^c \partial_c h_{ab} + 2\epsilon_1 \epsilon_2 c^c \partial_c (dh_{ab} + \nabla_{(a} c_{b)}) \\
&\quad - 2\epsilon_1 \epsilon_2 (dh_{ac} + \nabla_{(a} c_{c)}) \partial_b c^c - \epsilon_1 \epsilon_2 h_{ac} \partial_b (c^d \partial_d c^c) \\
&\quad - 2\epsilon_1 \epsilon_2 (dh_{bc} + \nabla_{(b} c_{c)}) \partial_a c^c - \epsilon_1 \epsilon_2 h_{bc} \partial_a (c^d \partial_d c^c) \\
\delta_2 \delta_1 h_{ab} &= 4\epsilon_1 \epsilon_2 d \nabla_{(a} c_{b)} - \epsilon_1 \epsilon_2 c^d \partial_d c^c \partial_c h_{ab} + 2\epsilon_1 \epsilon_2 c^c (d \partial_c h_{ab} + \partial_c \nabla_{(a} c_{b)}) \\
&\quad - 2\epsilon_1 \epsilon_2 (dh_{ac} + \nabla_{(a} c_{c)}) \partial_b c^c - \epsilon_1 \epsilon_2 h_{ac} \partial_b (c^d \partial_d c^c) \\
&\quad - 2\epsilon_1 \epsilon_2 (dh_{bc} + \nabla_{(b} c_{c)}) \partial_a c^c - \epsilon_1 \epsilon_2 h_{bc} \partial_a (c^d \partial_d c^c) \\
\delta_2 \delta_1 h_{ab} &= 4\epsilon_1 \epsilon_2 d \nabla_{(a} c_{b)} - 2\epsilon_1 \epsilon_2 d c^c \partial_c h_{ab} - 2\epsilon_1 \epsilon_2 d h_{ac} \partial_b c^c - 2\epsilon_1 \epsilon_2 d h_{bc} \partial_a c^c \\
&\quad - \epsilon_1 \epsilon_2 c^d \partial_d c^c \partial_c h_{ab} - \epsilon_1 \epsilon_2 h_{bc} \partial_a (c^d \partial_d c^c) - \epsilon_1 \epsilon_2 h_{ac} \partial_b (c^d \partial_d c^c) \\
&\quad - 2\epsilon_1 \epsilon_2 \partial_c \nabla_{(a} c_{b)} c^c - 2\epsilon_1 \epsilon_2 \nabla_{(a} c_{c)} \partial_b c^c - 2\epsilon_1 \epsilon_2 \nabla_{(b} c_{c)} \partial_a c^c \\
\delta_2 \delta_1 h_{ab} &= 4\epsilon_1 \epsilon_2 d \nabla_{(a} c_{b)} - 2\epsilon_1 \epsilon_2 d [c^c \partial_c h_{ab} + h_{ac} \partial_b c^c + h_{bc} \partial_a c^c] \\
&\quad - \epsilon_1 \epsilon_2 [c^d \partial_d c^c \partial_c h_{ab} + h_{bc} \partial_a (c^d \partial_d c^c) + h_{ac} \partial_b (c^d \partial_d c^c)] \\
&\quad - 2\epsilon_1 \epsilon_2 [\partial_c \nabla_{(a} c_{b)} c^c + \nabla_{(a} c_{c)} \partial_b c^c + \nabla_{(b} c_{c)} \partial_a c^c] \\
\delta_2 \delta_1 h_{ab} &= 4\epsilon_1 \epsilon_2 d \nabla_{(a} c_{b)} - 4\epsilon_1 \epsilon_2 d \nabla_{(a} c_{b)} \\
&\quad - 2\epsilon_1 \epsilon_2 \nabla_{(a} [c^c \nabla_{c} c_{b)}] \\
&\quad - 2\epsilon_1 \epsilon_2 [\nabla_c \nabla_{(a} c_{b)} c^c + \nabla_{(a} c_{c)} \nabla_b c^c + \nabla_{(b} c_{c)} \nabla_a c^c] \\
\delta_2 \delta_1 h_{ab} &= -2\epsilon_1 \epsilon_2 [\nabla_{(a} [c^c \nabla_{c} c_{b)}] + \nabla_c \nabla_{(a} c_{b)} c^c + \nabla_{(a} c_{c)} \nabla_b c^c + \nabla_{(b} c_{c)} \nabla_a c^c]
\end{aligned}$$

$$\begin{aligned}
\delta_2 \delta_1 h_{ab} &= -\epsilon_1 \epsilon_2 [\nabla_a (c^c \nabla_c c_b) + \nabla_b (c^c \nabla_c c_a) + \nabla_c \nabla_a c_b c^c + \nabla_c \nabla_b c_a c^c \\
&\quad + \nabla_a c_c \nabla_b c^c + \nabla_b c_c \nabla_a c^c + \nabla_c c_a \nabla_b c^c + \nabla_c c_b \nabla_a c^c] \\
\delta_2 \delta_1 h_{ab} &= -\epsilon_1 \epsilon_2 [c^c \nabla_a \nabla_c c_b + c^c \nabla_b \nabla_c c_a + \nabla_c \nabla_a c_b c^c + \nabla_c \nabla_b c_a c^c \\
&\quad + \nabla_a c^c \nabla_c c_b + \nabla_b c^c \nabla_c c_a + \nabla_a c_c \nabla_b c^c + \nabla_b c_c \nabla_a c^c + \nabla_c c_a \nabla_b c^c + \nabla_c c_b \nabla_a c^c] \\
\delta_2 \delta_1 h_{ab} &= -\epsilon_1 \epsilon_2 [c^c \nabla_a \nabla_c c_b + c^c \nabla_b \nabla_c c_a - c^c \nabla_c \nabla_a c_b - c^c \nabla_c \nabla_b c_a \\
&\quad + \nabla_a c^c \nabla_c c_b + \nabla_b c^c \nabla_c c_a - \nabla_b c^c \nabla_c c_a - \nabla_a c^c \nabla_c c_b] \\
\delta_2 \delta_1 h_{ab} &= -\epsilon_1 \epsilon_2 [c^c \nabla_a \nabla_c c_b + c^c \nabla_b \nabla_c c_a - c^c \nabla_c \nabla_a c_b - c^c \nabla_c \nabla_b c_a \\
&\quad + \nabla_a c^c \nabla_c c_b + \nabla_b c^c \nabla_c c_a - \nabla_b c^c \nabla_c c_a - \nabla_a c^c \nabla_c c_b] \\
\delta_2 \delta_1 h_{ab} &= -\epsilon_1 \epsilon_2 c^c [R^d_{bac} c_d + R^d_{abc} c_d] \\
\delta_2 \delta_1 h_{ab} &= -\epsilon_1 \epsilon_2 c^c c^d [R_{dbac} + R_{dabc}] \\
\delta_2 \delta_1 h_{ab} &= -\frac{1}{2} \epsilon_1 \epsilon_2 c^c c^d [R_{dbac} - R_{cbad} + R_{dabc} - R_{cabd}] \\
\delta_2 \delta_1 h_{ab} &= -\frac{1}{2} \epsilon_1 \epsilon_2 c^c c^d [R_{dbac} - R_{cbad} + R_{bcd a} - R_{bdca}] \\
\delta_2 \delta_1 h_{ab} &= -\frac{1}{2} \epsilon_1 \epsilon_2 c^c c^d [R_{dbac} - R_{cbad} + R_{cbad} - R_{dbac}] \\
\delta_2 \delta_1 h_{ab} &= 0
\end{aligned}$$

$$\begin{aligned}
\delta_2 \Gamma_{be}^d &= \delta_2 \left\{ \frac{1}{2} h^{df} (\partial_b h_{ef} + \partial_e h_{bf} - \partial_f h_{be}) \right\} \\
\delta_2 \Gamma_{be}^d &= -i\epsilon_2 (dh^{df} + \nabla^{(d} c^{f)}) (\partial_b h_{ef} + \partial_e h_{bf} - \partial_f h_{be}) \\
&\quad + i\epsilon_2 h^{df} (\partial_b [dh_{ef} + \nabla_{(e} c_{f)}] + \partial_e [dh_{bf} + \nabla_{(b} c_{f)}] - \partial_f [dh_{be} + \nabla_{(b} c_{e)}]) \\
\delta_2 \Gamma_{be}^d &= -i\epsilon_2 \nabla^{(d} c^{f)} (\partial_b h_{ef} + \partial_e h_{bf} - \partial_f h_{be}) \\
&\quad + i\epsilon_2 h^{df} (h_{ef} \partial_b d + \partial_b \nabla_{(e} c_{f)} + h_{bf} \partial_e d + \partial_e \nabla_{(b} c_{f)} - h_{be} \partial_f d - \partial_f \nabla_{(b} c_{e)}) \\
h_{da} \delta_2 \Gamma_{be}^d &= -i\epsilon_2 h_{da} \nabla^{(d} c^{f)} (\partial_b h_{ef} + \partial_e h_{bf} - \partial_f h_{be}) \\
&\quad + i\epsilon_2 h_{da} h^{df} (h_{ef} \partial_b d + \partial_b \nabla_{(e} c_{f)} + h_{bf} \partial_e d + \partial_e \nabla_{(b} c_{f)} - h_{be} \partial_f d - \partial_f \nabla_{(b} c_{e)}) \\
h_{da} \delta_2 \Gamma_{be}^d &= -i\epsilon_2 h_{da} \nabla^{(d} c^{f)} (\partial_b h_{ef} + \partial_e h_{bf} - \partial_f h_{be}) \\
&\quad + i\epsilon_2 (h_{ea} \partial_b d + \partial_b \nabla_{(a} c_{e)} + h_{ba} \partial_e d + \partial_e \nabla_{(b} c_{a)} - h_{eb} \partial_a d - \partial_a \nabla_{(b} c_{e)}) \\
h_{d(a} \delta_2 \Gamma_{b)e}^d &= -i\epsilon_2 h_{d(a} \nabla^{(d} c^{f)} (\partial_b) h_{ef} + \partial_e h_{bf} - \partial_f h_{be}) \\
&\quad + i\epsilon_2 (h_{ba} \partial_e d + \partial_e \nabla_{(b} c_{a)}) \\
h_{d(a} \delta_2 \Gamma_{b)e}^d &= -i\epsilon_2 \nabla_{(c} c_{(a)} h^{fc} (\partial_b) h_{ef} + \partial_e h_{bf} - \partial_f h_{be}) \\
&\quad + i\epsilon_2 (h_{ba} \partial_e d + \partial_e \nabla_{(b} c_{a)}) \\
h_{d(a} \delta_2 \Gamma_{b)e}^d &= -2i\epsilon_2 \nabla_{(c} c_{(a)} \Gamma_{b)e}^c + i\epsilon_2 (h_{ba} \partial_e d + \partial_e \nabla_{(b} c_{a)}) \\
h_{d(a} \delta_2 \Gamma_{b)e}^d &= -i\epsilon_2 [\nabla_c c_{(a} + \nabla_{(a} c_{c)}] \Gamma_{b)e}^c + i\epsilon_2 (h_{ba} \partial_e d + \partial_e \nabla_{(b} c_{a)}) \\
h_{d(a} \delta_2 \Gamma_{b)e}^d &= -\frac{1}{2} i\epsilon_2 [\nabla_c c_a + \nabla_a c_c] \Gamma_{be}^c - \frac{1}{2} i\epsilon_2 [\nabla_c c_b + \nabla_b c_c] \Gamma_{ae}^c + \frac{1}{2} i\epsilon_2 \partial_e \nabla_b c_a + \frac{1}{2} i\epsilon_2 \partial_e \nabla_a c_b \\
&\quad + i\epsilon_2 h_{ba} \partial_e d \\
h_{d(a} \delta_2 \Gamma_{b)e}^d &= \frac{1}{2} i\epsilon_2 [\partial_e \nabla_a c_b - \Gamma_{be}^c \nabla_a c_c - \Gamma_{ae}^c \nabla_c c_b] + \frac{1}{2} i\epsilon_2 [\partial_e \nabla_b c_a - \Gamma_{be}^c \nabla_c c_a - \Gamma_{ae}^c \nabla_b c_c] \\
&\quad + i\epsilon_2 h_{ba} \partial_e d \\
h_{d(a} \delta_2 \Gamma_{b)e}^d &= \frac{1}{2} i\epsilon_2 \nabla_e \nabla_a c_b + \frac{1}{2} i\epsilon_2 \nabla_e \nabla_b c_a + i\epsilon_2 h_{ba} \partial_e d
\end{aligned}$$



$$h_{d(a}\delta_2\Gamma_{b)e}^d = \mathrm{i}\epsilon_2\nabla_e\nabla_{(a}c_{b)} + \mathrm{i}\epsilon_2h_{ba}\partial_e d$$