ADTGR SEMINAR

VICENTE V. FIGUEIRA

1. Introduction

Einstein-Hilbert action:

$$S_{\rm EH} = \frac{1}{\kappa^2} \int\limits_{M} \mathrm{d}^4 x \, \sqrt{|g|} g^{ab} R^c_{\ acb}$$

The Vierbein/Tetrad formalism:

$$\eta_{\mu\nu} = \mathbf{g}(\mathbf{e}_{\mu}, \mathbf{e}_{\nu})$$
$$g_{ab} = e^{\mu}{}_{a}e^{\nu}{}_{b}\eta_{\mu\nu}$$
$$\eta_{\mu\nu} = e_{\mu}{}^{a}e_{\nu}{}^{b}g_{ab}$$

A connection is defined with respect to a vector basis:

$$\nabla_{\mathbf{X}}(\mathbf{e}_{\nu}) = \boldsymbol{\omega}(\mathbf{X})^{\mu}_{\nu}\mathbf{e}_{\mu}$$

$$\nabla_{a}e_{\nu}^{b} = \omega_{a}^{\mu}_{\nu}e_{\mu}^{b}$$

$$e_{\alpha}^{c}g_{cb}\nabla_{a}e_{\nu}^{b} = e_{\alpha}^{c}g_{cb}\omega_{a}^{\mu}_{\nu}e_{\mu}^{b}$$

$$e_{\alpha}^{c}g_{cb}\nabla_{a}e_{\nu}^{b} + e_{\alpha}^{c}e_{\nu}^{b}\nabla_{a}g_{cb} = \omega_{a\alpha\nu}$$

$$e_{\alpha}^{c}\nabla_{a}(g_{cb}e_{\nu}^{b}) = \omega_{a\alpha\nu}$$

$$\nabla_{a}(e_{\alpha}^{c}g_{cb}e_{\nu}^{b}) - g_{cb}e_{\nu}^{b}\nabla_{a}e_{\alpha}^{c} = \omega_{a\alpha\nu}$$

$$\nabla_{a}(\eta_{\alpha\nu}) - g_{cb}e_{\nu}^{b}\nabla_{a}e_{\alpha}^{c} = \omega_{a\alpha\nu}$$

$$-g_{cb}e_{\nu}^{b}\nabla_{a}e_{\alpha}^{c} = \omega_{a\alpha\nu}$$

$$-\omega_{a\nu\alpha} = \omega_{a\alpha\nu}, \quad \text{Metric compatibility}$$

Riemann curvature tensor, $\mathbf{Riem}(\mathbf{X}, \mathbf{Y}) : \mathfrak{X} \to \mathfrak{X}$:

$$\begin{split} \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= \left(\nabla_{\mathbf{X}}\nabla_{\mathbf{Y}} - \nabla_{\mathbf{Y}}\nabla_{\mathbf{X}} - \nabla_{[\mathbf{X},\mathbf{Y}]}\right)\mathbf{e}_{\mu} \\ \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= \left(\nabla_{\mathbf{X}}\left(Y^{b}\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) - \nabla_{\mathbf{Y}}\left(X^{a}\omega_{a}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) - [\mathbf{X},\mathbf{Y}]^{b}\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) \\ \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= \left(\nabla_{\mathbf{X}}\left(Y^{b}\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) - \nabla_{\mathbf{Y}}\left(X^{a}\omega_{a}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) - [\mathbf{X},\mathbf{Y}]^{b}\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) \\ \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= \left(Y^{b}\nabla_{\mathbf{X}}\left(\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) - X^{a}\nabla_{\mathbf{Y}}\left(\omega_{a}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) \\ + \nabla_{\mathbf{X}}\left(Y^{b}\right)\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu} - \nabla_{\mathbf{Y}}\left(X^{b}\right)\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu} - [\mathbf{X},\mathbf{Y}]^{b}\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) \\ \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= \left(X^{a}Y^{b}\nabla_{a}\left(\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) - X^{a}Y^{b}\nabla_{b}\left(\omega_{a}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) \\ + \left(\nabla_{\mathbf{X}}\left(Y^{b}\right) - \nabla_{\mathbf{Y}}\left(X^{b}\right)\right)\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu} - [\mathbf{X},\mathbf{Y}]^{b}\omega_{b}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu}\right) \\ \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= \left(X^{a}Y^{b}\nabla_{a}\left(\omega_{b}{}^{\nu}{}_{\mu}\right)\mathbf{e}_{\nu} + X^{a}Y^{b}\omega_{b}{}^{\nu}{}_{\mu}\omega_{a}{}^{\nu}{}_{\nu}\mathbf{e}_{\alpha} - X^{a}Y^{b}\nabla_{b}\left(\omega_{a}{}^{\nu}{}_{\mu}\right)\mathbf{e}_{\nu} - X^{a}Y^{b}\omega_{a}{}^{\nu}{}_{\mu}\omega_{a}\right) \\ \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= X^{a}Y^{b}\left(\nabla_{a}\left(\omega_{b}{}^{\nu}{}_{\mu}\right) + \omega_{b}{}^{\alpha}{}_{\mu}\omega_{a}{}^{\alpha}{}_{\alpha} - \nabla_{b}\left(\omega_{a}{}^{\nu}{}_{\mu}\right) - \omega_{a}{}^{\alpha}{}_{\mu}\omega_{b}{}^{\nu}{}_{\alpha}\right) \mathbf{e}_{\nu} \\ \operatorname{Riem}(\mathbf{X},\mathbf{Y})\mathbf{e}_{\mu} &= X^{a}Y^{b}R_{ab}{}^{\nu}{}_{\mu}\mathbf{e}_{\nu} \end{aligned}$$

Date: June 10, 2025.

(1.1)

References

- [1] Sidney Coleman and Jeffrey Mandula. "All Possible Symmetries of the S Matrix". In: Phys. Rev. 159 (5 1967), pp. 1251-1256. DOI: 10.1103/PhysRev.159.1251. URL: https://link.aps.org/doi/10.1103/PhysRev.159.1251.
- [2] Rudolf Haag, Jan T. Lopuszański, and Martin Sohnius. "All possible generators of supersymmetries of the S-matrix". In: Nuclear Physics B 88.2 (1975), pp. 257-274. ISSN: 0550-3213. DOI: https://doi.org/10.1016/0550-3213(75)90279-5. URL: https://www.sciencedirect.com/science/article/pii/0550321375902795.
- [3] H.J.W. Müller-Kirsten and A. Wiedemann. *Introduction to Supersymmetry*. G Reference, Information and Interdisciplinary Subjects Series. World Scientific, 2010. ISBN: 9789814293426. URL: https://books.google.com.br/books?id=65DkngEACAAJ.
- [4] M. Srednicki. Quantum Field Theory. Cambridge University Press, 2007. ISBN: 9781139462761. URL: https://books.google.com.br/books?id=50epxIG42B4C.
- [5] David Tong. Supersymmetric Field Theory. 2022. URL: http://www.damtp.cam.ac.uk/user/tong/susy.html (visited on 12/10/2024).
- [6] S. Weinberg. The Quantum Theory of Fields: Volume 3, Supersymmetry. Cambridge University Press, 2005. ISBN: 9781139643436. URL: https://books.google.com.br/books?id=QMkgAwAAQBAJ.
- [7] Edward Witten. "Constraints on supersymmetry breaking". In: Nuclear Physics B 202.2 (1982), pp. 253-316. ISSN: 0550-3213. DOI: https://doi.org/10.1016/0550-3213(82)90071-2. URL: https://www.sciencedirect.com/science/article/pii/0550321382900712.