

Q1- The elbow joint connects the arm with the:

[Difficulty: Easy] — General anatomy

A- Hand

B- Forearm

C- Shoulder

D- Wrist

E- Scapula

Answer: B

Q2- The elbow joint is classified as a:

[Difficulty: Easy] — Type of joint

A- Fibrous joint

B- Cartilaginous joint

C- Synovial hinge joint

D- Saddle joint

E- Pivot joint

Answer: C

Q3- Which bones form the elbow joint?

[Difficulty: Easy] — General anatomy

A- Humerus, radius, ulna

B- Radius, ulna, scapula

C- Humerus, clavicle, ulna

D- Radius, carpal bones, ulna

E- Humerus, femur, ulna

Answer: A

Q4- The humeroulnar joint is formed between the trochlea of humerus and:

[Difficulty: Easy] — Articular surfaces

A- Radial notch

B- Capitulum

C- Olecranon

D- Trochlear notch of ulna

E- Coronoid fossa

Answer: D

Q5- The radiohumeral joint is formed between the capitulum and:
[Difficulty: Easy] — Articular surfaces

- A- Ulna**
- B- Olecranon**
- C- Radial head**
- D- Coronoid process**
- E- Radial notch**

Answer: C

Q6- In full extension, the olecranon process is received by the:
[Difficulty: Easy] — Flexion and extension relations

- A- Coronoid fossa**
- B- Radial fossa**
- C- Olecranon fossa**
- D- Trochlear notch**
- E- Capitulum**

Answer: C

Q7- The annular ligament surrounds the:
[Difficulty: Easy] — Ligaments

- A- Trochlea**
- B- Capitulum**
- C- Radial head**
- D- Coronoid process**
- E- Olecranon**

Answer: C

Q8- The anterior and posterior fat pads become visible on:
[Difficulty: Easy] — Plain radiographs

- A- AP radiograph**
- B- Oblique radiograph**
- C- Lateral radiograph**
- D- MRI**
- E- CT scan**

Answer: C

Q9- Arthrography of the elbow joint is achieved by injection of contrast medium between the radial head and:

[Difficulty: Easy] — Arthrography

- A- Trochlea**
- B- Coronoid process**
- C- Olecranon**
- D- Capitulum**
- E- Medial epicondyle**

Answer: D

Q10- The ulnar collateral ligament is responsible for:

[Difficulty: Easy] — MRI anatomy

- A- Lateral stability**
- B- Medial elbow stability**
- C- Wrist stability**
- D- Finger movement**
- E- Shoulder stability**

Answer: B

Q11- The anterior bundle of the ulnar collateral ligament is best visualized in:

[Difficulty: Easy] — MRI anatomy

- A- Sagittal images**
- B- Coronal images**
- C- Axial images**
- D- Oblique images**
- E- AP images**

Answer: B

Q12- Injury to the annular ligament should be suspected in patients with:

[Difficulty: Easy] — Clinical application

- A- Femoral dislocation**
- B- Shoulder fracture**
- C- Radial head subluxation**
- D- Clavicular injury**
- E- Scaphoid fracture**

Answer: C

Q13- The elbow joint cavity includes all the following joints EXCEPT:

[Difficulty: Medium] — Components of elbow joint

- A- Humeroulnar joint**
- B- Humeroradial joint**
- C- Superior radioulnar joint**
- D- Inferior radioulnar joint**
- E- Radiohumeral joint**

Answer: D

Q14- Which structure is received by the coronoid fossa during full flexion?

[Difficulty: Medium] — Flexion and extension

- A- Olecranon process**
- B- Radial head**
- C- Coronoid process**
- D- Capitulum**
- E- Trochlea**

Answer: C

Q15- The radial head is received by which fossa during full flexion?

[Difficulty: Medium] — Flexion and

extension

- A- Olecranon fossa**
- B- Coronoid fossa**
- C- Radial fossa**
- D- Trochlear fossa**
- E- Supratrochlear fossa**

Answer: C

**Q16- The joint capsule is attached proximally on the humerus to include:
[Difficulty: Medium] — Joint capsule**

- A- Radial tuberosity only**
- B- Olecranon, coronoid, and radial fossae**
- C- Capitulum only**
- D- Trochlea only**
- E- Coracoid process**

Answer: B

**Q17- The radial and ulnar collateral ligaments are:
[Difficulty: Medium] — Ligaments**

- A- Separate tendons**
- B- Intra-articular discs**
- C- Lateral and medial thickenings of the capsule**
- D- Synovial folds**
- E- Cartilaginous structures**

Answer: C

**Q18- The annular ligament is attached to the:
[Difficulty: Medium] — Ligaments**

- A- Medial epicondyle**
- B- Olecranon process**
- C- Anterior and posterior edges of radial notch**

D- Capitulum

E- Trochlea

Answer: C

Q19- Elbow joint effusion causes:

[Difficulty: Medium] — Plain radiographs

A- Narrowing of ulna

B- Distension of the capsule anteriorly and posteriorly

C- Capitular erosion

D- Radial shortening

E- Wrist swelling

Answer: B

Q20- Fat pads act as:

[Difficulty: Medium] — Plain radiographs

A- Bone stabilizers

B- Shock absorbers

C- Ossification centers

D- Synovial membranes

E- Ligament substitutes

Answer: B

Q21- On lateral elbow radiographs, displaced fat pads appear as:

[Difficulty: Medium] — Plain radiographs

A- Dense calcifications

B- Linear lucencies

C- Radiopaque shadows

D- Metallic densities

E- Circular opacities

Answer: B

Q22- Arthrography outlines the:

[Difficulty: Medium] — Arthrography

- A- Bone marrow cavity**
- B- Synovial cavity**
- C- Medullary canal**
- D- Olecranon process**
- E- Radial tuberosity**

Answer: B

Q23- Arthrography of the elbow is most useful in demonstrating:

[Difficulty: Medium] — Arthrography

- A- Bone tumors**
- B- Lung lesions**
- C- Intra-articular loose bodies**
- D- Vertebral fractures**
- E- Pelvic masses**

Answer: C

Q24- Arthrography helps evaluate:

[Difficulty: Medium] — Arthrography

- A- Muscle thickness**
- B- Articular cartilaginous surfaces**
- C- Skin folds**
- D- Nail bed injuries**
- E- Tendon calcification only**

Answer: B

Q25- The ulnar collateral ligament is composed of:

[Difficulty: Medium] — MRI anatomy

- A- Two bundles**
- B- Three bundles**
- C- Four bundles**
- D- One bundle**
- E- Five bundles**

Answer: B

Q26- Which bundle contributes most to elbow stability?

[Difficulty: Medium] — MRI anatomy

- A- Posterior bundle**
- B- Transverse bundle**
- C- Anterior bundle**
- D- Radial bundle**
- E- Oblique bundle**

Answer: C

Q27- The anterior bundle of the UCL becomes taut during:

[Difficulty: Medium] — MRI anatomy

- A- Flexion**
- B- Extension**
- C- Supination**
- D- Pronation**
- E- Rotation**

Answer: B

Q28- The anterior bundle extends from the undersurface of the medial epicondyle to:

[Difficulty: Medium] — MRI anatomy

- A- Lateral capitulum**
- B- Olecranon tip**
- C- Medial aspect of ulnar coronoid process**
- D- Radial tuberosity**
- E- Trochlea**

Answer: C

Q29- The posterior and transverse bundles are identified best in:

[Difficulty: Medium] — MRI anatomy

- A- Coronal plane**
- B- Sagittal plane**
- C- Axial plane**

D- AP plane

E- Oblique plane

Answer: C

Q30- Injury to the lateral radial collateral ligament may complicate:

[Difficulty: Medium] — Clinical application

A- Acute varus strain

B- Femoral fracture

C- Tibial dislocation

D- Shoulder impingement

E- Wrist sprain

Answer: A

Q31- The radial collateral ligament appears on MRI as:

[Difficulty: Medium] — MRI anatomy

A- High-signal heterogeneous structure

B- Homogeneously low-signal structure

C- Calcified opacity

D- Fat-density band

E- Fluid collection

Answer: B

Q32- The radial collateral ligament courses at the:

[Difficulty: Medium] — MRI anatomy

A- Posteromedial humerus

B- Anterolateral aspect of humerus

C- Inferior ulna

D- Posterior radius

E- Medial trochlea

Answer: B

Q33- The annular ligament is best visualized in MRI on:

[Difficulty: Medium] — MRI anatomy

- A- Coronal plane**
- B- Sagittal plane**
- C- Axial plane**
- D- AP plane**
- E- Oblique plane**

Answer: C

Q34- Which ligament surrounds the radial head and rotates within it?

[Difficulty: Medium] — Ligaments

- A- Radial collateral ligament**
- B- Ulnar collateral ligament**
- C- Annular ligament**
- D- Cruciate ligament**
- E- Coracohumeral ligament**

Answer: C

Q35- All of the following are components of the elbow joint cavity EXCEPT:

[Difficulty: Hard] — Components

- A- Humeroulnar joint**
- B- Superior radioulnar joint**
- C- Humeroradial joint**
- D- Inferior radioulnar joint**
- E- Radiohumeral joint**

Answer: D

Q36- In a lateral elbow radiograph showing posterior displacement of fat pads, the most likely cause is:

[Difficulty: Hard] — Radiological reasoning

- A- Muscle hypertrophy**
- B- Joint effusion**
- C- Osteoporosis**
- D- Tendinitis**

E- Bone infarction

Answer: B

Q37- All of the following are true regarding fat pads EXCEPT:

[Difficulty: Hard] — Plain radiographs

A- Located within the joint capsule

B- Act as shock absorbers

C- Become displaced by effusion

D- Best seen on AP radiographs

E- Separated from bone by soft tissue density

Answer: D

Q38- Which of the following correctly matches the articulation?

[Difficulty: Hard] — Articular surfaces

A- Capitulum + trochlear notch

B- Trochlea + radial head

C- Radial head + radial notch of ulna

D- Olecranon + capitulum

E- Coronoid process + trochlea

Answer: C

Q39- During full flexion of the elbow:

[Difficulty: Hard] — Flexion and extension

A- Olecranon enters radial fossa

B- Coronoid process enters coronoid fossa

C- Radial head enters olecranon fossa

D- Trochlea enters radial notch

E- Capitulum enters coronoid fossa

Answer: B

Q40- During full extension, which process is received posteriorly?

[Difficulty: Hard] — Flexion and extension

- A- Coronoid process**
- B- Radial head**
- C- Olecranon process**
- D- Styloid process**
- E- Coracoid process**

Answer: C

Q41- Which statement regarding the joint capsule is CORRECT?

[Difficulty: Hard] — Joint capsule

- A- Attached distally only to humerus**
- B- Excludes the fossae from the joint**
- C- Attached to margins of radius and ulna articular surfaces**
- D- Covers only anterior humerus**
- E- Does not include radial fossa**

Answer: C

Q42- Which ligament contributes to medial elbow stability and resists valgus strain?

[Difficulty: Hard] — MRI anatomy

- A- Radial collateral ligament**
- B- Annular ligament**
- C- Ulnar collateral ligament**
- D- Coracoacromial ligament**
- E- Interosseous ligament**

Answer: C

Q43- The most frequently injured bundle of the UCL is:

[Difficulty: Hard] — MRI anatomy

- A- Posterior bundle**
- B- Transverse bundle**
- C- Anterior bundle**
- D- Radial bundle**
- E- Oblique bundle**

Answer: C

Q44- Which MRI plane best demonstrates the posterior and transverse bundles?

[Difficulty: Hard] — MRI anatomy

A- Coronal

B- Sagittal

C- Axial

D- AP

E- Lateral

Answer: C

Q45- The radial collateral ligament injury occurs less frequently than injury to:

[Difficulty: Hard] — MRI anatomy

A- Annular ligament

B- Medial ulnar collateral ligament

C- Coracohumeral ligament

D- Patellar ligament

E- Deltoid ligament

Answer: B

Q46- A patient with repeated radial head dislocation most likely has injury to:

[Difficulty: Hard] — Clinical application

A- Ulnar collateral ligament

B- Radial collateral ligament

C- Annular ligament

D- Cruciate ligament

E- Glenoid labrum

Answer: C

Q47- Arthrography is performed infrequently but is valuable in:

[Difficulty: Hard] — Arthrography

- A- Evaluating lung fields**
- B- Detecting intra-articular loose bodies**
- C- Diagnosing skull fractures**
- D- Imaging vertebral discs**
- E- Assessing renal stones**

Answer: B

Q48- In arthrography, the synovial cavity extends:

[Difficulty: Hard] — Arthrography

- A- Distal to the ulna only**
- B- Proximal to humeral articular surfaces**
- C- Into the wrist joint**
- D- To the clavicle**
- E- Beyond the shoulder joint**

Answer: B

Q49- Which imaging modality best visualizes the annular ligament in the axial plane?

[Difficulty: Hard] — MRI anatomy

- A- Plain radiography**
- B- Ultrasound**
- C- MRI**
- D- Arthrography**
- E- Fluoroscopy**

Answer: C

Q50- All the following statements about the anterior bundle are true EXCEPT:

[Difficulty: Hard] — MRI anatomy

- A- Most important for stability**
- B- Frequently injured**
- C- Best seen in coronal images**
- D- Loose during elbow extension**
- E- Extends to coronoid process of ulna**

Answer: D
