

# Circular Motion and Gravity

## Orbital Simulation

### PHYS 442

Dr. Schultz

October 7, 2015

Date Performed: September 18, 2015  
Partners: Whole class  
Instructor: Me

## 1 Objective

Explored the motion of a particle under the influence of a gravitational force. Specifically we look at attractive inverse square distance forces, Hookean forces, escape velocity, circular orbits, kinetic energy, potential energy and elliptical orbits. These are defined in 1.1:

### 1.1 Definitions

**Law of Universal Gravitation** The law of universal gravitation states the force of gravity between two point masses is directly proportional to each mass and inversely proportional to the distance between them. This is also true for masses outside of spherically symmetric mass distributions. (Homer, 2014)

$$F_g = \frac{mMG}{r^2}$$

**Hookean Forces** Inside a uniformly dense sphere of mass the force is Hookean, with an attractive force proportional to the displacement from equilibrium. The effective spring constant is  $K = \frac{mMG}{R^3}$ .

$$F_g = \frac{mMG}{R^3}r$$

**Gravitational Constant** The universal gravitation constant  $G$  determines the strength of the gravity force from a given mass. It may also be considered

as the force that 1 kg exerts on another 1 kg mass separated by 1 meter.

$$G = 6.67 \times 10^{-11} \frac{Nm^2}{kg^2}$$

**Escape Velocity** Escape velocity is the initial velocity required to escape gravitational attraction. An object launched at the escape velocity will never come back (escape).

$$v_{escape} = \sqrt{\frac{2MG}{r}}$$

**Kinetic Energy** Kinetic energy is the energy associated with motion.

$$KE = \frac{mv^2}{2}$$

**Potential Energy** The potential associated with the universal gravitation force is written as follows.

$$PE = -\frac{mMG}{r}$$

**Circular Orbit** A circular orbit is an orbit with a constant radius  $r$ .

**Elliptic Orbit** An elliptic orbit is a closed orbit with changing radius  $r$ .

## 2 Simulation

The simulation applies a central acceleration to the orbiting particle. Outside the boundary of the central mass we have the following acceleration.

$$a = \frac{K}{r^2}$$

Inside the boundary of the central mass ( $r < R$ ) we have the following acceleration.

$$a = \frac{K}{R^3}r$$

Here  $R$  is the radius of the central mass and  $K$  is a constant determined by the user of the simulation.  $K$  is  $MG$ . For this simulation the radius was set to  $R = 6$  and the constant was set to  $K = -0.1$  making the acceleration attractive.

The initial position  $\vec{r}_0$  and initial velocity  $\vec{v}_0$  are set by the user.

### 3 Sample Calculation

#### 3.1 Circular Orbit

Given  $\vec{r}_0 = (10, 0)$  and  $K = -0.1$  we find the  $\vec{v}_0$  for circular orbit.

$$\begin{aligned}F_{net} &= ma \\ \frac{mMG}{r^2} &= m \frac{v^2}{r} \\ \frac{K}{r^2} &= \frac{v^2}{r} \\ v &= \sqrt{\frac{K}{r}} \\ v &= \sqrt{\frac{0.1}{10}} = 0.1\end{aligned}$$

The velocity must be tangential and therefore  $\vec{v}_0$  must be perpendicular to  $\vec{r}_0$ .

$$\vec{v}_0 = (0, 0.1)$$

#### 3.2 Escape Velocity

Given  $\vec{r}_0 = (10, 0)$  and  $K = -0.1$  we find the  $\vec{v}_0$  for escape from the central mass' gravitational attraction. Escape is associated with a total mechanical energy of zero.

$$\begin{aligned}PE + KE &= 0 \\ -\frac{mMG}{r} + \frac{mv^2}{2} &= 0 \\ -\frac{K}{r} + \frac{v^2}{2} &= 0 \\ v_{escape} &= \sqrt{\frac{2K}{r}} \\ v_{escape} &= \sqrt{\frac{2(0.1)}{10}} = 0.14\end{aligned}$$

## 4 Results and Conclusions

### 4.1 Circular Orbit

For the conditions  $\vec{r}_0 = (10, 0)$  and  $K = -0.1$  we calculate an initial velocity  $\vec{v}_0 = (0, 0.1)$  will give a circular orbit. Running the simulation yields the following orbit. We can see it is circular.




Figure 1: Circular Orbit

### 4.2 Escape Velocity

For the conditions  $\vec{r}_0 = (10, 0)$  and  $K = -0.1$  we calculate an initial velocity  $\vec{v}_0 = (0, 0.14)$  will give an escape from the gravitational attraction. We can observe in the simulation the object indeed escapes and the total energy is zero. See the following figure.



Figure 2: Escape Velocity

### 4.3 Elliptical Orbit

For the conditions  $\vec{r}_0 = (10, 0)$  and  $K = -0.1$  we use an initial velocity  $\vec{v}_0 = (0, 0.12)$  that is between the one for circular orbit and escape. As the figure shows the resulting orbit is elliptical.




Figure 3: Elliptic Orbit

## 5 Discussion

I did change some section in this file. After this work, I have more knowledge about technology also about the orbital simulation, circular orbit and escape velocity. I know how to write in Latex now. It is very cool and useful. This technique can help me a lot in the future with my working experience. I want to know it better so i can use it competently. I am very interested in this kind of writing because it makes me feel like an IT engineer.

## References

Homer, J. (2014). *Physics*. Oxford, 3rd edition.