

# Common Information: Old and New

**Vincent Y. F. Tan and Lei Yu**

National University of Singapore and Nankai University

2021 International Symposium on Information Theory

# Style of Tutorial

- Based on an **upcoming monograph** by the present speakers

# Style of Tutorial

- Based on an **upcoming monograph** by the present speakers
- Will cover **classical stuff** and more recent advances based on the speaker's knowledge and preferences

# Style of Tutorial

- Based on an **upcoming monograph** by the present speakers
- Will cover **classical stuff** and more recent advances based on the speaker's knowledge and preferences
- Will not be able to touch all bases, e.g., everything I will talk about is **discrete** (and sorry if we don't cite you)

# Style of Tutorial

- Based on an **upcoming monograph** by the present speakers
- Will cover **classical stuff** and more recent advances based on the speaker's knowledge and preferences
- Will not be able to touch all bases, e.g., everything I will talk about is **discrete** (and sorry if we don't cite you)
- Will do some **proof sketches** (since this is a tutorial)

# Style of Tutorial


- Based on an **upcoming monograph** by the present speakers
- Will cover **classical stuff** and more recent advances based on the speaker's knowledge and preferences
- Will not be able to touch all bases, e.g., everything I will talk about is **discrete** (and sorry if we don't cite you)
- Will do some **proof sketches** (since this is a tutorial)
- May get a bit **technical** (no apologies for that)

# Style of Tutorial

- Based on an **upcoming monograph** by the present speakers
- Will cover **classical stuff** and more recent advances based on the speaker's knowledge and preferences
- Will not be able to touch all bases, e.g., everything I will talk about is **discrete** (and sorry if we don't cite you)
- Will do some **proof sketches** (since this is a tutorial)
- May get a bit **technical** (no apologies for that)
- But will try to provide as much **intuition** as possible

# Style of Tutorial

- Based on an **upcoming monograph** by the present speakers
- Will cover **classical stuff** and more recent advances based on the speaker's knowledge and preferences
- Will not be able to touch all bases, e.g., everything I will talk about is **discrete** (and sorry if we don't cite you)
- Will do some **proof sketches** (since this is a tutorial)
- May get a bit **technical** (no apologies for that)
- But will try to provide as much **intuition** as possible
- Prerequisite: Information theory at the level of Cover and Thomas



# Outline

- 1 Introduction: Measures of Information Among Two Random Variables
- 2 Wyner's Common Information
- 3 Rényi Common Information
- 4 Exact Common Information
- 5 Approximate and Exact Channel Synthesis
- 6 Nonnegative Matrix Factorization and Nonnegative Rank
- 7 Extensions of Gäcs–Körner–Witsenhausen's Common Information (L. Yu)

# Measures of Information Among Random Variables

- Given two random variables  $X \in \mathcal{X}$  and  $Y \in \mathcal{Y}$  with joint distribution  $\pi_{XY}$ , how **common** are they?

# Measures of Information Among Random Variables

- Given two random variables  $X \in \mathcal{X}$  and  $Y \in \mathcal{Y}$  with joint distribution  $\pi_{XY}$ , how **common** are they?
- One may conceive of the following measures of "**common information**".

# Measures of Information Among Random Variables

- Given two random variables  $X \in \mathcal{X}$  and  $Y \in \mathcal{Y}$  with joint distribution  $\pi_{XY}$ , how **common** are they?
- One may conceive of the following measures of "**common information**".
- Pearson correlation coefficient

$$\rho(X; Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \in [-1, 1].$$

# Measures of Information Among Random Variables

- Given two random variables  $X \in \mathcal{X}$  and  $Y \in \mathcal{Y}$  with joint distribution  $\pi_{XY}$ , how **common** are they?
- One may conceive of the following measures of "**common information**".
- Pearson correlation coefficient

$$\rho(X; Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \in [-1, 1].$$

- Mutual Information

$$I_\pi(X; Y) = \mathsf{E} \left[ \log \frac{\pi_{XY}(X, Y)}{\pi_X(X)\pi_Y(Y)} \right] = D(\pi_{XY} \| \pi_X\pi_Y).$$

# Measures of Information Among Random Variables

- Given two random variables  $X \in \mathcal{X}$  and  $Y \in \mathcal{Y}$  with joint distribution  $\pi_{XY}$ , how **common** are they?
- One may conceive of the following measures of "**common information**".
- Pearson correlation coefficient

$$\rho(X; Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \in [-1, 1].$$

- Mutual Information

$$I_\pi(X; Y) = \mathbb{E} \left[ \log \frac{\pi_{XY}(X, Y)}{\pi_X(X)\pi_Y(Y)} \right] = D(\pi_{XY} \| \pi_X\pi_Y).$$

- As information theorists, we like **operational interpretations**

# Measures of Information Among Random Variables

- Given two random variables  $X \in \mathcal{X}$  and  $Y \in \mathcal{Y}$  with joint distribution  $\pi_{XY}$ , how **common** are they?
- One may conceive of the following measures of "**common information**".
- Pearson correlation coefficient

$$\rho(X; Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \in [-1, 1].$$

- Mutual Information

$$I_\pi(X; Y) = \mathbb{E} \left[ \log \frac{\pi_{XY}(X, Y)}{\pi_X(X)\pi_Y(Y)} \right] = D(\pi_{XY} \| \pi_X\pi_Y).$$

- As information theorists, we like **operational interpretations**
- Wyner's CI** and **Gäcs–Körner–Witsenhausen's CI** are the two archetypal notions of information among RVs that admit **operational interpretations**.

# Outline

- 1 Introduction: Measures of Information Among Two Random Variables
- 2 Wyner's Common Information
- 3 Rényi Common Information
- 4 Exact Common Information
- 5 Approximate and Exact Channel Synthesis
- 6 Nonnegative Matrix Factorization and Nonnegative Rank
- 7 Extensions of Gäcs–Körner–Witsenhausen's Common Information (L. Yu)


# Wyner's Common Information [Wyner, 1975]

# Wyner's Common Information [Wyner, 1975]

$$\underline{M_n}$$

- $M_n$  is uniformly distributed over  $\mathcal{M}_n = [2^{nR}] := \{1, \dots, 2^{nR}\}$


# Wyner's Common Information [Wyner, 1975]



- $M_n$  is uniformly distributed over  $\mathcal{M}_n = [2^{nR}] := \{1, \dots, 2^{nR}\}$
- An  $(n, R)$ -synthesis code consists of

$$P_{X^n|M_n} : \mathcal{M}_n \rightarrow \mathcal{X}^n \quad \text{and} \quad P_{Y^n|M_n} : \mathcal{M}_n \rightarrow \mathcal{Y}^n.$$

# Wyner's Common Information [Wyner, 1975]




- $M_n$  is uniformly distributed over  $\mathcal{M}_n = [2^{nR}] := \{1, \dots, 2^{nR}\}$
- An  $(n, R)$ -synthesis code consists of

$$P_{X^n|M_n} : \mathcal{M}_n \rightarrow \mathcal{X}^n \quad \text{and} \quad P_{Y^n|M_n} : \mathcal{M}_n \rightarrow \mathcal{Y}^n.$$

- The distribution induced by the code  $(P_{X^n|M_n}, P_{Y^n|M_n})$  is

$$P_{X^n Y^n}(x^n, y^n) := \frac{1}{|\mathcal{M}_n|} \sum_{m \in \mathcal{M}_n} P_{X^n|M_n}(x^n|m) P_{Y^n|M_n}(y^n|m)$$

# Wyner's Common Information [Wyner, 1975]



- $M_n$  is uniformly distributed over  $\mathcal{M}_n = [2^{nR}] := \{1, \dots, 2^{nR}\}$
- An  $(n, R)$ -synthesis code consists of

$$P_{X^n|M_n} : \mathcal{M}_n \rightarrow \mathcal{X}^n \quad \text{and} \quad P_{Y^n|M_n} : \mathcal{M}_n \rightarrow \mathcal{Y}^n.$$

- The distribution induced by the code  $(P_{X^n|M_n}, P_{Y^n|M_n})$  is

$$P_{X^n Y^n}(x^n, y^n) := \frac{1}{|\mathcal{M}_n|} \sum_{m \in \mathcal{M}_n} P_{X^n|M_n}(x^n|m) P_{Y^n|M_n}(y^n|m)$$

- Desideratum:

$$P_{X^n Y^n} \approx \pi_{XY}^n \quad (\text{target distribution})$$

# Wyner's Common Information [Wyner, 1975]

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. IT-21, NO. 2, MARCH 1975

163

## The Common Information of Two Dependent Random Variables

AARON D. WYNER, SENIOR MEMBER, IEEE

# Wyner's Common Information [Wyner, 1975]

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. IT-21, NO. 2, MARCH 1975

163

## The Common Information of Two Dependent Random Variables

AARON D. WYNER, SENIOR MEMBER, IEEE

Normalized **relative entropy** to measure the “distance” between  $P_{X^n Y^n}$  and  $\pi_{XY}^n$

# Wyner's Common Information [Wyner, 1975]

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. IT-21, NO. 2, MARCH 1975

163

## The Common Information of Two Dependent Random Variables

AARON D. WYNER, SENIOR MEMBER, IEEE

Normalized **relative entropy** to measure the “distance” between  $P_{X^n Y^n}$  and  $\pi_{XY}^n$

### Theorem ([Wyner, 1975])

$$\inf \left\{ R : \frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0 \right\}$$

# Wyner's Common Information [Wyner, 1975]

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. IT-21, NO. 2, MARCH 1975

163

## The Common Information of Two Dependent Random Variables

AARON D. WYNER, SENIOR MEMBER, IEEE

Normalized **relative entropy** to measure the “distance” between  $P_{X^n Y^n}$  and  $\pi_{XY}^n$

### Theorem ([Wyner, 1975])

$$\begin{aligned} & \inf \left\{ R : \frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0 \right\} \\ &= \min_{P_W P_{X|W} P_{Y|W} : P_{XY} = \pi_{XY}} I(XY; W) \end{aligned}$$

# Wyner's Common Information [Wyner, 1975]

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. IT-21, NO. 2, MARCH 1975

163

## The Common Information of Two Dependent Random Variables

AARON D. WYNER, SENIOR MEMBER, IEEE

Normalized **relative entropy** to measure the “distance” between  $P_{X^n Y^n}$  and  $\pi_{XY}^n$

### Theorem ([Wyner, 1975])

$$\begin{aligned} & \inf \left\{ R : \frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0 \right\} \\ &= \min_{P_W P_{X|W} P_{Y|W} : P_{XY} = \pi_{XY}} I(XY; W) \\ &=: C_W(\pi_{XY}) \end{aligned}$$

# Wyner's Common Information [Wyner, 1975]

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. IT-21, NO. 2, MARCH 1975

163

## The Common Information of Two Dependent Random Variables

AARON D. WYNER, SENIOR MEMBER, IEEE

Normalized **relative entropy** to measure the “distance” between  $P_{X^n Y^n}$  and  $\pi_{XY}^n$

### Theorem ([Wyner, 1975])

$$\begin{aligned} & \inf \left\{ R : \frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0 \right\} \\ &= \min_{P_W P_{X|W} P_{Y|W} : P_{XY} = \pi_{XY}} I(XY; W) \\ &=: C_W(\pi_{XY}) \end{aligned}$$

# Wyner's Common Information [Wyner, 1975]

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. IT-21, NO. 2, MARCH 1975

163

## The Common Information of Two Dependent Random Variables

AARON D. WYNER, SENIOR MEMBER, IEEE

Normalized **relative entropy** to measure the “distance” between  $P_{X^n Y^n}$  and  $\pi_{XY}^n$

### Theorem ([Wyner, 1975])

$$\begin{aligned} & \inf \left\{ R : \frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0 \right\} \\ &= \min_{P_W P_{X|W} P_{Y|W} : P_{XY} = \pi_{XY}} I(XY; W) \\ &=: C_W(\pi_{XY}) \end{aligned}$$

where  $C_W(\pi_{XY})$  is named **Wyner's Common Information**.

# Sanity Check I

- So Wyner said that a **reasonable notion** of common information is

$$C_W(\pi_{XY}) = \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} I(XY; W).$$

# Sanity Check I

- So Wyner said that a **reasonable notion** of common information is

$$C_W(\pi_{XY}) = \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} I(XY; W).$$

- Let's test this on  $X = (\tilde{X}, V)$  and  $Y = (\tilde{Y}, V)$  with  $\tilde{X}, \tilde{Y}, V$  independent.

# Sanity Check I

- So Wyner said that a **reasonable notion** of common information is

$$C_W(\pi_{XY}) = \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} I(XY; W).$$

- Let's test this on  $X = (\tilde{X}, V)$  and  $Y = (\tilde{Y}, V)$  with  $\tilde{X}, \tilde{Y}, V$  independent.
- Intuitively, we should get  $H(V)$  as the common information. Do we?

# Sanity Check I

- So Wyner said that a **reasonable notion** of common information is

$$C_W(\pi_{XY}) = \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} I(XY; W).$$

- Let's test this on  $X = (\tilde{X}, V)$  and  $Y = (\tilde{Y}, V)$  with  $\tilde{X}, \tilde{Y}, V$  independent.
- Intuitively, we should get  $H(V)$  as the common information. Do we?
- Take  $W = V$ , satisfies  $X - W - Y$ . Then

$$I(XY; W) = I(XY; V) \leq H(V) \quad \text{so far so good...}$$



# Sanity Check II

- Now comes the other part, i.e., to show  $C_W(\pi_{XY}) \geq H(V)$ .

# Sanity Check II

- Now comes the other part, i.e., to show  $C_W(\pi_{XY}) \geq H(V)$ .
- Obviously  $X = (\tilde{X}, V)$  and  $Y = (\tilde{Y}, V)$  and so

$$V - X - W - Y - V.$$

## Sanity Check II

- Now comes the other part, i.e., to show  $C_W(\pi_{XY}) \geq H(V)$ .
- Obviously  $X = (\tilde{X}, V)$  and  $Y = (\tilde{Y}, V)$  and so

$$V - X - W - Y - V.$$

- So  $V$  is a function of  $W$  and

$$I(X, Y; W) = I(\tilde{X}, \tilde{Y}, V; W, V) \geq H(V)$$

## Sanity Check II

- Now comes the other part, i.e., to show  $C_W(\pi_{XY}) \geq H(V)$ .
- Obviously  $X = (\tilde{X}, V)$  and  $Y = (\tilde{Y}, V)$  and so

$$V - X - W - Y - V.$$

- So  $V$  is a function of  $W$  and

$$I(X, Y; W) = I(\tilde{X}, \tilde{Y}, V; W, V) \geq H(V)$$

- Minimize over  $X - W - Y$  so

$$C_W(\pi_{XY}) \geq H(V)$$



# Proof Idea of the Achievability Part

Lemma (Soft-covering lemma [Wyner, 1975] [Cuff, 2012])

Let  $(U, W) \sim P_{UW}$  have mutual information  $I(U; W)$ . For any

$$R > I(U; W),$$

there exists a sequence of codebooks  $\mathcal{C}_n = \{w^n(m) : m \in [2^{nR}]\}$  such that the synthesized distribution

$$P_{U^n}(u^n) = \frac{1}{2^{nR}} \sum_{m=1}^{2^{nR}} P_{U|W}^n(u^n | w^n(m)) \quad \forall n \in \mathbb{N}$$

# Proof Idea of the Achievability Part

Lemma (Soft-covering lemma [Wyner, 1975] [Cuff, 2012])

Let  $(U, W) \sim P_{UW}$  have mutual information  $I(U; W)$ . For any

$$\textcolor{red}{R} > I(U; W),$$

there exists a sequence of codebooks  $\mathcal{C}_n = \{w^n(m) : m \in [2^{\textcolor{red}{nR}}]\}$  such that the synthesized distribution

$$P_{U^n}(u^n) = \frac{1}{2^{nR}} \sum_{m=1}^{2^{nR}} P_{U|W}^n(u^n | w^n(m)) \quad \forall n \in \mathbb{N}$$

satisfies

$$\lim_{n \rightarrow \infty} \frac{1}{n} D(P_{U^n} \| P_U^n) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} |P_{U^n} - P_U^n| = 0 \quad (\text{TV dist}).$$

# Proof Idea of the Achievability Part

Lemma (Soft-covering lemma [Wyner, 1975] [Cuff, 2012])

Let  $(U, W) \sim P_{UW}$  have mutual information  $I(U; W)$ . For any

$$R > I(U; W),$$

there exists a sequence of codebooks  $\mathcal{C}_n = \{w^n(m) : m \in [2^{nR}]\}$  such that the synthesized distribution


$$P_{U^n}(u^n) = \frac{1}{2^{nR}} \sum_{m=1}^{2^{nR}} P_{U|W}^n(u^n | w^n(m)) \quad \forall n \in \mathbb{N}$$

satisfies

$$\lim_{n \rightarrow \infty} \frac{1}{n} D(P_{U^n} \| P_U^n) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} |P_{U^n} - P_U^n| = 0 \quad (\text{TV dist}).$$

Also known as **resolvability** [Han and Verdú, 1993] [Hayashi, 2006].

# Proof Idea of the Achievability Part



**Figure:** If  $M = 2^{nR}$  and  $R > I(U; W)$ , then  $\frac{1}{n} D(P_{U^n} \| P_U^n) \rightarrow 0$ .

# Proof Idea of the Achievability Part





Figure: If  $M = 2^{nR}$  and  $R > I(U; W)$ , then  $\frac{1}{n} D(P_{U^n} \| P_U^n) \rightarrow 0$ .


Now take  $\mathbf{U} = (X, Y) \sim \pi_{XY}$  and note by Markovity  $X - W - Y$  that

$$P_{X^n|M_n}(x^n|m)P_{Y^n|M_n}(y^n|m) = P_{U^n|W^n}(u^n|w^n(m)) \text{ and } I(W; \mathbf{U}) = I(W; XY).$$

# Alternative Interpretation of Wyner's Common Information




# Alternative Interpretation of Wyner's Common Information




- An  $(n, R_0, R_1, R_2)$ -Gray-Wyner code [Gray and Wyner, 1974] consists of

# Alternative Interpretation of Wyner's Common Information




- An  $(n, R_0, R_1, R_2)$ -Gray-Wyner code [Gray and Wyner, 1974] consists of
  - ▶ Three encoders  $f_i : \mathcal{X}^n \times \mathcal{Y}^n \rightarrow [2^{nR_i}]$  where  $i = 0, 1, 2$ ;

# Alternative Interpretation of Wyner's Common Information



- An  $(n, R_0, R_1, R_2)$ -Gray-Wyner code [Gray and Wyner, 1974] consists of
  - ▶ Three encoders  $f_i : \mathcal{X}^n \times \mathcal{Y}^n \rightarrow [2^{nR_i}]$  where  $i = 0, 1, 2$ ;
  - ▶ Two decoders  $\varphi_1 : [2^{nR_0}] \times [2^{nR_1}] \rightarrow \mathcal{X}^n$  and  $\varphi_2 : [2^{nR_0}] \times [2^{nR_2}] \rightarrow \mathcal{Y}^n$ .

# Alternative Interpretation of Wyner's Common Information



- An **( $n, R_0, R_1, R_2$ )-Gray-Wyner code** [Gray and Wyner, 1974] consists of
  - ▶ Three encoders  $f_i : \mathcal{X}^n \times \mathcal{Y}^n \rightarrow [2^{nR_i}]$  where  $i = 0, 1, 2$ ;
  - ▶ Two decoders  $\varphi_1 : [2^{nR_0}] \times [2^{nR_1}] \rightarrow \mathcal{X}^n$  and  $\varphi_2 : [2^{nR_0}] \times [2^{nR_2}] \rightarrow \mathcal{Y}^n$ .
- The **probability of error** of the code is

$$\Pr((\varphi_1(M_0, M_1), \varphi_2(M_0, M_2)) \neq (X^n, Y^n)).$$

where  $M_i = f_i(X^n, Y^n)$  for  $i = 0, 1, 2$ .

# Alternative Interpretation of Wyner's Common Information

Common information based on the Gray-Wyner system  $T_{\text{GW}}(\pi_{XY})$  for  $(X, Y) \sim \pi_{XY}$



Smallest common rate  $R_0$  such that for all  $\epsilon > 0$ , there exists sequence of  $(n, R_0, R_1, R_2)$  Gray-Wyner codes  $\{(f_{0,n}, f_{1,n}, f_{2,n}, \varphi_{1,n}, \varphi_{2,n})\}_{n=1}^{\infty}$  such that

$$R_0 + R_1 + R_2 \leq H(XY) + \epsilon$$

and the probability of error vanishes.

# Alternative Interpretation of Wyner's Common Information

Common information based on the Gray-Wyner system  $T_{\text{GW}}(\pi_{XY})$  for  $(X, Y) \sim \pi_{XY}$



Smallest common rate  $R_0$  such that for all  $\epsilon > 0$ , there exists sequence of  $(n, R_0, R_1, R_2)$  Gray-Wyner codes  $\{(f_{0,n}, f_{1,n}, f_{2,n}, \varphi_{1,n}, \varphi_{2,n})\}_{n=1}^{\infty}$  such that

$$R_0 + R_1 + R_2 \leq H(XY) + \epsilon$$

and the probability of error vanishes.

Theorem ([Wyner, 1975])

$$T_{\text{GW}}(\pi_{XY}) = C_{\text{W}}(\pi_{XY})$$

# Example: Doubly Symmetric Binary Source (DSBS)

- Consider a DSBS  $(X, Y) \in \{0, 1\}^2$  which is defined for  $p \in (0, 1/2)$  by


$$\pi_{XY} = \begin{bmatrix} (1-p)/2 & p/2 \\ p/2 & (1-p)/2 \end{bmatrix}$$

# Example: Doubly Symmetric Binary Source (DSBS)

- Consider a DSBS  $(X, Y) \in \{0, 1\}^2$  which is defined for  $p \in (0, 1/2)$  by

$$\pi_{XY} = \begin{bmatrix} (1-p)/2 & p/2 \\ p/2 & (1-p)/2 \end{bmatrix}$$

- Interpretation in terms of  $X - W - Y$




# Example: Doubly Symmetric Binary Source (DSBS)

- Consider a DSBS  $(X, Y) \in \{0, 1\}^2$  which is defined for  $p \in (0, 1/2)$  by

$$\pi_{XY} = \begin{bmatrix} (1-p)/2 & p/2 \\ p/2 & (1-p)/2 \end{bmatrix}$$

- Interpretation in terms of  $X - W - Y$





# Example: Doubly Symmetric Binary Source (DSBS)

- Consider a DSBS  $(X, Y) \in \{0, 1\}^2$  which is defined for  $p \in (0, 1/2)$  by

$$\pi_{XY} = \begin{bmatrix} (1-p)/2 & p/2 \\ p/2 & (1-p)/2 \end{bmatrix}$$

- Interpretation in terms of  $X - W - Y$





# Example: Doubly Symmetric Binary Source (DSBS)

- Consider a DSBS  $(X, Y) \in \{0, 1\}^2$  which is defined for  $p \in (0, 1/2)$  by

$$\pi_{XY} = \begin{bmatrix} (1-p)/2 & p/2 \\ p/2 & (1-p)/2 \end{bmatrix}$$

- Interpretation in terms of  $X - W - Y$



- Here,  $a * a = p$  and

$$a = \frac{1 - \sqrt{1 - 2p}}{2} \in (0, 1/2).$$

# Example: DSBS





Figure: Plots of Wyner's common information for the DSBS in terms of  $p$  and  $a$

# Outline

- 1 Introduction: Measures of Information Among Two Random Variables
- 2 Wyner's Common Information
- 3 Rényi Common Information
- 4 Exact Common Information
- 5 Approximate and Exact Channel Synthesis
- 6 Nonnegative Matrix Factorization and Nonnegative Rank
- 7 Extensions of Gäcs–Körner–Witsenhausen's Common Information (L. Yu)

# Motivation for Alternative Measures

- Wyner used the **normalized** relative entropy, i.e.,

$$\inf \left\{ R : \lim_{n \rightarrow \infty} \frac{D(P_{X^n Y^n} \| \pi_{XY}^n)}{n} = 0 \right\} = C_W(\pi_{XY}) = \min_{X-W-Y} I(W; XY).$$

# Motivation for Alternative Measures

- Wyner used the **normalized** relative entropy, i.e.,

$$\inf \left\{ R : \lim_{n \rightarrow \infty} \frac{D(P_{X^n Y^n} \| \pi_{XY}^n)}{n} = 0 \right\} = C_W(\pi_{XY}) = \min_{X-W-Y} I(W; XY).$$

- What if we do not normalize?

$$\tilde{T}(\pi_{XY}) := \inf \left\{ R : \lim_{n \rightarrow \infty} D(P_{X^n Y^n} \| \pi_{XY}^n) = 0 \right\} \geq C_W(\pi_{XY}).$$

We get a **stronger** measure of dependence.

# Motivation for Alternative Measures

- Wyner used the **normalized** relative entropy, i.e.,

$$\inf \left\{ R : \lim_{n \rightarrow \infty} \frac{D(P_{X^n Y^n} \| \pi_{XY}^n)}{n} = 0 \right\} = C_W(\pi_{XY}) = \min_{X-W-Y} I(W; XY).$$

- What if we do not normalize?

$$\tilde{T}(\pi_{XY}) := \inf \left\{ R : \lim_{n \rightarrow \infty} D(P_{X^n Y^n} \| \pi_{XY}^n) = 0 \right\} \geq C_W(\pi_{XY}).$$

We get a **stronger** measure of dependence.

- What if we want an **even stronger** measure of dependence?

# Motivation for Alternative Measures

- Wyner used the **normalized** relative entropy, i.e.,

$$\inf \left\{ R : \lim_{n \rightarrow \infty} \frac{D(P_{X^n Y^n} \| \pi_{XY}^n)}{n} = 0 \right\} = C_W(\pi_{XY}) = \min_{X-W-Y} I(W; XY).$$

- What if we do not normalize?

$$\tilde{T}(\pi_{XY}) := \inf \left\{ R : \lim_{n \rightarrow \infty} D(P_{X^n Y^n} \| \pi_{XY}^n) = 0 \right\} \geq C_W(\pi_{XY}).$$

We get a **stronger** measure of dependence.

- What if we want an **even stronger** measure of dependence?
- Rényi common information for orders  $\geq 1$  [Yu and Tan, 2018]!

$$T_{1+s}(\pi_{XY}) := \inf \left\{ R : \lim_{n \rightarrow \infty} \frac{D_{1+s}(P_{X^n Y^n} \| \pi_{XY}^n)}{n} = 0 \right\}$$

$$\tilde{T}_{1+s}(\pi_{XY}) := \inf \left\{ R : \lim_{n \rightarrow \infty} D_{1+s}(P_{X^n Y^n} \| \pi_{XY}^n) = 0 \right\}$$

# Rényi Common Information

- Rényi divergence

$$D_{1+s}(P\|Q) := \frac{1}{s} \log \sum_{x \in \text{supp}(P)} P(x) \left( \frac{P(x)}{Q(x)} \right)^s \quad s \in [-1, \infty)$$
$$D_\infty(P\|Q) := \log \max_{x \in \text{supp}(P)} \frac{P(x)}{Q(x)}.$$

# Rényi Common Information

- Rényi divergence

$$D_{1+s}(P\|Q) := \frac{1}{s} \log \sum_{x \in \text{supp}(P)} P(x) \left( \frac{P(x)}{Q(x)} \right)^s \quad s \in [-1, \infty)$$
$$D_\infty(P\|Q) := \log \max_{x \in \text{supp}(P)} \frac{P(x)}{Q(x)}.$$

- The Rényi divergence is monotonically non-decreasing, i.e.,

$$D_{1+s}(P\|Q) \leq D_{1+t}(P\|Q) \quad s \leq t.$$

# Rényi Common Information

- Rényi divergence

$$D_{1+s}(P\|Q) := \frac{1}{s} \log \sum_{x \in \text{supp}(P)} P(x) \left( \frac{P(x)}{Q(x)} \right)^s \quad s \in [-1, \infty)$$
$$D_\infty(P\|Q) := \log \max_{x \in \text{supp}(P)} \frac{P(x)}{Q(x)}.$$

- The Rényi divergence is monotonically non-decreasing, i.e.,

$$D_{1+s}(P\|Q) \leq D_{1+t}(P\|Q) \quad s \leq t.$$

- Hence, the Rényi common information is also non-decreasing, i.e.,

$$\text{(normalized)} \quad T_{1+s}(\pi_{XY}) \leq T_{1+t}(\pi_{XY}) \quad s \leq t.$$

and

$$\text{(unnormalized)} \quad \tilde{T}_{1+s}(\pi_{XY}) \leq \tilde{T}_{1+t}(\pi_{XY}) \quad s \leq t.$$

# Rényi Common Information

- Rényi divergence

$$D_{1+s}(P\|Q) := \frac{1}{s} \log \sum_{x \in \text{supp}(P)} P(x) \left( \frac{P(x)}{Q(x)} \right)^s \quad s \in [-1, \infty)$$

$$D_\infty(P\|Q) := \log \max_{x \in \text{supp}(P)} \frac{P(x)}{Q(x)}.$$

- The Rényi divergence is monotonically non-decreasing, i.e.,

$$D_{1+s}(P\|Q) \leq D_{1+t}(P\|Q) \quad s \leq t.$$

- Hence, the Rényi common information is also non-decreasing, i.e.,

$$\text{(normalized)} \quad T_{1+s}(\pi_{XY}) \leq T_{1+t}(\pi_{XY}) \quad s \leq t.$$

and

$$\text{(unnormalized)} \quad \tilde{T}_{1+s}(\pi_{XY}) \leq \tilde{T}_{1+t}(\pi_{XY}) \quad s \leq t.$$

- And for a fixed order  $1 + s \in [0, \infty]$ ,

$$T_{1+s}(\pi_{XY}) \leq \tilde{T}_{1+s}(\pi_{XY}).$$

# Are we doing math for the sake of doing math?

# Are we doing math for the sake of doing math?

- The sceptic in you might wonder whether we are **just** doing math.

# Are we doing math for the sake of doing math?

- The sceptic in you might wonder whether we are **just** doing math.
- In fact not! We show in the sequel that

$$\tilde{T}_\infty(\pi_{XY}) = \text{Exact Common Information of } \pi_{XY}.$$

Exact Common Information was introduced by [Kumar et al., 2014].

# Are we doing math for the sake of doing math?

- The sceptic in you might wonder whether we are **just** doing math.
- In fact not! We show in the sequel that

$$\tilde{T}_\infty(\pi_{XY}) = \text{Exact Common Information of } \pi_{XY}.$$

Exact Common Information was introduced by [Kumar et al., 2014].

- And it is through this unexpected connection that we show that

$$\text{Exact Common Information of } \pi_{XY} > C_W(\pi_{XY})$$

for some joint sources  $\pi_{XY}$ .

# Are we doing math for the sake of doing math?

- The sceptic in you might wonder whether we are **just** doing math.
- In fact not! We show in the sequel that

$$\tilde{T}_\infty(\pi_{XY}) = \text{Exact Common Information of } \pi_{XY}.$$

Exact Common Information was introduced by [Kumar et al., 2014].

- And it is through this unexpected connection that we show that

$$\text{Exact Common Information of } \pi_{XY} > C_W(\pi_{XY})$$

for some joint sources  $\pi_{XY}$ .

- But let's **soldier on** and tackle the Rényi common information for now.



# Rényi Common Information: The Weaker Case

# Rényi Common Information: The Weaker Case

Let's start with a simple exercise. Consider the case  $s \in (-1, 0]$  in which

$$T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$$

# Rényi Common Information: The Weaker Case

Let's start with a simple exercise. Consider the case  $s \in (-1, 0]$  in which

$$T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$$

Theorem ([Yu and Tan, 2018] [Yu and Tan, 2020a])

For Rényi orders in  $(0, 1]$  (i.e.,  $s \in (-1, 0]$ ),

$$T_{1+s}(\pi_{XY}) = \tilde{T}_{1+s}(\pi_{XY}) = C_W(\pi_{XY}).$$

# Rényi Common Information: The Weaker Case

Let's start with a simple exercise. Consider the case  $s \in (-1, 0]$  in which

$$T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$$

**Theorem ([Yu and Tan, 2018] [Yu and Tan, 2020a])**

For Rényi orders in  $(0, 1]$  (i.e.,  $s \in (-1, 0]$ ),

$$T_{1+s}(\pi_{XY}) = \tilde{T}_{1+s}(\pi_{XY}) = C_W(\pi_{XY}).$$

Our stepping stone...

# Rényi Common Information: The Weaker Case

Let's start with a simple exercise. Consider the case  $s \in (-1, 0]$  in which

$$T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$$

**Theorem ([Yu and Tan, 2018] [Yu and Tan, 2020a])**

For Rényi orders in  $(0, 1]$  (i.e.,  $s \in (-1, 0]$ ),

$$T_{1+s}(\pi_{XY}) = \tilde{T}_{1+s}(\pi_{XY}) = C_W(\pi_{XY}).$$

Our stepping stone... Total variation distance  $|P - Q| := \frac{1}{2} \sum_x |P(x) - Q(x)|$ .

# Rényi Common Information: The Weaker Case

Let's start with a simple exercise. Consider the case  $s \in (-1, 0]$  in which

$$T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$$

**Theorem ([Yu and Tan, 2018] [Yu and Tan, 2020a])**

For Rényi orders in  $(0, 1]$  (i.e.,  $s \in (-1, 0]$ ),

$$T_{1+s}(\pi_{XY}) = \tilde{T}_{1+s}(\pi_{XY}) = C_W(\pi_{XY}).$$

Our stepping stone... Total variation distance  $|P - Q| := \frac{1}{2} \sum_x |P(x) - Q(x)|$ .

**Theorem ([Yu and Tan, 2018])**


For any  $\varepsilon \in [0, 1)$ ,

$$T_\varepsilon^{\text{TV}}(\pi_{XY}) = C_W(\pi_{XY}), \quad (\text{Strong converse})$$


where  $T_\varepsilon^{\text{TV}}(\pi_{XY})$  is the minimum simulation rate required to ensure

$$\limsup_{n \rightarrow \infty} |P_{X^n Y^n} - \pi_{XY}^n| \leq \varepsilon.$$


# Total Variation Common Information



# Total Variation Common Information




# Total Variation Common Information



In fact, we have an **exponential strong converse**, i.e., if  $R < C_W(\pi_{XY})$ ,

$$|P_{X^n Y^n} - \pi_{XY}^n| \geq 1 - 2^{-nE} \quad \text{for some } E > 0.$$

# Total Variation Common Information



In fact, we have an **exponential strong converse**, i.e., if  $R < C_W(\pi_{XY})$ ,

$$|P_{X^n Y^n} - \pi_{XY}^n| \geq 1 - 2^{-nE} \quad \text{for some } E > 0.$$

Amenable to **second-order**?

# Total Variation Common Information

- Achievability part follows from the soft-covering lemma.

If  $R > I(XY; W)$  then  $\lim_{n \rightarrow \infty} |P_{X^n Y^n} - \pi_{XY}^n| = 0$ .

# Total Variation Common Information

- Achievability part follows from the soft-covering lemma.

$$\text{If } R > I(XY; W) \text{ then } \lim_{n \rightarrow \infty} |P_{X^n Y^n} - \pi_{XY}^n| = 0.$$

- Converse requires a very cool information spectrum, single-letterization idea from [Oohama, 2018].



Article

## Exponential Strong Converse for Source Coding with Side Information at the Decoder<sup>†</sup>

Yasutada Oohama

Department of Communication Engineering and Informatics, University of Electro-Communications, Tokyo 182-8585, Japan; oohama@uec.ac.jp; Tel.: +81-42-443-5358

<sup>†</sup> This paper is an extended version of our paper published in 2016 International Symposium on Information Theory and Its Applications, Monterey, CA, USA, 6–9 November 2016; pp. 171–175.

Received: 31 January 2018; Accepted: 20 April 2018; Published: 8 May 2018



# Going Back to Rényi CI: The Weaker Case $s \in (-1, 0]$

- Because  $T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$ , only have to prove the converse.

# Going Back to Rényi CI: The Weaker Case $s \in (-1, 0]$

- Because  $T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$ , only have to prove the converse.
- Main idea is a **Pinsker-type inequality** due to [Sason, 2016].

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. 62, NO. 1, JANUARY 2016

23

## On the Rényi Divergence, Joint Range of Relative Entropies, and a Channel Coding Theorem

Igal Sason, *Senior Member, IEEE*

# Going Back to Rényi CI: The Weaker Case $s \in (-1, 0]$

- Because  $T_{1+s}(\pi_{XY}) \leq C_W(\pi_{XY})$ , only have to prove the converse.
- Main idea is a **Pinsker-type inequality** due to [Sason, 2016].

IEEE TRANSACTIONS ON INFORMATION THEORY, VOL. 62, NO. 1, JANUARY 2016

23

## On the Rényi Divergence, Joint Range of Relative Entropies, and a Channel Coding Theorem

Igal Sason, *Senior Member, IEEE*

### Lemma

For any  $s \in (-1, 0]$ ,

$$\inf_{P_X, Q_X : |P_X - Q_X| \geq \epsilon} D_{1+s}(P_X \| Q_X) = \inf_{q \in [0, 1-\epsilon]} d_{1+s}(q + \epsilon \| q)$$

and

$$\inf_{q \in [0, 1-\epsilon]} d_{1+s}(q + \epsilon \| q) \geq \left[ \min \left\{ 1, \frac{1+s}{s} \right\} \log \frac{1}{1-\epsilon} + \frac{1}{s} \log 2 \right]^+$$

# Going Back to Rényi CI: The Weaker Case $s \in (-1, 0]$

- From [Sason, 2016], we have

$$\inf_{P_X, Q_X : |P_X - Q_X| \geq \epsilon} D_{1+s}(P_X \| Q_X) \geq \left[ \min \left\{ 1, \frac{1+s}{s} \right\} \log \frac{1}{1-\epsilon} + \frac{1}{s} \log 2 \right]^+$$

# Going Back to Rényi CI: The Weaker Case $s \in (-1, 0]$

- From [Sason, 2016], we have

$$\inf_{P_X, Q_X : |P_X - Q_X| \geq \epsilon} D_{1+s}(P_X \| Q_X) \geq \left[ \min \left\{ 1, \frac{1+s}{s} \right\} \log \frac{1}{1-\epsilon} + \frac{1}{s} \log 2 \right]^+$$

- If  $R < C_W(\pi_{XY})$ , exponential strong converse to TV CI says

$$|P_{X^n Y^n} - \pi_{XY}^n| \geq 1 - 2^{-nE} \quad \text{for some } E > 0.$$

# Going Back to Rényi CI: The Weaker Case $s \in (-1, 0]$

- From [Sason, 2016], we have

$$\inf_{P_X, Q_X : |P_X - Q_X| \geq \epsilon} D_{1+s}(P_X \| Q_X) \geq \left[ \min \left\{ 1, \frac{1+s}{s} \right\} \log \frac{1}{1-\epsilon} + \frac{1}{s} \log 2 \right]^+$$

- If  $R < C_W(\pi_{XY})$ , exponential strong converse to TV CI says

$$|P_{X^n Y^n} - \pi_{XY}^n| \geq 1 - 2^{-nE} \quad \text{for some } E > 0.$$

- Thus, if  $R < C_W(\pi_{XY})$

$$\frac{1}{n} \inf_{P_X, Q_X : |P_X - Q_X| \geq \epsilon} D_{1+s}(P_X \| Q_X) \geq \frac{1}{n} \left[ \min \left\{ 1, \frac{1+s}{s} \right\} nE + \frac{1}{s} \log 2 \right]^+$$

and the normalized Rényi divergence cannot vanish.

# Rényi CI: The Stronger Case $s \in (0, 1] \cup \{\infty\}$

- For  $s \in (0, 1] \cup \{\infty\}$ ,

$$C_W(\pi_{XY}) \leq T_{1+s}(\pi_{XY}).$$

# Rényi CI: The Stronger Case $s \in (0, 1] \cup \{\infty\}$

- For  $s \in (0, 1] \cup \{\infty\}$ ,

$$C_W(\pi_{XY}) \leq T_{1+s}(\pi_{XY}).$$

- We only discuss the case  $s = \infty$  in this tutorial.

# Rényi CI: The Stronger Case $s \in (0, 1] \cup \{\infty\}$

- For  $s \in (0, 1] \cup \{\infty\}$ ,

$$C_W(\pi_{XY}) \leq T_{1+s}(\pi_{XY}).$$

- We only discuss the case  $s = \infty$  in this tutorial.
- For the other cases (i.e.,  $s \geq 1$  finite), see our upcoming monograph.

# Rényi CI: The Stronger Case $s \in (0, 1] \cup \{\infty\}$

- For  $s \in (0, 1] \cup \{\infty\}$ ,

$$C_W(\pi_{XY}) \leq T_{1+s}(\pi_{XY}).$$

- We only discuss the case  $s = \infty$  in this tutorial.
- For the other cases (i.e.,  $s \geq 1$  finite), see our upcoming monograph.

## Definition

The **maximal cross entropy** w.r.t.  $(X, Y) \sim \pi_{XY}$  over couplings of  $(P_X, P_Y)$  is

$$H_\infty(P_X, P_Y \| \pi_{XY}) := \max_{Q_{XY} \in \mathcal{C}(P_X, P_Y)} \sum_{x,y} Q_{XY}(x, y) \log \frac{1}{\pi_{XY}(x, y)},$$

where

$$\mathcal{C}(P_X, P_Y) := \{Q_{XY} \in \mathcal{P}(\mathcal{X} \times \mathcal{Y}) : Q_X = P_X, Q_Y = P_Y\}.$$

# Rényi CI: The Stronger Case $s \in (0, 1] \cup \{\infty\}$

- For  $s \in (0, 1] \cup \{\infty\}$ ,

$$C_W(\pi_{XY}) \leq T_{1+s}(\pi_{XY}).$$

- We only discuss the case  $s = \infty$  in this tutorial.
- For the other cases (i.e.,  $s \geq 1$  finite), see our upcoming monograph.

## Definition

The **maximal cross entropy** w.r.t.  $(X, Y) \sim \pi_{XY}$  over couplings of  $(P_X, P_Y)$  is

$$H_\infty(P_X, P_Y \| \pi_{XY}) := \max_{Q_{XY} \in \mathcal{C}(P_X, P_Y)} \sum_{x,y} Q_{XY}(x, y) \log \frac{1}{\pi_{XY}(x, y)},$$

where

$$\mathcal{C}(P_X, P_Y) := \{Q_{XY} \in \mathcal{P}(\mathcal{X} \times \mathcal{Y}) : Q_X = P_X, Q_Y = P_Y\}.$$

- $H_\infty(\pi_X, \pi_Y \| \pi_{XY}) \geq H_\pi(X; Y)$  with equality iff  $\pi_{XY} = \pi_X \pi_Y$ .

# Intuition for the Maximal Cross Entropy

- Take a sequence of  $n$ -types  $T_X^{(n)} \in \mathcal{P}_n(\mathcal{X})$  and  $T_Y^{(n)} \in \mathcal{P}_n(\mathcal{Y})$ .

# Intuition for the Maximal Cross Entropy

- Take a sequence of  $n$ -types  $T_X^{(n)} \in \mathcal{P}_n(\mathcal{X})$  and  $T_Y^{(n)} \in \mathcal{P}_n(\mathcal{Y})$ .
- Let them converge as

$$T_X^{(n)} \rightarrow P_X \quad \text{and} \quad T_Y^{(n)} \rightarrow P_Y.$$

# Intuition for the Maximal Cross Entropy

- Take a sequence of  $n$ -types  $T_X^{(n)} \in \mathcal{P}_n(\mathcal{X})$  and  $T_Y^{(n)} \in \mathcal{P}_n(\mathcal{Y})$ .
- Let them converge as

$$T_X^{(n)} \rightarrow P_X \quad \text{and} \quad T_Y^{(n)} \rightarrow P_Y.$$

- What's the **minimum  $\pi_{XY}^n$ -probability** of  $(x^n, y^n)$  where  $x^n$  has type  $T_X^{(n)}$  and  $y^n$  has type  $T_Y^{(n)}$ , i.e.,

$$\min_{T_{x^n} = T_X^{(n)}, T_{y^n} = T_Y^{(n)}} \pi_{XY}^n(x^n, y^n)?$$

# Intuition for the Maximal Cross Entropy

- Take a sequence of  $n$ -types  $T_X^{(n)} \in \mathcal{P}_n(\mathcal{X})$  and  $T_Y^{(n)} \in \mathcal{P}_n(\mathcal{Y})$ .
- Let them converge as
$$T_X^{(n)} \rightarrow P_X \quad \text{and} \quad T_Y^{(n)} \rightarrow P_Y.$$
- What's the **minimum  $\pi_{XY}^n$ -probability** of  $(x^n, y^n)$  where  $x^n$  has type  $T_X^{(n)}$  and  $y^n$  has type  $T_Y^{(n)}$ , i.e.,

$$\min_{T_{x^n} = T_X^{(n)}, T_{y^n} = T_Y^{(n)}} \pi_{XY}^n(x^n, y^n)?$$

- By type gymnastics,

$$\min_{T_{x^n} = T_X^{(n)}, T_{y^n} = T_Y^{(n)}} \pi_{XY}^n(x^n, y^n) \doteq \exp\left(-H_\infty(P_X, P_Y \| \pi_{XY})\right).$$

# Intuition for the Maximal Cross Entropy

- Take a sequence of  $n$ -types  $T_X^{(n)} \in \mathcal{P}_n(\mathcal{X})$  and  $T_Y^{(n)} \in \mathcal{P}_n(\mathcal{Y})$ .
- Let them converge as

$$T_X^{(n)} \rightarrow P_X \quad \text{and} \quad T_Y^{(n)} \rightarrow P_Y.$$

- What's the **minimum  $\pi_{XY}^n$ -probability** of  $(x^n, y^n)$  where  $x^n$  has type  $T_X^{(n)}$  and  $y^n$  has type  $T_Y^{(n)}$ , i.e.,

$$\min_{T_{x^n} = T_X^{(n)}, T_{y^n} = T_Y^{(n)}} \pi_{XY}^n(x^n, y^n)?$$

- By type gymnastics,

$$\min_{T_{x^n} = T_X^{(n)}, T_{y^n} = T_Y^{(n)}} \pi_{XY}^n(x^n, y^n) \doteq \exp(-H_\infty(P_X, P_Y \| \pi_{XY})).$$

- So  $H_\infty(P_X, P_Y \| \pi_{XY})$  is the exponential decay rate of this probability.

# Upper and Lower Pseudo Common Informations

## Definition

The **upper pseudo-common information** is

$$\bar{\Gamma}_\infty(\pi_{XY}) := \min_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) + \mathbb{E}_{P_W} [\mathsf{H}_\infty(P_{X|W}, P_{Y|W} \| \pi_{XY})]$$

# Upper and Lower Pseudo Common Informations

## Definition

The upper pseudo-common information is

$$\bar{\Gamma}_{\infty}(\pi_{XY}) := \min_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) + \mathsf{E}_{P_W} [\mathsf{H}_{\infty}(P_{X|W}, P_{Y|W} \| \pi_{XY})]$$

Contrast to Wyner's common information

$$C_W(\pi_{XY}) = \min_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) + \mathsf{H}(XY).$$

# Upper and Lower Pseudo Common Informations

## Definition

The **upper pseudo-common information** is

$$\bar{\Gamma}_{\infty}(\pi_{XY}) := \min_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) + \mathsf{E}_{P_W} [\mathsf{H}_{\infty}(P_{X|W}, P_{Y|W} \| \pi_{XY})]$$

Contrast to Wyner's common information

$$C_W(\pi_{XY}) = \min_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) + \mathsf{H}(XY).$$

## Definition

The **lower pseudo-common information** is

$$\begin{aligned} \underline{\Gamma}_{\infty}(\pi_{XY}) &:= \inf_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) \\ &+ \inf_{Q_{WW'} \in \mathcal{C}(P_W, P_W)} \mathsf{E}_{Q_{WW'}} [\mathsf{H}_{\infty}(P_{X|W}, P_{Y|W'} \| \pi_{XY})]. \end{aligned}$$

# Upper and Lower Pseudo Common Informations

## Definition

The upper pseudo-common information is

$$\bar{\Gamma}_{\infty}(\pi_{XY}) := \min_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) + \mathbb{E}_{P_W} [\mathsf{H}_{\infty}(P_{X|W}, P_{Y|W} \| \pi_{XY})]$$

Contrast to Wyner's common information

$$C_W(\pi_{XY}) = \min_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) + H(XY).$$

## Definition

The lower pseudo-common information is

$$\begin{aligned} \underline{\Gamma}_{\infty}(\pi_{XY}) &:= \inf_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} -H(XY|W) \\ &+ \inf_{Q_{WW'} \in \mathcal{C}(P_W, P_W)} \mathbb{E}_{Q_{WW'}} [\mathsf{H}_{\infty}(P_{X|W}, P_{Y|W'} \| \pi_{XY})]. \end{aligned}$$

# Rényi Common Information of order $\infty$

Theorem ([Yu and Tan, 2020a] [Yu and Tan, 2020c])

*The order- $\infty$  Rényi common information admits the following single-letter bounds*

$$\tilde{T}_\infty(\pi_{XY}) \geq T_\infty(\pi_{XY}) \geq \max \{\underline{\Gamma}_\infty(\pi_{XY}), C_W(\pi_{XY})\}$$

and

$$T_\infty(\pi_{XY}) \leq \tilde{T}_\infty(\pi_{XY}) \leq \bar{\Gamma}_\infty(\pi_{XY}).$$

# Rényi Common Information of order $\infty$

Theorem ([Yu and Tan, 2020a] [Yu and Tan, 2020c])

*The order- $\infty$  Rényi common information admits the following single-letter bounds*

$$\tilde{T}_\infty(\pi_{XY}) \geq T_\infty(\pi_{XY}) \geq \max \{\underline{\Gamma}_\infty(\pi_{XY}), C_W(\pi_{XY})\}$$

and

$$T_\infty(\pi_{XY}) \leq \tilde{T}_\infty(\pi_{XY}) \leq \bar{\Gamma}_\infty(\pi_{XY}).$$

**Achievability:** Rényi soft-covering [Yu and Tan, 2019] and truncated product distributions.

# Rényi Common Information of order $\infty$

Theorem ([Yu and Tan, 2020a] [Yu and Tan, 2020c])


The order- $\infty$  Rényi common information admits the following single-letter bounds

$$\tilde{T}_\infty(\pi_{XY}) \geq T_\infty(\pi_{XY}) \geq \max \{\underline{\Gamma}_\infty(\pi_{XY}), C_W(\pi_{XY})\}$$

and

$$T_\infty(\pi_{XY}) \leq \tilde{T}_\infty(\pi_{XY}) \leq \bar{\Gamma}_\infty(\pi_{XY}).$$

**Achievability:** Rényi soft-covering [Yu and Tan, 2019] and truncated product distributions.



Product distribution

$$P_W^n(w^n) = \prod_{i=1}^n P_W(w_i)$$

# Rényi Common Information of order $\infty$

Theorem ([Yu and Tan, 2020a] [Yu and Tan, 2020c])


The order- $\infty$  Rényi common information admits the following single-letter bounds

$$\tilde{T}_\infty(\pi_{XY}) \geq T_\infty(\pi_{XY}) \geq \max \{\underline{\Gamma}_\infty(\pi_{XY}), C_W(\pi_{XY})\}$$

and

$$T_\infty(\pi_{XY}) \leq \tilde{T}_\infty(\pi_{XY}) \leq \bar{\Gamma}_\infty(\pi_{XY}).$$

**Achievability:** Rényi soft-covering [Yu and Tan, 2019] and truncated product distributions.



Truncated product distribution


$$P_{W^n}(w^n) \propto \left( \prod_{i=1}^n P_W(w_i) \right) \mathbb{1}\{w^n \in \mathcal{S}\}$$

# Rényi Common Information of other orders $\in (1, \infty)$ ?

- Can obtain similar bounds [Yu and Tan, 2020a]


# Rényi Common Information of other orders $\in (1, \infty)$ ?

- Can obtain similar bounds [Yu and Tan, 2020a]
- For the DSBS, for  $1 + s \in [0, 2]$ , after some calculations, we get



# Rényi Common Information of other orders $\in (1, \infty)$ ?


- Can obtain similar bounds [Yu and Tan, 2020a]
- For the DSBS, for  $1 + s \in [0, 2]$ , after some calculations, we get



- Rényi common information for the DSBS **increases** with  $1 + s \in [1, 2]!$

# Rényi Common Information of other orders $\in (1, \infty)$ ?

- Can obtain similar bounds [Yu and Tan, 2020a]
- For the DSBS, for  $1 + s \in [0, 2]$ , after some calculations, we get



- Rényi common information for the DSBS **increases** with  $1 + s \in [1, 2]$ !

Does this have more **profound** implications?



# Outline

- 1 Introduction: Measures of Information Among Two Random Variables
- 2 Wyner's Common Information
- 3 Rényi Common Information
- 4 Exact Common Information
- 5 Approximate and Exact Channel Synthesis
- 6 Nonnegative Matrix Factorization and Nonnegative Rank
- 7 Extensions of Gäcs–Körner–Witsenhausen's Common Information (L. Yu)

# Exact Common Information?

- In the distributed source simulation problem à la Wyner, we mandated that

$$\frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0.$$

# Exact Common Information?

- In the distributed source simulation problem à la Wyner, we mandated that

$$\frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0.$$

- What if we require

$$P_{X^n Y^n} = \pi_{XY}^n \quad \text{for some } n \in \mathbb{N}?$$

# Exact Common Information?

- In the distributed source simulation problem à la Wyner, we mandated that

$$\frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0.$$

- What if we require

$$P_{X^n Y^n} = \pi_{XY}^n \quad \text{for some } n \in \mathbb{N}?$$

- Using **fixed-length block codes**, we need rate  $\lim_{n \rightarrow \infty} \frac{1}{n} \log |\mathcal{W}_n|$  over  $W \in \mathcal{W}_n$  such that  $X^n - W - Y^n$ ! Potentially up to  $\min\{\log |\mathcal{X}|, \log |\mathcal{Y}|\}$ .



# Exact Common Information?

- In the distributed source simulation problem à la Wyner, we mandated that

$$\frac{1}{n} D(P_{X^n Y^n} \| \pi_{XY}^n) \rightarrow 0.$$

- What if we require

$$P_{X^n Y^n} = \pi_{XY}^n \quad \text{for some } n \in \mathbb{N}?$$

- Using **fixed-length block codes**, we need rate  $\lim_{n \rightarrow \infty} \frac{1}{n} \log |\mathcal{W}_n|$  over  $W \in \mathcal{W}_n$  such that  $X^n - W - Y^n$ ! Potentially up to  $\min\{\log |\mathcal{X}|, \log |\mathcal{Y}|\}$ .



- In come [Kumar et al., 2014], who introduced

2014 IEEE International Symposium on Information Theory


## Exact Common Information

Gowtham Ramani Kumar  
Electrical Engineering  
Stanford University  
Email: gowthamr@stanford.edu


Cheuk Ting Li  
Electrical Engineering  
Stanford University  
Email: ctli@stanford.edu

Abbas El Gamal  
Electrical Engineering  
Stanford University  
Email: abbas@stanford.edu

# Exact Common Information




# Exact Common Information




- A synthesis code ( $P_{W_n}$ ,  $P_{X^n|W_n}$ ,  $P_{Y^n|W_n}$ )

# Exact Common Information



- A synthesis code ( $P_{W_n}, P_{X^n|W_n}, P_{Y^n|W_n}$ )
- $W_n$  can be any (not necessarily uniform) discrete random variable


# Exact Common Information



- A synthesis code ( $P_{W_n}$ ,  $P_{X^n|W_n}$ ,  $P_{Y^n|W_n}$ )
- $W_n$  can be any (not necessarily uniform) discrete random variable
- Distribution induced by the code is

$$P_{X^n Y^n}(x^n, y^n) := \sum_w P_{W_n}(w) P_{X^n|W_n}(x^n|w) P_{Y^n|W_n}(y^n|w).$$

# Exact Common Information



- A synthesis code ( $P_{W_n}$ ,  $P_{X^n|W_n}$ ,  $P_{Y^n|W_n}$ )
- $W_n$  can be any (not necessarily uniform) discrete random variable
- Distribution induced by the code is

$$P_{X^n Y^n}(x^n, y^n) := \sum_w P_{W_n}(w) P_{X^n|W_n}(x^n|w) P_{Y^n|W_n}(y^n|w).$$

- Require

$$P_{X^n Y^n} = \pi_{XY}^n \quad \text{for some } n \in \mathbb{N}.$$

# Exact Common Information

Asymptotic rate induced by the code is

$$\lim_{n \rightarrow \infty} \frac{H(W_n)}{n}$$

# Exact Common Information

Asymptotic rate induced by the code is

$$\lim_{n \rightarrow \infty} \frac{H(W_n)}{n}$$

- Compress  $W_n$  by a prefix-free, zero-error **variable-length code** (e.g., Shannon-Fano or Huffman code)

$$f : \mathcal{W}_n \rightarrow \{0, 1\}^* := \bigcup_{n \geq 1} \{0, 1\}^n$$

- Let the length of  $W_n$  be  $\ell(W_n)$ .

# Exact Common Information

Asymptotic rate induced by the code is

$$\lim_{n \rightarrow \infty} \frac{H(W_n)}{n}$$

- Compress  $W_n$  by a prefix-free, zero-error **variable-length code** (e.g., Shannon-Fano or Huffman code)

$$f : \mathcal{W}_n \rightarrow \{0, 1\}^* := \bigcup_{n \geq 1} \{0, 1\}^n$$

- Let the length of  $W_n$  be  $\ell(W_n)$ .
- Then, by **Shannon's zero-error compression theorem**, the optimal expected codeword length  $L(W_n) = \mathbb{E}[\ell(W_n)]$  satisfies

$$H(W_n) \leq L(W_n) < H(W_n) + 1$$

which implies that

$$\lim_{n \rightarrow \infty} \frac{L(W_n)}{n} = \lim_{n \rightarrow \infty} \frac{H(W_n)}{n}.$$

# Exact Common Information

## Definition

The exact common information is defined as

$$T_{\text{Ex}}(\pi_{XY}) := \inf \left\{ \lim_{n \rightarrow \infty} \frac{L(W_n)}{n} : P_{X^n Y^n} = \pi_{XY}^n \text{ for some } n \geq 1 \right\}$$

# Exact Common Information

## Definition

The exact common information is defined as

$$T_{\text{Ex}}(\pi_{XY}) := \inf \left\{ \lim_{n \rightarrow \infty} \frac{L(W_n)}{n} : P_{X^n Y^n} = \pi_{XY}^n \text{ for some } n \geq 1 \right\}$$

## Theorem ([Kumar et al., 2014])

$$T_{\text{Ex}}(\pi_{XY}) = \lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n).$$

# Exact Common Information

## Definition

The exact common information is defined as

$$T_{\text{Ex}}(\pi_{XY}) := \inf \left\{ \lim_{n \rightarrow \infty} \frac{L(W_n)}{n} : P_{X^n Y^n} = \pi_{XY}^n \text{ for some } n \geq 1 \right\}$$

## Theorem ([Kumar et al., 2014])

$$T_{\text{Ex}}(\pi_{XY}) = \lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n).$$

- Multi-letter characterization!

# Exact Common Information

## Definition

The exact common information is defined as

$$T_{\text{Ex}}(\pi_{XY}) := \inf \left\{ \lim_{n \rightarrow \infty} \frac{L(W_n)}{n} : P_{X^n Y^n} = \pi_{XY}^n \text{ for some } n \geq 1 \right\}$$

## Theorem ([Kumar et al., 2014])

$$T_{\text{Ex}}(\pi_{XY}) = \lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n).$$

- Multi-letter characterization!
- Exact CI  $\geq$  Wyner's CI

# Exact Common Information

## Definition

The exact common information is defined as

$$T_{\text{Ex}}(\pi_{XY}) := \inf \left\{ \lim_{n \rightarrow \infty} \frac{L(W_n)}{n} : P_{X^n Y^n} = \pi_{XY}^n \text{ for some } n \geq 1 \right\}$$

## Theorem ([Kumar et al., 2014])

$$T_{\text{Ex}}(\pi_{XY}) = \lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n).$$

- Multi-letter characterization!
- Exact CI  $\geq$  Wyner's CI
- Exact CI  $>$  Wyner's CI?
- Open problem posed by [Kumar et al., 2014]

# Exact Common Information

## Definition

The exact common information is defined as

$$T_{\text{Ex}}(\pi_{XY}) := \inf \left\{ \lim_{n \rightarrow \infty} \frac{L(W_n)}{n} : P_{X^n Y^n} = \pi_{XY}^n \text{ for some } n \geq 1 \right\}$$

## Theorem ([Kumar et al., 2014])

$$T_{\text{Ex}}(\pi_{XY}) = \lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n).$$

- Multi-letter characterization!
- Exact CI  $\geq$  Wyner's CI
- Exact CI  $>$  Wyner's CI?
- Open problem posed by [Kumar et al., 2014]

As expected the exact common information rate is greater than or equal to the Wyner common information.

### Proposition 3.

$$\bar{G}(X; Y) \geq J(X; Y).$$

In the following section, we show that they are equal for the SBES in Example 1. We do not know if this is the case in general, however.

From [Kumar et al., 2014]

# Exact Common Information

## Definition

The exact common information is defined as

$$T_{\text{Ex}}(\pi_{XY}) := \inf \left\{ \lim_{n \rightarrow \infty} \frac{L(W_n)}{n} : P_{X^n Y^n} = \pi_{XY}^n \text{ for some } n \geq 1 \right\}$$

## Theorem ([Kumar et al., 2014])

$$T_{\text{Ex}}(\pi_{XY}) = \lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n).$$

- Multi-letter characterization!
- Exact CI  $\geq$  Wyner's CI
- Exact CI  $>$  Wyner's CI?
- Open problem posed by [Kumar et al., 2014]

the exact common information rate. While this multiletter characterization is in general greater than or equal to the Wyner common information, we showed that they are equal for the SBES. The main open question is whether the exact common information rate has a single letter characterization in general. Is it always equal to the Wyner common information? Is there an example 2-DMS for which the exact common information rate is strictly larger than the Wyner common information? It would also be interesting to further explore the application to machine learning.

From [Kumar et al., 2014]

# Surprising Equivalence: $\infty$ -Rényi CI and Exact CI

Theorem ([Yu and Tan, 2020c])

For a bivariate source  $\pi_{XY}$  on a finite alphabet,


$$T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_\infty(\pi_{XY}).$$

# Surprising Equivalence: $\infty$ -Rényi CI and Exact CI

Theorem ([Yu and Tan, 2020c])

For a bivariate source  $\pi_{XY}$  on a finite alphabet,

$$T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_\infty(\pi_{XY}).$$




# Surprising Equivalence: $\infty$ -Rényi CI and Exact CI

Theorem ([Yu and Tan, 2020c])

For a bivariate source  $\pi_{XY}$  on a finite alphabet,

$$T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_\infty(\pi_{XY}).$$




# Surprising Equivalence: $\infty$ -Rényi CI and Exact CI

Theorem ([Yu and Tan, 2020c])

For a bivariate source  $\pi_{XY}$  on a finite alphabet,

$$T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_\infty(\pi_{XY}).$$




# Surprising Equivalence: $\infty$ -Rényi CI and Exact CI

Theorem ([Yu and Tan, 2020c])

For a bivariate source  $\pi_{XY}$  on a finite alphabet,

$$T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_\infty(\pi_{XY}).$$




## Surprising Equivalence: $\infty$ -Rényi CI and Exact CI

### Theorem ([Yu and Tan, 2020c])

For a bivariate source  $\pi_{XY}$  on a finite alphabet,

$$T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_\infty(\pi_{XY}).$$



# Proof of $\implies$ Part of Equivalence Theorem

Lemma ([Kumar et al., 2014], [Vellambi and Kliewer, 2016])

$\exists$  rate- $R$   $\infty$ -Rényi CI code  $\implies \exists$  rate- $R$  Exact CI code

# Proof of $\implies$ Part of Equivalence Theorem

Lemma ([Kumar et al., 2014], [Vellambi and Kliewer, 2016])

$\exists$  rate- $R$   $\infty$ -Rényi CI code  $\implies \exists$  rate- $R$  Exact CI code

- $\exists$  rate- $R$   $\infty$ -Rényi CI code

$$D_{\infty}(P_{X^n Y^n} \| \pi_{XY}^n) < \epsilon \implies P_{X^n Y^n}(x^n, y^n) < 2^{\epsilon} \pi_{XY}^n(x^n, y^n)$$

# Proof of $\implies$ Part of Equivalence Theorem

Lemma ([Kumar et al., 2014], [Vellambi and Kliewer, 2016])

$\exists$  rate- $R$   $\infty$ -Rényi CI code  $\implies \exists$  rate- $R$  Exact CI code

- $\exists$  rate- $R$   $\infty$ -Rényi CI code

$$D_{\infty}(P_{X^n Y^n} \| \pi_{XY}^n) < \epsilon \implies P_{X^n Y^n}(x^n, y^n) < 2^{\epsilon} \pi_{XY}^n(x^n, y^n)$$

- Define

$$\hat{P}_{X^n Y^n}(x^n, y^n) := \frac{2^{\epsilon} \pi_{XY}^n(x^n, y^n) - P_{X^n Y^n}(x^n, y^n)}{2^{\epsilon} - 1},$$

then obviously,  $\hat{P}_{X^n Y^n}(x^n, y^n)$  is a valid distribution.

# Proof of $\implies$ Part of Equivalence Theorem

Lemma ([Kumar et al., 2014], [Vellambi and Kliewer, 2016])

$\exists$  rate- $R$   $\infty$ -Rényi CI code  $\implies \exists$  rate- $R$  Exact CI code

- $\exists$  rate- $R$   $\infty$ -Rényi CI code

$$D_\infty(P_{X^n Y^n} \| \pi_{XY}^n) < \epsilon \implies P_{X^n Y^n}(x^n, y^n) < 2^\epsilon \pi_{XY}^n(x^n, y^n)$$

- Define

$$\hat{P}_{X^n Y^n}(x^n, y^n) := \frac{2^\epsilon \pi_{XY}^n(x^n, y^n) - P_{X^n Y^n}(x^n, y^n)}{2^\epsilon - 1},$$

then obviously,  $\hat{P}_{X^n Y^n}(x^n, y^n)$  is a valid distribution.

- Hence  $\pi_{XY}^n$  can be written as a mixture distribution

$$\pi_{XY}^n(x^n, y^n) = 2^{-\epsilon} P_{X^n Y^n}(x^n, y^n) + (1 - 2^{-\epsilon}) \hat{P}_{X^n Y^n}(x^n, y^n)$$

# Proof of $\implies$ Part of Equivalence Theorem

$$\pi_{XY}^n(x^n, y^n) = 2^{-\epsilon} P_{X^n Y^n}(x^n, y^n) + (1 - 2^{-\epsilon}) \hat{P}_{X^n Y^n}(x^n, y^n)$$

- A time-sharing variable-length scheme:

- ▶ The encoder first generates  $U \sim \text{Bern}(2^{-\epsilon})$ , and transmits it to two generators using 1 bit
- ▶ If  $U = 1$ , then the encoder and two generators use the rate- $R$   $\infty$ -Rényi CI code to generate  $P_{X^n Y^n}$
- ▶ If  $U = 0$ , then the encoder generates  $(X^n, Y^n) \sim \hat{P}_{X^n Y^n}$ , and compresses it with rate  $\log(|\mathcal{X}||\mathcal{Y}|)$  to generate  $\hat{P}_{X^n Y^n}$

# Proof of $\implies$ Part of Equivalence Theorem

$$\pi_{XY}^n(x^n, y^n) = 2^{-\epsilon} P_{X^n Y^n}(x^n, y^n) + (1 - 2^{-\epsilon}) \hat{P}_{X^n Y^n}(x^n, y^n)$$

- A time-sharing variable-length scheme:
  - ▶ The encoder first generates  $U \sim \text{Bern}(2^{-\epsilon})$ , and transmits it to two generators using 1 bit
  - ▶ If  $U = 1$ , then the encoder and two generators use the rate- $R$   $\infty$ -Rényi CI code to generate  $P_{X^n Y^n}$
  - ▶ If  $U = 0$ , then the encoder generates  $(X^n, Y^n) \sim \hat{P}_{X^n Y^n}$ , and compresses it with rate  $\log(|\mathcal{X}||\mathcal{Y}|)$  to generate  $\hat{P}_{X^n Y^n}$
- The induced distribution is  $\pi_{XY}^n$  exactly

# Proof of $\implies$ Part of Equivalence Theorem

$$\pi_{XY}^n(x^n, y^n) = 2^{-\epsilon} P_{X^n Y^n}(x^n, y^n) + (1 - 2^{-\epsilon}) \hat{P}_{X^n Y^n}(x^n, y^n)$$

- A time-sharing variable-length scheme:
  - ▶ The encoder first generates  $U \sim \text{Bern}(2^{-\epsilon})$ , and transmits it to two generators using 1 bit
  - ▶ If  $U = 1$ , then the encoder and two generators use the rate- $R$   $\infty$ -Rényi CI code to generate  $P_{X^n Y^n}$
  - ▶ If  $U = 0$ , then the encoder generates  $(X^n, Y^n) \sim \hat{P}_{X^n Y^n}$ , and compresses it with rate  $\log(|\mathcal{X}||\mathcal{Y}|)$  to generate  $\hat{P}_{X^n Y^n}$
- The induced distribution is  $\pi_{XY}^n$  exactly
- The total code rate

$$\leq \frac{1}{n} + 2^{-\epsilon} R + (1 - 2^{-\epsilon}) \log(|\mathcal{X}||\mathcal{Y}|) \rightarrow R$$

as  $n \rightarrow \infty, \epsilon \rightarrow 0$

# Proof of $\Leftarrow$ Part of Equivalence Theorem

## Lemma

$\exists$  rate- $R$   $\infty$ -Rényi CI code  $\Leftarrow \exists$  rate- $R$  Exact CI code

# Proof of $\Leftarrow$ Part of Equivalence Theorem

## Lemma

$\exists$  rate- $R$   $\infty$ -Rényi CI code  $\Leftarrow \exists$  rate- $R$  Exact CI code

- Let  $\{(P_{W_k}, P_{X^k|W_k}, P_{Y^k|W_k})\}_{k \in \mathbb{N}}$  be rate- $R$  exact CI codes such that

$$\lim_{k \rightarrow \infty} \frac{1}{k} H(P_{W_k}) = R$$

but  $W_k$  is not uniform.



# Proof of $\Leftarrow$ Part of Equivalence Theorem

## Lemma

$\exists$  rate- $R$   $\infty$ -Rényi CI code  $\Leftarrow \exists$  rate- $R$  Exact CI code


- Let  $\{(P_{W_k}, P_{X^k|W_k}, P_{Y^k|W_k})\}_{k \in \mathbb{N}}$  be rate- $R$  exact CI codes such that

$$\lim_{k \rightarrow \infty} \frac{1}{k} H(P_{W_k}) = R$$


but  $W_k$  is not uniform.




- Simulate  $W_k^n$  using two Rényi source resolvability codes!



# Proof of $\Leftarrow$ Part of Equivalence Theorem




# Proof of $\Leftarrow$ Part of Equivalence Theorem




Succeed in the sense of  $D_{\infty}(P_{f(M)} \| P_{W_k}^n) \rightarrow 0$  if [Yu and Tan, 2019]

$$R > \frac{1}{k} H(P_{W_k})$$

# Proof of $\Leftarrow$ Part of Equivalence Theorem



# Proof of $\Leftarrow$ Part of Equivalence Theorem




- For the given stochastic kernel (channel)  $P_{X^k|W_k}^n P_{Y^k|W_k}^n$ ,

$$P_W^n \longrightarrow P_{X^k|W_k}^n P_{Y^k|W_k}^n \longrightarrow \pi_{XY}^{kn}$$

$$P_{f(M)} \longrightarrow P_{X^k|W_k}^n P_{Y^k|W_k}^n \longrightarrow P_{X^{kn}Y^{kn}}$$

# Proof of $\Leftarrow$ Part of Equivalence Theorem



- For the given stochastic kernel (channel)  $P_{X^k|W_k}^n P_{Y^k|W_k}^n$ ,

$$P_W^n \longrightarrow P_{X^k|W_k}^n P_{Y^k|W_k}^n \longrightarrow \pi_{XY}^{kn}$$

$$P_{f(M)} \longrightarrow P_{X^k|W_k}^n P_{Y^k|W_k}^n \longrightarrow P_{X^{kn}Y^{kn}}$$

- By the data processing inequality (DPI) for Rényi divergence,

$$D_\infty(P_{X^{kn}Y^{kn}} \| \pi_{XY}^{kn}) \leq D_\infty(P_{f(M)} \| P_{W_k}^n) \xrightarrow{n \rightarrow \infty} 0$$

# Combining with Single-Letter Bounds from Rényi CI

Theorem ([Yu and Tan, 2020c])

For  $(X, Y) \sim \pi_{XY}$  on a finite alphabet,

$$\underline{\Gamma}_{\infty}(\pi_{XY}) \leq T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_{\infty}(\pi_{XY}) \leq \overline{\Gamma}_{\infty}(\pi_{XY}).$$

# Combining with Single-Letter Bounds from Rényi CI

## Theorem ([Yu and Tan, 2020c])

For  $(X, Y) \sim \pi_{XY}$  on a finite alphabet,

$$\underline{\Gamma}_\infty(\pi_{XY}) \leq T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_\infty(\pi_{XY}) \leq \bar{\Gamma}_\infty(\pi_{XY}).$$

- Gone from a **multi-letter expression** by [Kumar et al., 2014]

$$\lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n)$$

to **single-letter bounds**.

# Combining with Single-Letter Bounds from Rényi CI

## Theorem ([Yu and Tan, 2020c])

For  $(X, Y) \sim \pi_{XY}$  on a finite alphabet,

$$\underline{\Gamma}_{\infty}(\pi_{XY}) \leq T_{\text{Ex}}(\pi_{XY}) = \tilde{T}_{\infty}(\pi_{XY}) \leq \bar{\Gamma}_{\infty}(\pi_{XY}).$$


- Gone from a **multi-letter expression** by [Kumar et al., 2014]

$$\lim_{n \rightarrow \infty} \frac{1}{n} \min_{\substack{P_{W_n} P_{X^n|W_n} P_{Y^n|W_n}: \\ P_{X^n Y^n} = \pi_{XY}^n}} H(W_n)$$



to **single-letter bounds**.

- Presumably the bounds are more amenable to numerical evaluation?



# Revisiting the DBSS



# Revisiting the DBSS



# Revisiting the DBSS




## Theorem (Evaluation of Upper and Lower Bounds for $\text{DSBS}(p)$ )

For a DSBS  $(X, Y) \sim \text{DSBS}(p)$  with crossover probability  $p \in (0, 1/2)$ ,


$$\begin{aligned}\tilde{T}_\infty(\pi_{XY}) &= T_{\text{Ex}}(\pi_{XY}) \\ &= -2h(a) - (1-2a)\log\left[\frac{1}{2}(a^2 + (1-a)^2)\right] - 2a\log[a(1-a)],\end{aligned}$$

where  $a := \frac{1-\sqrt{1-2p}}{2} \in (0, \frac{1}{2})$  and  $h(a) := -a\log a - (1-a)\log(1-a)$ .

# Numerical Results — DSBS



# Numerical Results — DSBS




$$T_{\text{Ex}}(\text{DSBS}(p)) > C_W(\text{DSBS}(p)) \quad \forall p \in (0, 1/2).$$


Answers the open question in [Kumar et al., 2014].

# Why is Exact CI (or $\infty$ -Rényi CI) > Wyner's CI?

# Why is Exact CI (or $\infty$ -Rényi CI) > Wyner's CI?




# Why is Exact CI (or $\infty$ -Rényi CI) > Wyner's CI?




Wyner's common information requires

$$\frac{P_{X^n Y^n}(x^n, y^n)}{\pi_{XY}^n(x^n, y^n)} = 1 + o(1) \quad \text{for almost all } (x^n, y^n) \in \mathcal{T}_\epsilon^{(n)}(\pi_{XY})$$

# Why is Exact CI (or $\infty$ -Rényi CI) > Wyner's CI?




# Why is Exact CI (or $\infty$ -Rényi CI) > Wyner's CI?



Rényi CI of order  $\infty$  or Exact CI requires

$$\frac{P_{X^n Y^n}(x^n, y^n)}{\pi_{XY}^n(x^n, y^n)} = 1 + o(1) \quad \text{for all } (x^n, y^n) \in \text{supp}(P_{X^n Y^n})$$


# Why is Exact CI (or $\infty$ -Rényi CI) > Wyner's CI?



Rényi CI of order  $\infty$  or Exact CI requires

$$\frac{P_{X^n Y^n}(x^n, y^n)}{\pi_{XY}^n(x^n, y^n)} = 1 + o(1) \quad \text{for all } (x^n, y^n) \in \bigcup_w \text{supp}\left(P_{X^n|W_n}(\cdot|w)P_{Y^n|W_n}(\cdot|w)\right)$$

# Why is Exact CI (or $\infty$ -Rényi CI) > Wyner's CI?




Rényi CI of order  $\infty$  or Exact CI requires


$$\frac{P_{X^n Y^n}(x^n, y^n)}{\pi_{XY}^n(x^n, y^n)} = 1 + o(1) \quad \text{for all } (x^n, y^n) \in \bigcup_w \text{supp}\left(P_{X^n|W_n}(\cdot|w)P_{Y^n|W_n}(\cdot|w)\right)$$

# When is Exact CI (or $\infty$ -Rényi CI) = Wyner's CI?

# When is Exact CI (or $\infty$ -Rényi CI) = Wyner's CI?




# When is Exact CI (or $\infty$ -Rényi CI) = Wyner's CI?



Sufficient Condition [Vellambi and Kliewer, 2016]


# When is Exact CI (or $\infty$ -Rényi CI) = Wyner's CI?



Sufficient Condition [Vellambi and Kliewer, 2016]

$$H(X|W=w)H(Y|W=w) = 0 \quad \text{for each } w$$

# When is Exact CI (or $\infty$ -Rényi CI) = Wyner's CI?




Sufficient Condition [Vellambi and Kliewer, 2016]

$$H(X|W=w)H(Y|W=w) = 0 \quad \text{for each } w$$

$$\iff \mathcal{C}(P_{X|W}, P_{Y|W}) = \{P_{X|W}P_{Y|W}\}$$


# When is Exact CI (or $\infty$ -Rényi CI) = Wyner's CI?



Sufficient Condition [Vellambi and Kliewer, 2016]

$$\begin{aligned} & H(X|W=w)H(Y|W=w) = 0 \quad \text{for each } w \\ \iff & \mathcal{C}(P_{X|W}, P_{Y|W}) = \{P_{X|W}P_{Y|W}\} \\ \iff & \mathcal{T}_{\epsilon}^{(n)}(P_{XY}) \approx \bigcup_{w^n \in \mathcal{C}} \left( \mathcal{T}_{\epsilon}^{(n)}(P_{XW}|w^n) \times \mathcal{T}_{\epsilon}^{(n)}(P_{YW}|w^n) \right) \end{aligned}$$

# When is Exact CI (or $\infty$ -Rényi CI) = Wyner's CI?



Sufficient Condition [Vellambi and Kliewer, 2016]

$$H(X|W=w)H(Y|W=w) = 0 \quad \text{for each } w$$

$$\iff \mathcal{C}(P_{X|W}, P_{Y|W}) = \{P_{X|W}P_{Y|W}\}$$

$$\iff \mathcal{T}_{\epsilon}^{(n)}(P_{XY}) \approx \bigcup_{w^n \in \mathcal{C}} \left( \mathcal{T}_{\epsilon}^{(n)}(P_{XW}|w^n) \times \mathcal{T}_{\epsilon}^{(n)}(P_{YW}|w^n) \right)$$

$$\iff \mathcal{T}_{\epsilon}^{(n)}(P_{XY}) \approx \text{supp}(P_{X^n Y^n}) \quad (\text{No type overflow})$$

# When Exact CI (or $\infty$ -Rényi CI) = Wyner's CI

Example for Sufficient Condition:



$$H(X|W=w)H(Y|W=w) = 0 \quad \forall w$$

# When Exact CI (or $\infty$ -Rényi CI) = Wyner's CI

Example for Sufficient Condition:

$$H(X|W=w)H(Y|W=w) = 0 \quad \forall w$$

- Symmetric Binary Erasure Source (SBES)





# When Exact CI (or $\infty$ -Rényi CI) = Wyner's CI

Example for Sufficient Condition:

$$H(X|W=w)H(Y|W=w) = 0 \quad \forall w$$

- Symmetric Binary Erasure Source (SBES)




- $(1 - p_1)(1 - p_2) = 1 - p$ .

# When Exact CI (or $\infty$ -Rényi CI) = Wyner's CI

Example for Sufficient Condition:

$$H(X|W=w)H(Y|W=w) = 0 \quad \forall w$$

- Symmetric Binary Erasure Source (SBES)



- $(1 - p_1)(1 - p_2) = 1 - p$ .
- The Exact CI is equal to Wyner's CI and

$$\tilde{T}_\infty(\pi_{XY}) = T_{\text{Exact}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY}) = \begin{cases} 1 & p \leq \frac{1}{2} \\ H(p) & p > \frac{1}{2} \end{cases}.$$


# Outline

- 1 Introduction: Measures of Information Among Two Random Variables
- 2 Wyner's Common Information
- 3 Rényi Common Information
- 4 Exact Common Information
- 5 Approximate and Exact Channel Synthesis
- 6 Nonnegative Matrix Factorization and Nonnegative Rank
- 7 Extensions of Gács–Körner–Witsenhausen's Common Information (L. Yu)

# Channel Synthesis

- Given  $\pi_{XY} = \pi_X \pi_{Y|X}$  consider the following task:


$$K_n \sim \text{Unif}[2^{nR_0}] \quad (\text{Shared Key})$$



# Channel Synthesis

- Given  $\pi_{XY} = \pi_X \pi_{Y|X}$  consider the following task:

$$K_n \sim \text{Unif}[2^{nR_0}] \quad (\text{Shared Key})$$




- Goal: Ensure that

$$P_{X^n Y^n} \approx \pi_{XY}^n \text{ (Approximate)} \quad \text{or} \quad P_{X^n Y^n} = \pi_{XY}^n \text{ (Exact).}$$

# Channel Synthesis

- Given  $\pi_{XY} = \pi_X \pi_{Y|X}$  consider the following task:

$$K_n \sim \text{Unif}[2^{nR_0}] \quad (\text{Shared Key})$$



- Goal: Ensure that

$$P_{X^n Y^n} \approx \pi_{XY}^n \text{ (Approximate)} \quad \text{or} \quad P_{X^n Y^n} = \pi_{XY}^n \text{ (Exact)}.$$


- Equivalently,

$$P_{Y^n|X^n} \approx \pi_{Y|X}^n \text{ (Approximate)} \quad \text{or} \quad P_{Y^n|X^n} = \pi_{Y|X}^n \text{ (Exact)}.$$

# Channel Synthesis

- Given  $\pi_{XY} = \pi_X \pi_{Y|X}$  consider the following task:

$$K_n \sim \text{Unif}[2^{nR_0}] \quad (\text{Shared Key})$$



- Goal: Ensure that

$$P_{X^n Y^n} \approx \pi_{XY}^n \text{ (Approximate)} \quad \text{or} \quad P_{X^n Y^n} = \pi_{XY}^n \text{ (Exact)}.$$

- Equivalently,

$$P_{Y^n | X^n} \approx \pi_{Y|X}^n \text{ (Approximate)} \quad \text{or} \quad P_{Y^n | X^n} = \pi_{Y|X}^n \text{ (Exact)}.$$

- Known as **channel synthesis** [Cuff, 2012].

# Approximate Channel Synthesis

- Consider approximate channel synthesis under TV criterion, i.e.,

$$\lim_{n \rightarrow \infty} |P_{X^n Y^n} - \pi_{XY}^n| = 0.$$

# Approximate Channel Synthesis

- Consider approximate channel synthesis under TV criterion, i.e.,

$$\lim_{n \rightarrow \infty} |P_{X^n Y^n} - \pi_{XY}^n| = 0.$$


- Let the region of achievable rate pairs  $(R, R_0)$  be  $\mathcal{R}_W(\pi_{XY})$ .

# Approximate Channel Synthesis

- Consider approximate channel synthesis under TV criterion, i.e.,

$$\lim_{n \rightarrow \infty} |P_{X^n Y^n} - \pi_{XY}^n| = 0.$$

- Let the region of achievable rate pairs  $(R, R_0)$  be  $\mathcal{R}_W(\pi_{XY})$ .
- When  $R_0 = 0$ , problem reduces to approximate distributed source simulation



so the minimum compression rate is Wyner's common information


$$R^*(R_0 = 0 | \pi_{XY}) = C_W(\pi_{XY})$$

# Approximate Channel Synthesis

- Consider approximate channel synthesis under TV criterion, i.e.,

$$\lim_{n \rightarrow \infty} |P_{X^n Y^n} - \pi_{XY}^n| = 0.$$

- Let the region of achievable rate pairs  $(R, R_0)$  be  $\mathcal{R}_W(\pi_{XY})$ .
- When  $R_0 = 0$ , problem reduces to approximate distributed source simulation



so the minimum compression rate is Wyner's common information

$$R^*(R_0 = 0 | \pi_{XY}) = C_W(\pi_{XY})$$

- When  $R_0 = \infty$ ,

$$R^*(R_0 = \infty | \pi_{XY}) = I_\pi(X; Y)$$

# Approximate Channel Synthesis


It was shown in [Cuff, 2012] that

$$\mathcal{R}_W(\pi_{XY}) := \bigcup_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} \left\{ (R, R_0) : \begin{array}{l} R \geq I(X; W) \\ R + R_0 \geq I(XY; W) \end{array} \right\}.$$

# Approximate Channel Synthesis


It was shown in [Cuff, 2012] that

$$\mathcal{R}_W(\pi_{XY}) := \bigcup_{\substack{P_W P_{X|W} P_{Y|W}: \\ P_{XY} = \pi_{XY}}} \left\{ (R, R_0) : \begin{array}{l} R \geq I(X; W) \\ R + R_0 \geq I(XY; W) \end{array} \right\}.$$




# Exact Channel Synthesis

$K_n \sim \text{Unif}[2^{nR_0}]$  (Shared Key)



# Exact Channel Synthesis




- Now, similarly to **exact common information**, we demand that

$$P_{X^n Y^n} = \pi_{XY}^n \text{ for some large enough } n \in \mathbb{N}$$

but just like exact CI, we allow **variable-length codes** for  $W_n$ .

# Exact Channel Synthesis

$$K_n \sim \text{Unif}[2^{nR_0}] \quad (\text{Shared Key})$$



- Now, similarly to **exact common information**, we demand that


$$P_{X^n Y^n} = \pi_{XY}^n \text{ for some large enough } n \in \mathbb{N}$$

but just like exact CI, we allow **variable-length codes** for  $W_n$ .

- If  $R_0 = \infty$ , [Bennett et al., 2002] showed that the minimum  $R$  is  $I(X; Y)$ .

# Exact Channel Synthesis

$$K_n \sim \text{Unif}[2^{nR_0}] \quad (\text{Shared Key})$$




- Now, similarly to **exact common information**, we demand that

$$P_{X^n Y^n} = \pi_{XY}^n \text{ for some large enough } n \in \mathbb{N}$$


but just like exact CI, we allow **variable-length codes** for  $W_n$ .

- If  $R_0 = \infty$ , [Bennett et al., 2002] showed that the minimum  $R$  is  $I(X; Y)$ .
- Best **tradeoff between  $R$  and  $R_0$**  in the non-extremal cases considered by [Yu and Tan, 2020b].

# Doubly Binary Symmetric Sources




# Doubly Binary Symmetric Sources




Exact channel synthesis region is **strictly smaller than**  $\mathcal{R}_W(\pi_{XY})$

# Doubly Binary Symmetric Sources



Exact channel synthesis region is **strictly smaller than**  $\mathcal{R}_W(\pi_{XY})$

# Doubly Binary Symmetric Sources



Exact channel synthesis region is strictly smaller than  $\mathcal{R}_W(\pi_{XY})$

# Outline

- 1 Introduction: Measures of Information Among Two Random Variables
- 2 Wyner's Common Information
- 3 Rényi Common Information
- 4 Exact Common Information
- 5 Approximate and Exact Channel Synthesis
- 6 Nonnegative Matrix Factorization and Nonnegative Rank
- 7 Extensions of Gäcs–Körner–Witsenhausen's Common Information (L. Yu)

# Nonnegative Matrix Factorization

- Given a matrix  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , find  $\mathbf{U} \in \mathbb{R}_+^{m \times r}$  and  $\mathbf{V} \in \mathbb{R}_+^{r \times k}$  such that

$$\mathbf{M} \approx \mathbf{UV} \quad \text{or} \quad \mathbf{M} = \mathbf{UV}.$$

Many applications. See [Cichocki et al., 2009] or [Gillis, 2020].


# Nonnegative Matrix Factorization

- Given a matrix  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , find  $\mathbf{U} \in \mathbb{R}_+^{m \times r}$  and  $\mathbf{V} \in \mathbb{R}_+^{r \times k}$  such that

$$\mathbf{M} \approx \mathbf{UV} \quad \text{or} \quad \mathbf{M} = \mathbf{UV}.$$

Many applications. See [Cichocki et al., 2009] or [Gillis, 2020].

- Dimensionality reduction:




# Nonnegative Matrix Factorization

- Given a matrix  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , find  $\mathbf{U} \in \mathbb{R}_+^{m \times r}$  and  $\mathbf{V} \in \mathbb{R}_+^{r \times k}$  such that

$$\mathbf{M} \approx \mathbf{UV} \quad \text{or} \quad \mathbf{M} = \mathbf{UV}.$$

Many applications. See [Cichocki et al., 2009] or [Gillis, 2020].

- Dimensionality reduction:




# Nonnegative Matrix Factorization

- Given a matrix  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , find  $\mathbf{U} \in \mathbb{R}_+^{m \times r}$  and  $\mathbf{V} \in \mathbb{R}_+^{r \times k}$  such that

$$\mathbf{M} \approx \mathbf{UV} \quad \text{or} \quad \mathbf{M} = \mathbf{UV}.$$

Many applications. See [Cichocki et al., 2009] or [Gillis, 2020].

- Dimensionality reduction:



- Only interested in **exact** factorization.


# Nonnegative Matrix Factorization

- Given a matrix  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , find  $\mathbf{U} \in \mathbb{R}_+^{m \times r}$  and  $\mathbf{V} \in \mathbb{R}_+^{r \times k}$  such that

$$\mathbf{M} \approx \mathbf{UV} \quad \text{or} \quad \mathbf{M} = \mathbf{UV}.$$

Many applications. See [Cichocki et al., 2009] or [Gillis, 2020].

- Dimensionality reduction:



- Only interested in **exact** factorization.
- What is the **minimum  $r$**  to achieve exact factorization? Is this connected to **information theory**?

# Nonnegative Rank

## Definition

The **nonnegative rank** of  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , denoted as  $\text{rank}_+(\mathbf{M})$ , is the **smallest integer  $r$**  such that

$$\mathbf{M} = \sum_{w=1}^r \mathbf{u}_w \mathbf{v}_w^\top$$

for some **nonnegative vectors**  $\mathbf{u}_w \in \mathbb{R}_+^m$  and  $\mathbf{v}_w \in \mathbb{R}_+^k$ .

# Nonnegative Rank

## Definition

The **nonnegative rank** of  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , denoted as  $\text{rank}_+(\mathbf{M})$ , is the **smallest integer  $r$**  such that

$$\mathbf{M} = \sum_{w=1}^r \mathbf{u}_w \mathbf{v}_w^\top$$

for some **nonnegative vectors**  $\mathbf{u}_w \in \mathbb{R}_+^m$  and  $\mathbf{v}_w \in \mathbb{R}_+^k$ .

- Obviously,  $\text{rank}(\mathbf{M}) \leq \text{rank}_+(\mathbf{M})$

# Nonnegative Rank

## Definition

The **nonnegative rank** of  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , denoted as  $\text{rank}_+(\mathbf{M})$ , is the **smallest integer  $r$**  such that

$$\mathbf{M} = \sum_{w=1}^r \mathbf{u}_w \mathbf{v}_w^\top$$

for some **nonnegative vectors**  $\mathbf{u}_w \in \mathbb{R}_+^m$  and  $\mathbf{v}_w \in \mathbb{R}_+^k$ .

- Obviously,  $\text{rank}(\mathbf{M}) \leq \text{rank}_+(\mathbf{M})$
- Gap can be **large**. Fix  $\{a_1, \dots, a_m\} \subset \mathbb{R}$  and consider **distance matrix**

$$\mathbf{M} = \begin{bmatrix} 0 & (a_1 - a_2)^2 & (a_1 - a_3)^2 & \dots & (a_1 - a_m)^2 \\ (a_2 - a_1)^2 & 0 & (a_2 - a_3)^2 & \dots & (a_2 - a_m)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ (a_m - a_1)^2 & (a_m - a_2)^2 & (a_m - a_3)^2 & \dots & 0 \end{bmatrix}.$$

# Nonnegative Rank

## Definition

The **nonnegative rank** of  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$ , denoted as  $\text{rank}_+(\mathbf{M})$ , is the **smallest integer  $r$**  such that

$$\mathbf{M} = \sum_{w=1}^r \mathbf{u}_w \mathbf{v}_w^\top$$

for some **nonnegative vectors**  $\mathbf{u}_w \in \mathbb{R}_+^m$  and  $\mathbf{v}_w \in \mathbb{R}_+^k$ .

- Obviously,  $\text{rank}(\mathbf{M}) \leq \text{rank}_+(\mathbf{M})$
- Gap can be **large**. Fix  $\{a_1, \dots, a_m\} \subset \mathbb{R}$  and consider **distance matrix**

$$\mathbf{M} = \begin{bmatrix} a_1^2 & 1 & -2a_1 \\ a_2^2 & 1 & -2a_2 \\ \vdots & \ddots & \vdots \\ a_m^2 & 1 & -2a_m \end{bmatrix} \begin{bmatrix} 1 & 1 & \dots & 1 \\ a_1^2 & a_2^2 & \dots & a_m^2 \\ a_1 & a_2 & \dots & a_m \end{bmatrix}$$

- $\text{rank}(\mathbf{M}) \leq 3$ . [Beasley and Laffey, 2009] showed  $\text{rank}_+(\mathbf{M}) = \Omega(\log m)$ .

# Wyner's CI for Normalized Nonnegative Matrices

- Let  $\mathbf{M} \in \mathbb{R}_+^{r \times k}$  be a nonnegative matrix.

# Wyner's CI for Normalized Nonnegative Matrices

- Let  $\mathbf{M} \in \mathbb{R}_+^{r \times k}$  be a nonnegative matrix.
- We may define

$$\pi_{XY}(x, y) := \frac{M_{x,y}}{\|\mathbf{M}\|_1} \quad (x, y) \in [m] \times [k] = \mathcal{X} \times \mathcal{Y}.$$

# Wyner's CI for Normalized Nonnegative Matrices

- Let  $\mathbf{M} \in \mathbb{R}_+^{r \times k}$  be a nonnegative matrix.
- We may define

$$\pi_{XY}(x, y) := \frac{M_{x,y}}{\|\mathbf{M}\|_1} \quad (x, y) \in [m] \times [k] = \mathcal{X} \times \mathcal{Y}.$$

- A discrete r.v.  $W \in \mathcal{W}$  is a **seed** for  $\pi_{XY}$ , or equivalently  $\mathbf{M}$ , if

$$X - W - Y.$$

# Wyner's CI for Normalized Nonnegative Matrices

- Let  $\mathbf{M} \in \mathbb{R}_+^{r \times k}$  be a nonnegative matrix.
- We may define

$$\pi_{XY}(x, y) := \frac{M_{x,y}}{\|\mathbf{M}\|_1} \quad (x, y) \in [m] \times [k] = \mathcal{X} \times \mathcal{Y}.$$

- A discrete r.v.  $W \in \mathcal{W}$  is a **seed** for  $\pi_{XY}$ , or equivalently  $\mathbf{M}$ , if

$$X - W - Y.$$

- Every NMF of

$$\mathbf{M} = \sum_w \mathbf{u}_w \mathbf{v}_w^\top$$

induces a seed  $W$  for  $\mathbf{M}$ .

# Wyner's CI for Normalized Nonnegative Matrices

- Let  $\mathbf{M} \in \mathbb{R}_+^{r \times k}$  be a nonnegative matrix.
- We may define

$$\pi_{XY}(x, y) := \frac{M_{x,y}}{\|\mathbf{M}\|_1} \quad (x, y) \in [m] \times [k] = \mathcal{X} \times \mathcal{Y}.$$

- A discrete r.v.  $W \in \mathcal{W}$  is a **seed** for  $\pi_{XY}$ , or equivalently  $\mathbf{M}$ , if

$$X - W - Y.$$

- Every NMF of

$$\mathbf{M} = \sum_w \mathbf{u}_w \mathbf{v}_w^\top$$

induces a seed  $W$  for  $\mathbf{M}$ .

- Wyner's common information for  $\mathbf{M}$  is

$$C_W(\mathbf{M}) := C_W(\pi_{XY}).$$

# Playing With Definitions

Theorem ([Jain et al., 2013], [Braun and Pokutta, 2013])

$$C_W(\mathbf{M}) \leq \log \text{rank}_+(\mathbf{M}).$$

# Playing With Definitions

Theorem ([Jain et al., 2013], [Braun and Pokutta, 2013])

$$C_W(\mathbf{M}) \leq \log \text{rank}_+(\mathbf{M}).$$

Proof.

Let  $\mathbf{M}$  have an **optimal NMF**  $\mathbf{M} = \sum_w \mathbf{u}_w \mathbf{v}_w^\top$ . Define seed  $W$  as

$$P_{W|XY}(w|x, y) = \begin{cases} \frac{[\mathbf{u}_w]_x [\mathbf{v}_w]_y}{M_{x,y}} & M_{x,y} > 0 \\ \text{arbitrary} & M_{x,y} = 0 \end{cases}.$$

# Playing With Definitions

Theorem ([Jain et al., 2013], [Braun and Pokutta, 2013])

$$C_W(\mathbf{M}) \leq \log \text{rank}_+(\mathbf{M}).$$

Proof.

Let  $\mathbf{M}$  have an **optimal NMF**  $\mathbf{M} = \sum_w \mathbf{u}_w \mathbf{v}_w^\top$ . Define seed  $W$  as

$$P_{W|XY}(w|x, y) = \begin{cases} \frac{[\mathbf{u}_w]_x [\mathbf{v}_w]_y}{M_{x,y}} & M_{x,y} > 0 \\ \text{arbitrary} & M_{x,y} = 0 \end{cases}.$$

By Bayes rule,

$$P_{XY|W}(x, y|w) = \frac{[\mathbf{u}_w]_x [\mathbf{v}_w]_y}{\sum_{x',y'} [\mathbf{u}_w]_{x'} [\mathbf{v}_w]_{y'}} \quad (x, y) \in \mathcal{X} \times \mathcal{Y}.$$

# Playing With Definitions

Theorem ([Jain et al., 2013], [Braun and Pokutta, 2013])

$$C_W(\mathbf{M}) \leq \log \text{rank}_+(\mathbf{M}).$$

Proof.

Let  $\mathbf{M}$  have an **optimal NMF**  $\mathbf{M} = \sum_w \mathbf{u}_w \mathbf{v}_w^\top$ . Define seed  $W$  as

$$P_{W|XY}(w|x, y) = \begin{cases} \frac{[\mathbf{u}_w]_x [\mathbf{v}_w]_y}{M_{x,y}} & M_{x,y} > 0 \\ \text{arbitrary} & M_{x,y} = 0 \end{cases}.$$

By Bayes rule,

$$P_{XY|W}(x, y|w) = \frac{[\mathbf{u}_w]_x [\mathbf{v}_w]_y}{\sum_{x',y'} [\mathbf{u}_w]_{x'} [\mathbf{v}_w]_{y'}} \quad (x, y) \in \mathcal{X} \times \mathcal{Y}.$$

So,  $X - W - Y$  and

$$C_W(\mathbf{M}) \leq I_P(XY; W) \leq H(W) \leq \log |\mathcal{W}| = \log \text{rank}_+(\mathbf{M}).$$



# Gap Between $C_W(\mathbf{M})$ and $\log \text{rank}_+(\mathbf{M})$ ?

- Consider the diagonal matrix

$$\mathbf{M} = \frac{1}{\sum_{j=1}^m 2^j} \begin{bmatrix} 2^1 & 0 & 0 & \dots & 0 \\ 0 & 2^2 & 0 & \dots & 0 \\ 0 & 0 & 2^3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 2^m \end{bmatrix}.$$

# Gap Between $C_W(\mathbf{M})$ and $\log \text{rank}_+(\mathbf{M})$ ?

- Consider the diagonal matrix

$$\mathbf{M} = \frac{1}{\sum_{j=1}^m 2^j} \begin{bmatrix} 2^1 & 0 & 0 & \dots & 0 \\ 0 & 2^2 & 0 & \dots & 0 \\ 0 & 0 & 2^3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 2^m \end{bmatrix}.$$

- $\text{rank}_+(\mathbf{M}) = m$

# Gap Between $C_W(\mathbf{M})$ and $\log \text{rank}_+(\mathbf{M})$ ?

- Consider the diagonal matrix

$$\mathbf{M} = \frac{1}{\sum_{j=1}^m 2^j} \begin{bmatrix} 2^1 & 0 & 0 & \dots & 0 \\ 0 & 2^2 & 0 & \dots & 0 \\ 0 & 0 & 2^3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 2^m \end{bmatrix}.$$

- $\text{rank}_+(\mathbf{M}) = m$
- But

$$\begin{aligned} C_W(\mathbf{M}) &\leq H_\pi(XY) = H(\pi_X) \\ &= H\left(\frac{2}{\sum_{j \in [m]} 2^j}, \frac{2^2}{\sum_{j \in [m]} 2^j}, \dots, \frac{2^m}{\sum_{j \in [m]} 2^j}\right) \\ &= - \sum_{i \in [m]} \frac{2^i}{\sum_{j \in [m]} 2^j} \log\left(\frac{2^i}{\sum_{j \in [m]} 2^j}\right) \leq 2 \quad \forall m \in \mathbb{N}. \end{aligned}$$

# Gap Between $C_W(\mathbf{M})$ and $\log \text{rank}_+(\mathbf{M})$ ?

- Consider the diagonal matrix

$$\mathbf{M} = \frac{1}{\sum_{j=1}^m 2^j} \begin{bmatrix} 2^1 & 0 & 0 & \dots & 0 \\ 0 & 2^2 & 0 & \dots & 0 \\ 0 & 0 & 2^3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 2^m \end{bmatrix}.$$

- $\text{rank}_+(\mathbf{M}) = m$
- But

$$\begin{aligned} C_W(\mathbf{M}) &\leq H_\pi(XY) = H(\pi_X) \\ &= H\left(\frac{2}{\sum_{j \in [m]} 2^j}, \frac{2^2}{\sum_{j \in [m]} 2^j}, \dots, \frac{2^m}{\sum_{j \in [m]} 2^j}\right) \\ &= - \sum_{i \in [m]} \frac{2^i}{\sum_{j \in [m]} 2^j} \log\left(\frac{2^i}{\sum_{j \in [m]} 2^j}\right) \leq 2 \quad \forall m \in \mathbb{N}. \end{aligned}$$

- Gap can be arbitrarily large.

# Gap Between $C_W(\mathbf{M})$ and $\log \text{rank}_+(\mathbf{M})$ ?

- Consider the diagonal matrix

$$\mathbf{M} = \frac{1}{\sum_{j=1}^m 2^j} \begin{bmatrix} 2^1 & 0 & 0 & \dots & 0 \\ 0 & 2^2 & 0 & \dots & 0 \\ 0 & 0 & 2^3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 2^m \end{bmatrix}.$$

- $\text{rank}_+(\mathbf{M}) = m$
- But

$$\begin{aligned} C_W(\mathbf{M}) &\leq H_\pi(XY) = H(\pi_X) \\ &= H\left(\frac{2}{\sum_{j \in [m]} 2^j}, \frac{2^2}{\sum_{j \in [m]} 2^j}, \dots, \frac{2^m}{\sum_{j \in [m]} 2^j}\right) \\ &= - \sum_{i \in [m]} \frac{2^i}{\sum_{j \in [m]} 2^j} \log\left(\frac{2^i}{\sum_{j \in [m]} 2^j}\right) \leq 2 \quad \forall m \in \mathbb{N}. \end{aligned}$$

- Gap can be arbitrarily large.
- Is the relation between  $C_W(\mathbf{M})$  and  $\log \text{rank}_+(\mathbf{M})$  fundamental?

# Amortization Comes to the Rescue

## Theorem ([Braun et al., 2017])

Let  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$  be such that  $\|\mathbf{M}\|_1 = \sum_{x,y} M_{x,y} = 1$ . For any  $\epsilon, \delta > 0$ , if  $n \geq n_0(\epsilon, \delta, m, k, C_W(\mathbf{M}))$  is sufficiently large, there exists  $\mathbf{M}_{\epsilon, \delta, n} \in \mathbb{R}_+^{m^n \times k^n}$  with

$$\|\mathbf{M}^{\otimes n} - \mathbf{M}_{\epsilon, \delta, n}\|_1 \leq \delta.$$

and

$$\lim_{\epsilon \downarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \text{rank}_+(\mathbf{M}_{\epsilon, \delta, n}) = C_W(\mathbf{M}).$$

# Amortization Comes to the Rescue

## Theorem ([Braun et al., 2017])

Let  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$  be such that  $\|\mathbf{M}\|_1 = \sum_{x,y} M_{x,y} = 1$ . For any  $\epsilon, \delta > 0$ , if  $n \geq n_0(\epsilon, \delta, m, k, C_W(\mathbf{M}))$  is sufficiently large, there exists  $\mathbf{M}_{\epsilon, \delta, n} \in \mathbb{R}_+^{m^n \times k^n}$  with

$$\|\mathbf{M}^{\otimes n} - \mathbf{M}_{\epsilon, \delta, n}\|_1 \leq \delta.$$

and

$$\lim_{\epsilon \downarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \text{rank}_+(\mathbf{M}_{\epsilon, \delta, n}) = C_W(\mathbf{M}).$$

- Normalized logarithm of the nonnegative rank of an  $\ell_1$ -perturbed version of  $\mathbf{M}^{\otimes n}$  for large enough  $n$ .

# Amortization Comes to the Rescue

## Theorem ([Braun et al., 2017])

Let  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$  be such that  $\|\mathbf{M}\|_1 = \sum_{x,y} M_{x,y} = 1$ . For any  $\epsilon, \delta > 0$ , if  $n \geq n_0(\epsilon, \delta, m, k, C_W(\mathbf{M}))$  is sufficiently large, there exists  $\mathbf{M}_{\epsilon, \delta, n} \in \mathbb{R}_+^{m^n \times k^n}$  with

$$\|\mathbf{M}^{\otimes n} - \mathbf{M}_{\epsilon, \delta, n}\|_1 \leq \delta.$$

and

$$\lim_{\epsilon \downarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \text{rank}_+(\mathbf{M}_{\epsilon, \delta, n}) = C_W(\mathbf{M}).$$

- Normalized logarithm of the nonnegative rank of an  $\ell_1$ -perturbed version of  $\mathbf{M}^{\otimes n}$  for large enough  $n$ .
- TV common information = Wyner's common information [Cuff, 2012].

# Amortization Comes to the Rescue

## Theorem ([Braun et al., 2017])

Let  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$  be such that  $\|\mathbf{M}\|_1 = \sum_{x,y} M_{x,y} = 1$ . For any  $\epsilon, \delta > 0$ , if  $n \geq n_0(\epsilon, \delta, m, k, C_W(\mathbf{M}))$  is sufficiently large, there exists  $\mathbf{M}_{\epsilon, \delta, n} \in \mathbb{R}_+^{m^n \times k^n}$  with

$$\|\mathbf{M}^{\otimes n} - \mathbf{M}_{\epsilon, \delta, n}\|_1 \leq \delta.$$

and

$$\lim_{\epsilon \downarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \text{rank}_+(\mathbf{M}_{\epsilon, \delta, n}) = C_W(\mathbf{M}).$$

- Normalized logarithm of the nonnegative rank of an  $\ell_1$ -perturbed version of  $\mathbf{M}^{\otimes n}$  for large enough  $n$ .
- TV common information = Wyner's common information [Cuff, 2012].

# Amortization Comes to the Rescue

## Theorem ([Braun et al., 2017])

Let  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$  be such that  $\|\mathbf{M}\|_1 = \sum_{x,y} M_{x,y} = 1$ . For any  $\epsilon, \delta > 0$ , if  $n \geq n_0(\epsilon, \delta, m, k, C_W(\mathbf{M}))$  is sufficiently large, there exists  $\mathbf{M}_{\epsilon, \delta, n} \in \mathbb{R}_+^{m^n \times k^n}$  with

$$\|\mathbf{M}^{\otimes n} - \mathbf{M}_{\epsilon, \delta, n}\|_1 \leq \delta.$$

and

$$\lim_{\epsilon \downarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \text{rank}_+(\mathbf{M}_{\epsilon, \delta, n}) = C_W(\mathbf{M}).$$

- Normalized logarithm of the nonnegative rank of an  $\ell_1$ -perturbed version of  $\mathbf{M}^{\otimes n}$  for large enough  $n$ .
- TV common information = Wyner's common information [Cuff, 2012].

# Amortization Comes to the Rescue

Theorem ([Braun et al., 2017])

Let  $\mathbf{M} \in \mathbb{R}_+^{m \times k}$  be such that  $\|\mathbf{M}\|_1 = \sum_{x,y} M_{x,y} = 1$ . For any  $\epsilon, \delta > 0$ , if  $n \geq n_0(\epsilon, \delta, m, k, C_W(\mathbf{M}))$  is sufficiently large, there exists  $\mathbf{M}_{\epsilon, \delta, n} \in \mathbb{R}_+^{m^n \times k^n}$  with

$$\|\mathbf{M}^{\otimes n} - \mathbf{M}_{\epsilon, \delta, n}\|_1 \leq \delta.$$

and

$$\lim_{\epsilon \downarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \text{rank}_+(\mathbf{M}_{\epsilon, \delta, n}) = C_W(\mathbf{M}).$$

- Normalized logarithm of the nonnegative rank of an  $\ell_1$ -perturbed version of  $\mathbf{M}^{\otimes n}$  for large enough  $n$ .
- TV common information = Wyner's common information [Cuff, 2012].



# Outline

- 1 Introduction: Measures of Information Among Two Random Variables
- 2 Wyner's Common Information
- 3 Rényi Common Information
- 4 Exact Common Information
- 5 Approximate and Exact Channel Synthesis
- 6 Nonnegative Matrix Factorization and Nonnegative Rank
- 7 Extensions of Gács–Körner–Witsenhausen's Common Information (L. Yu)

# References I

-  Beasley, L. B. and Laffey, T. J. (2009).  
Real rank versus nonnegative rank.  
*Linear Algebra and its Applications*, 431(12):2330–2335.
-  Bennett, C. H., Shor, P. W., Smolin, J. A., and Thapliyal, A. V. (2002).  
Entanglement-assisted capacity of a quantum channel and the reverse shannon theorem.  
*IEEE Trans. on Inform. Th.*, 48(10):2637–2655.
-  Braun, G., Jain, R., Lee, T., and Pokutta, S. (2017).  
Information-theoretic approximations of the nonnegative rank.  
*Computational Complexity*, 26:147–197.
-  Braun, G. and Pokutta, S. (2013).  
Common information and unique disjointness.  
In *Proc. of the 54th IEEE Symposium on Foundations of Computer Science*, pages 688–697.
-  Cichocki, A., Zdunek, R., Phan, A. H., and i. Amari, S. (2009).  
*Nonnegative Matrix and Tensor Factorizations: Applications to Exploratory Multi-way Data Analysis and Blind Source Separation*.  
Wiley.
-  Cuff, P. (2012).  
Distributed channel synthesis.  
*IEEE Trans. on Inform. Th.*, 59(11):7071–7096.
-  Gillis, N. (2020).  
*Nonnegative Matrix Factorization*.  
Society for Industrial & Applied Mathematics.

# References II

-  Gray, R. M. and Wyner, A. D. (1974).  
Source coding for a simple network.  
*The Bell Systems Technical Journal*, 53:1681–1721.
-  Han, T. S. and Verdú, S. (1993).  
Approximation theory of output statistics.  
*IEEE Trans. on Inform. Th.*, 39(3):752–772.
-  Hayashi, M. (2006).  
General nonasymptotic and asymptotic formulas in channel resolvability and identification capacity and their application to the wiretap channel.  
*IEEE Trans. on Inform. Th.*, 52(4):1562–1575.
-  Jain, R., Shi, Y., Wei, Z., and Zhang, S. (2013).  
Efficient protocols for generating bipartite classical distributions and quantum states.  
In *Proc. of SODA*.
-  Kumar, G. R., Li, C. T., and El Gamal, A. (2014).  
Exact common information.  
In *Proc. IEEE Int. Symp. Inform. Theory*, pages 161–165, Honolulu, Hawaii.
-  Oohama, Y. (2018).  
Exponential strong converse for source coding with side information at the decoder.  
*Entropy*, 20(5):352.
-  Sason, I. (2016).  
On the Rényi divergence, joint range of relative entropies, and a channel coding theorem.  
*IEEE Trans. on Inform. Th.*, 62(1):23–34.

# References III



Vellambi, B. N. and Kliewer, J. (2016).

Sufficient conditions for the equality of exact and wyner common information.

In *Proceedings of Allerton Conference on Communication, Control, and Computing*, Monticello, IL.



Wyner, A. D. (1975).

The common information of two dependent random variables.

*IEEE Trans. on Inform. Th.*, 21(2):163–179.



Yu, L. and Tan, V. Y. F. (2018).

Wyner's common information under Rényi divergence measures.

*IEEE Trans. on Inform. Th.*, 64(5):3616–3623.



Yu, L. and Tan, V. Y. F. (2019).

Simulation of random variables under Rényi divergence measures of all orders.

*IEEE Trans. on Inform. Th.*, 65(6):3349–3383.



Yu, L. and Tan, V. Y. F. (2020a).

Corrections to “Wyner's common information under Rényi divergence measures”.

*IEEE Trans. on Inform. Th.*, 66(4):2599–2608.



Yu, L. and Tan, V. Y. F. (2020b).

Exact channel synthesis.

*IEEE Trans. on Inform. Th.*, 66(5):2299–2818.



Yu, L. and Tan, V. Y. F. (2020c).

On exact and  $\infty$ -Rényi common information.

*IEEE Trans. on Inform. Th.*, 66(6):3366–3406.

# Common Information: Old and New

## Part II: Gács-Körner-Witsenhausen's CI and Extensions

**Vincent Y. F. Tan and Lei Yu**

National University of Singapore and Nankai University

2021 International Symposium on Information Theory

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Gács–Körner–Witsenhausen's System



# Gács–Körner–Witsenhausen's System



- $(\mathbf{X}, \mathbf{Y}) \sim P_{XY}^n$ : a pair of correlated sources

# Gács–Körner–Witsenhausen's System



- $(\mathbf{X}, \mathbf{Y}) \sim P_{XY}^n$ : a pair of correlated sources
- Define one-sided  $\epsilon$ -common information:

$$T_X(\epsilon) := \liminf_{n \rightarrow \infty} \max_{f, g: \mathbb{P}[f(\mathbf{X}) \neq g(\mathbf{Y})] \leq \epsilon} \frac{1}{n} H(f(\mathbf{X}))$$

$$T_Y(\epsilon) := \liminf_{n \rightarrow \infty} \max_{f, g: \mathbb{P}[f(\mathbf{X}) \neq g(\mathbf{Y})] \leq \epsilon} \frac{1}{n} H(g(\mathbf{Y}))$$

Theorem ([Gács and Körner, 1973])

$$\lim_{\epsilon \downarrow 0} T_X(\epsilon) = \lim_{\epsilon \downarrow 0} T_Y(\epsilon) = C_{\text{GKW}}(X; Y),$$

where

$$C_{\text{GKW}}(X; Y) := \max_{f, g: f(X) = g(Y)} H(f(X))$$

# Gács–Körner–Witsenhausen's CI

Theorem ([Gács and Körner, 1973])

$$\lim_{\epsilon \downarrow 0} T_X(\epsilon) = \lim_{\epsilon \downarrow 0} T_Y(\epsilon) = C_{\text{GKW}}(X; Y),$$

where

$$C_{\text{GKW}}(X; Y) := \max_{f, g: f(X) = g(Y)} H(f(X))$$

- $C_{\text{GKW}}(X; Y)$  called Gács–Körner–Witsenhausen's (GKW's) CI

# Gács–Körner–Witsenhausen's CI

Theorem ([Gács and Körner, 1973])

$$\lim_{\epsilon \downarrow 0} T_X(\epsilon) = \lim_{\epsilon \downarrow 0} T_Y(\epsilon) = C_{\text{GKW}}(X; Y),$$

where

$$C_{\text{GKW}}(X; Y) := \max_{f, g: f(X) = g(Y)} H(f(X))$$

- $C_{\text{GKW}}(X; Y)$  called Gács–Körner–Witsenhausen's (GKW's) CI
- Notice: This is a brief version of GKW's system introduced in [Csiszár and Narayan, 2000]

# Undesired Properties of GKW's CI

- Fact: GKW's CI = 0 for Gaussian sources and doubly symmetric binary sources (DSBSes)
  - More unfortunately, we cannot extract even one pair of identical bits from  $(X, Y)$ , if  $(X, Y)$  is jointly Gaussian or if  $(X, Y)$  is a DSBS.

# Undesired Properties of GKW's CI

- Fact: GKW's CI = 0 for Gaussian sources and doubly symmetric binary sources (DSBSes)
  - More unfortunately, we cannot extract even one pair of identical bits from  $(X, Y)$ , if  $(X, Y)$  is jointly Gaussian or if  $(X, Y)$  is a DSBS.
- How to measure “common information” for this case?

# Undesired Properties of GKW's CI

- Fact: GKW's CI = 0 for Gaussian sources and doubly symmetric binary sources (DSBSes)
  - More unfortunately, we cannot extract even one pair of identical bits from  $(X, Y)$ , if  $(X, Y)$  is jointly Gaussian or if  $(X, Y)$  is a DSBS.
- How to measure "common information" for this case?
- Literally, "common information"  $\iff$  "correlated bits"

# Undesired Properties of GKW's CI

- Fact: GKW's CI = 0 for Gaussian sources and doubly symmetric binary sources (DSBSes)
  - More unfortunately, we cannot extract even one pair of identical bits from  $(X, Y)$ , if  $(X, Y)$  is jointly Gaussian or if  $(X, Y)$  is a DSBS.
- How to measure "common information" for this case?
- Literally, "common information"  $\iff$  "correlated bits"

**A Variant of CI: What is the maximal possible correlation of a pair of bits that can be extracted from  $X, Y$  individually?**

# Undesired Properties of GKW's CI

- Fact: GKW's CI = 0 for Gaussian sources and doubly symmetric binary sources (DSBSes)
  - More unfortunately, we cannot extract even one pair of identical bits from  $(X, Y)$ , if  $(X, Y)$  is jointly Gaussian or if  $(X, Y)$  is a DSBS.
- How to measure "common information" for this case?
- Literally, "common information"  $\iff$  "correlated bits"

**A Variant of CI: What is the maximal possible correlation of a pair of bits that can be extracted from  $X, Y$  individually?**

- Coined the **binary decision** problem [Witsenhausen, 1975],  
the **noninteractive correlation distillation (NICD)** problem [Mossel et al., 2006],  
the **noninteractive binary simulation** problem [Kamath and Anantharam, 2016],  
an **isoperimetric** problem,

...

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Doubly Symmetric Binary Source (DSBS)

Throughout this talk, we mainly consider the DSBS, Unless Otherwise Specified!!!

$$P_{XY} = \begin{array}{c|cc} X \setminus Y & 0 & 1 \\ \hline 0 & \frac{1+\rho}{4} & \frac{1-\rho}{4} \\ 1 & \frac{1-\rho}{4} & \frac{1+\rho}{4} \end{array}$$

with correlation  $\rho \in (0, 1)$ , and

$$(\mathbf{X}, \mathbf{Y}) \sim P_{XY}^n$$

# Doubly Symmetric Binary Source (DSBS)

Throughout this talk, we mainly consider the DSBS, Unless Otherwise Specified!!!


$$P_{XY} = \begin{array}{c|cc} X \setminus Y & 0 & 1 \\ \hline 0 & \frac{1+\rho}{4} & \frac{1-\rho}{4} \\ 1 & \frac{1-\rho}{4} & \frac{1+\rho}{4} \end{array}$$

with correlation  $\rho \in (0, 1)$ , and

$$(\mathbf{X}, \mathbf{Y}) \sim P_{XY}^n$$

- If you are interested in other sources, please refer to [Ahlswede and Gács, 1976, Carlen and Cordero-Erausquin, 2009, Beigi and Nair, 2016, Yu et al., 2021, Yu, 2021c]...

# Noninteractive Correlation Distillation



$$\max \mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y})) \quad \text{or equivalently,} \quad \max \mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1)$$

# Noninteractive Correlation Distillation

- Formally, for  $a, b \in [0, 1]$ , define the **Forward Joint Probability** as

$$\begin{aligned}\bar{\Gamma}^{(n)}(a, b) &:= \max_{\substack{f, g: \{0,1\}^n \rightarrow \{0,1\}: \mathbb{P}(f(\mathbf{X})=1) \leq a, \\ \mathbb{P}(g(\mathbf{Y})=1) \leq b}} \mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \\ &= \max_{\substack{A, B \subseteq \{0,1\}^n: P_X^n(A) \leq a, \\ P_Y^n(B) \leq b}} P_{XY}^n(A \times B), \quad (f = 1_A, g = 1_B)\end{aligned}$$

# Noninteractive Correlation Distillation

- Formally, for  $a, b \in [0, 1]$ , define the **Forward Joint Probability** as

$$\begin{aligned}\bar{\Gamma}^{(n)}(a, b) &:= \max_{\substack{f, g: \{0,1\}^n \rightarrow \{0,1\}: \mathbb{P}(f(\mathbf{X})=1) \leq a, \\ \mathbb{P}(g(\mathbf{Y})=1) \leq b}} \mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \\ &= \max_{\substack{A, B \subseteq \{0,1\}^n: P_X^n(A) \leq a, \\ P_Y^n(B) \leq b}} P_{XY}^n(A \times B), \quad (f = 1_A, g = 1_B)\end{aligned}$$

- Define the **Reverse Joint Probability** as

$$\underline{\Gamma}^{(n)}(a, b) := \min_{\substack{A, B \subseteq \{0,1\}^n: P_X^n(A) \geq a, \\ P_Y^n(B) \geq b}} P_{XY}^n(A \times B)$$

# Noninteractive Correlation Distillation

- Formally, for  $a, b \in [0, 1]$ , define the **Forward Joint Probability** as

$$\begin{aligned}\bar{\Gamma}^{(n)}(a, b) &:= \max_{\substack{f, g: \{0,1\}^n \rightarrow \{0,1\}: \mathbb{P}(f(\mathbf{X})=1) \leq a, \\ \mathbb{P}(g(\mathbf{Y})=1) \leq b}} \mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \\ &= \max_{\substack{A, B \subseteq \{0,1\}^n: P_X^n(A) \leq a, \\ P_Y^n(B) \leq b}} P_{XY}^n(A \times B), \quad (f = 1_A, g = 1_B)\end{aligned}$$

- Define the **Reverse Joint Probability** as

$$\underline{\Gamma}^{(n)}(a, b) := \min_{\substack{A, B \subseteq \{0,1\}^n: P_X^n(A) \geq a, \\ P_Y^n(B) \geq b}} P_{XY}^n(A \times B)$$

- For  $a = \frac{M}{2^n}, b = \frac{N}{2^n}$  (with integers  $M, N$ ), the “inequalities” in the constraints can be replaced by “equalities”

# Noninteractive Correlation Distillation

- Formally, for  $a, b \in [0, 1]$ , define the **Forward Joint Probability** as

$$\begin{aligned}\bar{\Gamma}^{(n)}(a, b) &:= \max_{\substack{f, g: \{0,1\}^n \rightarrow \{0,1\}: \mathbb{P}(f(\mathbf{X})=1) \leq a, \\ \mathbb{P}(g(\mathbf{Y})=1) \leq b}} \mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \\ &= \max_{\substack{A, B \subseteq \{0,1\}^n: P_X^n(A) \leq a, \\ P_Y^n(B) \leq b}} P_{XY}^n(A \times B), \quad (f = 1_A, g = 1_B)\end{aligned}$$

- Define the **Reverse Joint Probability** as

$$\underline{\Gamma}^{(n)}(a, b) := \min_{\substack{A, B \subseteq \{0,1\}^n: P_X^n(A) \geq a, \\ P_Y^n(B) \geq b}} P_{XY}^n(A \times B)$$

- For  $a = \frac{M}{2^n}, b = \frac{N}{2^n}$  (with integers  $M, N$ ), the “inequalities” in the constraints can be replaced by “equalities”
- Equivalence:

$$\bar{\Gamma}^{(\infty)}(1-a, b) = b - \underline{\Gamma}^{(\infty)}(a, b),$$

where  $\bar{\Gamma}^{(\infty)}, \underline{\Gamma}^{(\infty)}$  denote the pointwise limits of  $\bar{\Gamma}^{(n)}, \underline{\Gamma}^{(n)}$  as  $n \rightarrow \infty$ .

# Interesting Cases

Limit cases as  $n \rightarrow \infty$

- Central Limit (CL) regime:  $a = 2^{-\alpha}, b = 2^{-\beta}$  are fixed

# Interesting Cases

Limit cases as  $n \rightarrow \infty$

- Central Limit (CL) regime:  $a = 2^{-\alpha}, b = 2^{-\beta}$  are fixed
- Large Deviation (LD) regime:  $a = 2^{-n\alpha}, b = 2^{-n\beta}$  are exponentially small

# Interesting Cases

Limit cases as  $n \rightarrow \infty$

- Central Limit (CL) regime:  $a = 2^{-\alpha}, b = 2^{-\beta}$  are fixed
- Large Deviation (LD) regime:  $a = 2^{-n\alpha}, b = 2^{-n\beta}$  are exponentially small
- Moderate Deviation (MD) regime:  $a = 2^{-\theta_n \alpha}, b = 2^{-\theta_n \beta}$  are “subexponentially” small with  
 $\theta_n \rightarrow \infty, \frac{\theta_n}{n} \rightarrow 0$ 
  - interpolates between the CL and LD regimes

# Interesting Cases

Limit cases as  $n \rightarrow \infty$

- Central Limit (CL) regime:  $a = 2^{-\alpha}, b = 2^{-\beta}$  are fixed
- Large Deviation (LD) regime:  $a = 2^{-n\alpha}, b = 2^{-n\beta}$  are exponentially small
- Moderate Deviation (MD) regime:  $a = 2^{-\theta_n \alpha}, b = 2^{-\theta_n \beta}$  are “subexponentially” small with  
 $\theta_n \rightarrow \infty, \frac{\theta_n}{n} \rightarrow 0$ 
  - interpolates between the CL and LD regimes
- Notice: In all the above cases,  $\alpha, \beta > 0$  are fixed.

# Definitions of Several Exponents

- (Forward and Reverse) CL Exponents: For  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \bar{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right), \quad \overline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \underline{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right)$$

# Definitions of Several Exponents

- (Forward and Reverse) **CL Exponents**: For  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \bar{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right), \quad \overline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \underline{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right)$$

- (Forward and Reverse) **LD Exponents**: For  $\alpha, \beta \in (0, 1)$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \bar{\Gamma}^{(n)}\left(e^{-n\alpha}, e^{-n\beta}\right), \quad \overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \underline{\Gamma}^{(n)}\left(e^{-n\alpha}, e^{-n\beta}\right)$$

# Definitions of Several Exponents

- (Forward and Reverse) **CL Exponents**: For  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \bar{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right), \quad \bar{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \underline{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right)$$

- (Forward and Reverse) **LD Exponents**: For  $\alpha, \beta \in (0, 1)$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \bar{\Gamma}^{(n)}\left(e^{-\textcolor{red}{n}\alpha}, e^{-\textcolor{red}{n}\beta}\right), \quad \bar{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \underline{\Gamma}^{(n)}\left(e^{-n\alpha}, e^{-n\beta}\right)$$

- (Forward and Reverse) **MD Exponents**: For a seq.  $\{\theta_n\}$  s.t.  $\theta_n \rightarrow \infty$  and  $\frac{\theta_n}{n} \rightarrow 0$ , for  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) := -\frac{1}{\theta_n} \log \bar{\Gamma}^{(n)}\left(e^{-\theta_n\alpha}, e^{-\theta_n\beta}\right), \quad \bar{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) := -\frac{1}{\theta_n} \log \underline{\Gamma}^{(n)}\left(e^{-\theta_n\alpha}, e^{-\theta_n\beta}\right)$$

# Definitions of Several Exponents

- (Forward and Reverse) **CL Exponents**: For  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \bar{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right), \quad \bar{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \underline{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right)$$

- (Forward and Reverse) **LD Exponents**: For  $\alpha, \beta \in (0, 1)$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \bar{\Gamma}^{(n)}\left(e^{-n\alpha}, e^{-n\beta}\right), \quad \bar{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \underline{\Gamma}^{(n)}\left(e^{-n\alpha}, e^{-n\beta}\right)$$

- (Forward and Reverse) **MD Exponents**: For a seq.  $\{\theta_n\}$  s.t.  $\theta_n \rightarrow \infty$  and  $\frac{\theta_n}{n} \rightarrow 0$ , for  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) := -\frac{1}{\theta_n} \log \bar{\Gamma}^{(n)}\left(e^{-\theta_n \alpha}, e^{-\theta_n \beta}\right), \quad \bar{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) := -\frac{1}{\theta_n} \log \underline{\Gamma}^{(n)}\left(e^{-\theta_n \alpha}, e^{-\theta_n \beta}\right)$$

- Denote  $\underline{\Theta}_{\text{CL}}^{(\infty)}, \bar{\Theta}_{\text{CL}}^{(\infty)}, \underline{\Theta}_{\text{LD}}^{(\infty)}, \bar{\Theta}_{\text{LD}}^{(\infty)}, \underline{\Theta}_{\text{MD}}^{(\infty)}, \bar{\Theta}_{\text{MD}}^{(\infty)}$  as the pointwise limits of the above as  $n \rightarrow \infty$ .

# Definitions of Several Exponents

- (Forward and Reverse) **CL Exponents**: For  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \bar{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right), \quad \bar{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta) := -\log \underline{\Gamma}^{(n)}\left(e^{-\alpha}, e^{-\beta}\right)$$

- (Forward and Reverse) **LD Exponents**: For  $\alpha, \beta \in (0, 1)$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \bar{\Gamma}^{(n)}\left(e^{-n\alpha}, e^{-n\beta}\right), \quad \bar{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) := -\frac{1}{n} \log \underline{\Gamma}^{(n)}\left(e^{-n\alpha}, e^{-n\beta}\right)$$

- (Forward and Reverse) **MD Exponents**: For a seq.  $\{\theta_n\}$  s.t.  $\theta_n \rightarrow \infty$  and  $\frac{\theta_n}{n} \rightarrow 0$ , for  $\alpha, \beta \in (0, \infty)$ ,

$$\underline{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) := -\frac{1}{\theta_n} \log \bar{\Gamma}^{(n)}\left(e^{-\theta_n\alpha}, e^{-\theta_n\beta}\right), \quad \bar{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) := -\frac{1}{\theta_n} \log \underline{\Gamma}^{(n)}\left(e^{-\theta_n\alpha}, e^{-\theta_n\beta}\right)$$

- Denote  $\underline{\Theta}_{\text{CL}}^{(\infty)}, \bar{\Theta}_{\text{CL}}^{(\infty)}, \underline{\Theta}_{\text{LD}}^{(\infty)}, \bar{\Theta}_{\text{LD}}^{(\infty)}, \underline{\Theta}_{\text{MD}}^{(\infty)}, \bar{\Theta}_{\text{MD}}^{(\infty)}$  as the pointwise limits of the above as  $n \rightarrow \infty$ .
- Equivalence: For finite  $n$ ,  $\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) = \frac{1}{n} \underline{\Theta}_{\text{CL}}^{(n)}(n\alpha, n\beta)$  and  $\bar{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) = \frac{1}{\theta_n} \bar{\Theta}_{\text{CL}}^{(n)}(\theta_n\alpha, \theta_n\beta)$ ; Similar formulas hold for the reverse part.

# Outline

## 1 Gács–Körner–Witsenhausen's CI


## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability


- Formulation
- Balanced Case
- MD and LD Regimes

# Achievability: Hamming Subcubes




- An  $(n - k)$ -subcube  $C_{n-k}$  is a set of  $\mathbf{x}$  with  $k$  components fixed (e.g.,  $\{1_k\} \times \{0, 1\}^{n-k}$  whose indicator  $\mathbf{x} \mapsto \prod_{i=1}^k x_i$ )

# Achievability: Hamming Subcubes




- An  $(n - k)$ -subcube  $C_{n-k}$  is a set of  $\mathbf{x}$  with  $k$  components fixed (e.g.,  $\{1_k\} \times \{0, 1\}^{n-k}$  whose indicator  $\mathbf{x} \mapsto \prod_{i=1}^k x_i$ )
  - Special case  $C_{n-1}$ : e.g.,  $\{1\} \times \{0, 1\}^{n-1}$  (Indicator  $\mathbf{x} \mapsto x_1$  called a **dictator** function)

# Achievability: Hamming Subcubes




- An  $(n - k)$ -subcube  $C_{n-k}$  is a set of  $\mathbf{x}$  with  $k$  components fixed (e.g.,  $\{1_k\} \times \{0, 1\}^{n-k}$  whose indicator  $\mathbf{x} \mapsto \prod_{i=1}^k x_i$ )
  - Special case  $C_{n-1}$ : e.g.,  $\{1\} \times \{0, 1\}^{n-1}$  (Indicator  $\mathbf{x} \mapsto x_1$  called a **dictator** function)
- Case of  $a = b = 2^{-k}$ :  $A = B = C_{n-k}$  (**identical**)  $\implies P_{XY}^n(A \times B) = P_{XY}(1, 1)^k = \left(\frac{1+\rho}{4}\right)^k$ ;  
 $A = 1 - B = C_{n-k}$  (**anti-symmetric**)  $\implies P_{XY}^n(A \times B) = P_{XY}(1, 0)^k = \left(\frac{1-\rho}{4}\right)^k$

# Achievability: Hamming Subcubes




- An  $(n - k)$ -subcube  $C_{n-k}$  is a set of  $\mathbf{x}$  with  $k$  components fixed (e.g.,  $\{1_k\} \times \{0, 1\}^{n-k}$  whose indicator  $\mathbf{x} \mapsto \prod_{i=1}^k x_i$ )
  - Special case  $C_{n-1}$ : e.g.,  $\{1\} \times \{0, 1\}^{n-1}$  (Indicator  $\mathbf{x} \mapsto x_1$  called a **dictator** function)
- Case of  $a = b = 2^{-k}$ :  $A = B = C_{n-k}$  (**identical**)  $\implies P_{XY}^n(A \times B) = P_{XY}(1, 1)^k = \left(\frac{1+\rho}{4}\right)^k$ ;  
 $A = 1 - B = C_{n-k}$  (**anti-symmetric**)  $\implies P_{XY}^n(A \times B) = P_{XY}(1, 0)^k = \left(\frac{1-\rho}{4}\right)^k$
- Case of  $a = 2^{-k_1}, b = 2^{-k_2}, k_1 \leq k_2$ :  $P_{XY}^n(A \times B) = \left(\frac{1}{2}\right)^{k_2 - k_1} \left(\frac{1+\rho}{4}\right)^{k_1}$  (resp.  $\left(\frac{1}{2}\right)^{k_2 - k_1} \left(\frac{1-\rho}{4}\right)^{k_1}$ ) for “almost identical” (resp. “almost anti-symmetric”) subcubes  $A, B$

# Achievability: Hamming Balls (CL Regime)




- Hamming Ball: For  $r \in [0, n]$ ,  $\mathbb{B}_r(\mathbf{0}) := \{\mathbf{x} : d_H(\mathbf{x}, \mathbf{0}) \leq r\} \iff \{\mathbf{x} : \sum_{i=1}^n x_i \leq r\}$

# Achievability: Hamming Balls (CL Regime)



- Hamming Ball: For  $r \in [0, n]$ ,  $\mathbb{B}_r(\mathbf{0}) := \{\mathbf{x} : d_H(\mathbf{x}, \mathbf{0}) \leq r\} \iff \{\mathbf{x} : \sum_{i=1}^n x_i \leq r\}$
- CL regime: Choose  $A = \mathbb{B}_{r_n}(\mathbf{0})$ ,  $B = \mathbb{B}_{s_n}(\mathbf{0})$  with  $r_n = \frac{n}{2} + \frac{\lambda\sqrt{n}}{2}$ ,  $s_n = \frac{n}{2} + \frac{\mu\sqrt{n}}{2}$  where  $\lambda, \mu \in \mathbb{R}$

# Achievability: Hamming Balls (CL Regime)



- Hamming Ball: For  $r \in [0, n]$ ,  $\mathbb{B}_r(\mathbf{0}) := \{\mathbf{x} : d_H(\mathbf{x}, \mathbf{0}) \leq r\} \iff \{\mathbf{x} : \sum_{i=1}^n x_i \leq r\}$
- CL regime: Choose  $A = \mathbb{B}_{r_n}(\mathbf{0})$ ,  $B = \mathbb{B}_{s_n}(\mathbf{0})$  with  $r_n = \frac{n}{2} + \frac{\lambda\sqrt{n}}{2}$ ,  $s_n = \frac{n}{2} + \frac{\mu\sqrt{n}}{2}$  where  $\lambda, \mu \in \mathbb{R}$
- By the univariate and multivariate CL theorems,

$$P_X^n(A) \rightarrow \Phi(\lambda), \quad P_Y^n(B) \rightarrow \Phi(\mu), \quad P_{XY}^n(A \times B) \rightarrow \Phi_\rho(\lambda, \mu)$$

where  $\Phi$  is the CDF of the standard Gaussian, and  $\Phi_\rho(\cdot, \cdot)$  is the CDF of the zero-mean bivariate Gaussian with covariance matrix  $\begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}$ .

# Achievability: Hamming Balls (CL Regime)

- Achievable CL probabilities:

$$\overline{\Gamma}^{(\infty)}(a, b) \geq \Lambda_{\rho}(a, b) \text{ (by concentric balls)}$$

- Bivariate normal copula (or Gaussian quadrant probability function):

$$\Lambda_{\rho}(a, b) := \Phi_{\rho}\left(\Phi^{-1}(a), \Phi^{-1}(b)\right)$$

# Achievability: Hamming Balls (CL Regime)

- Achievable CL probabilities:

$$\overline{\Gamma}^{(\infty)}(a, b) \geq \Lambda_{\rho}(a, b) \text{ (by concentric balls)}$$

- Bivariate normal copula (or Gaussian quadrant probability function):

$$\Lambda_{\rho}(a, b) := \Phi_{\rho}\left(\Phi^{-1}(a), \Phi^{-1}(b)\right)$$

- $\iff$

$$\underline{\Gamma}^{(\infty)}(a, b) \leq \Lambda_{-\rho}(a, b) \text{ (by anti-concentric balls)}$$

- $\Lambda_{-\rho}(a, b)$  is attained by anti-concentric balls  $A = \mathbb{B}_{r_n}(\mathbf{0}), B = \mathbb{B}_{s_n}(1)$

# Achievability: Hamming Balls (CL Regime)

- Achievable CL probabilities:

$$\overline{\Gamma}^{(\infty)}(a, b) \geq \Lambda_{\rho}(a, b) \text{ (by concentric balls)}$$

- Bivariate normal copula (or Gaussian quadrant probability function):

$$\Lambda_{\rho}(a, b) := \Phi_{\rho}\left(\Phi^{-1}(a), \Phi^{-1}(b)\right)$$

•  $\iff$

$$\underline{\Gamma}^{(\infty)}(a, b) \leq \Lambda_{-\rho}(a, b) \text{ (by anti-concentric balls)}$$

- $\Lambda_{-\rho}(a, b)$  is attained by anti-concentric balls  $A = \mathbb{B}_{r_n}(\mathbf{0}), B = \mathbb{B}_{s_n}(1)$


•  $\iff$

$$\underline{\Theta}_{\text{CL}}^{(\infty)}(\alpha, \beta) \leq \underline{\Theta}_{\text{CL}}(\alpha, \beta), \quad \overline{\Theta}_{\text{CL}}^{(\infty)}(\alpha, \beta) \geq \overline{\Theta}_{\text{CL}}(\alpha, \beta)$$

- Exponents of  $\Lambda_{\rho}$  and  $\Lambda_{-\rho}$ :


$$\underline{\Theta}_{\text{CL}}(\alpha, \beta) := -\log \Lambda_{\rho}\left(e^{-\alpha}, e^{-\beta}\right), \quad \overline{\Theta}_{\text{CL}}(\alpha, \beta) := -\log \Lambda_{-\rho}\left(e^{-\alpha}, e^{-\beta}\right)$$

# Achievability: Hamming Spheres (LD Regime)




- Hamming Sphere: For  $r \in [0 : n]$ ,  $\mathbb{S}_r(\mathbf{0}) := \{\mathbf{x} : d_H(\mathbf{x}, \mathbf{0}) = r\} \iff \{\mathbf{x} : \sum_{i=1}^n x_i = r\}$

# Achievability: Hamming Spheres (LD Regime)



- Hamming Sphere: For  $r \in [0 : n]$ ,  $\mathbb{S}_r(\mathbf{0}) := \{\mathbf{x} : d_H(\mathbf{x}, \mathbf{0}) = r\} \iff \{\mathbf{x} : \sum_{i=1}^n x_i = r\}$ 
  - It can be regarded as a type class with type  $(\lambda, \bar{\lambda})$  in Hamming space, where  $\lambda := \frac{r}{n}$  and  $\bar{\lambda} := 1 - \lambda$

# Achievability: Hamming Spheres (LD Regime)



- Hamming Sphere: For  $r \in [0 : n]$ ,  $\mathbb{S}_r(\mathbf{0}) := \{\mathbf{x} : d_H(\mathbf{x}, \mathbf{0}) = r\} \iff \{\mathbf{x} : \sum_{i=1}^n x_i = r\}$ 
  - It can be regarded as a type class with type  $(\lambda, \bar{\lambda})$  in Hamming space, where  $\lambda := \frac{r}{n}$  and  $\bar{\lambda} := 1 - \lambda$
- LD regime: Choose  $A = \mathbb{S}_{r_n}(\mathbf{0}), B = \mathbb{S}_{s_n}(\mathbf{0})$  with  $r_n = \lambda n, s_n = \mu n$  where  $\lambda, \mu \in [0, 1]$

# Achievability: Hamming Spheres (LD Regime)

- By LD theory (or Sanov's theorem),

$$-\frac{1}{n} \log P_X^n(A) \rightarrow D((\lambda, \bar{\lambda}) \| P_X) = 1 - H_2(\lambda)$$

$$-\frac{1}{n} \log P_Y^n(B) \rightarrow D((\mu, \bar{\mu}) \| P_Y) = 1 - H_2(\mu)$$

$$-\frac{1}{n} \log P_{XY}^n(A \times B) \rightarrow \mathbb{D}((\lambda, \bar{\lambda}), (\mu, \bar{\mu}) \| P_{XY})$$

# Achievability: Hamming Spheres (LD Regime)

- By LD theory (or Sanov's theorem),

$$-\frac{1}{n} \log P_X^n(A) \rightarrow D((\lambda, \bar{\lambda}) \| P_X) = 1 - H_2(\lambda)$$

$$-\frac{1}{n} \log P_Y^n(B) \rightarrow D((\mu, \bar{\mu}) \| P_Y) = 1 - H_2(\mu)$$

$$-\frac{1}{n} \log P_{XY}^n(A \times B) \rightarrow \mathbb{D}((\lambda, \bar{\lambda}), (\mu, \bar{\mu}) \| P_{XY})$$

- relative entropy:  $D(Q \| P) := \sum_x Q(x) \log \frac{Q(x)}{P(x)}$
- binary entropy function:  $H_2 : t \in [0, 1] \mapsto -t \log_2 t - (1-t) \log_2(1-t)$
- **minimum-relative-entropy** over couplings of  $(Q_X, Q_Y)$ :

$$\mathbb{D}(Q_X, Q_Y \| P_{XY}) := \min_{Q_{XY} \in C(Q_X, Q_Y)} D(Q_{XY} \| P_{XY})$$

with  $C(Q_X, Q_Y) := \{Q_{XY} \text{ with marginals } Q_X, Q_Y\}$  denoting the coupling set

# Achievability: Hamming Spheres (LD Regime)

- Optimizing  $\mathbb{D}(Q_X, Q_Y \| P_{XY})$  over feasible  $Q_X := (\lambda, \bar{\lambda}), Q_Y := (\mu, \bar{\mu}) \implies$

$$\underline{\Theta}_{\text{LD}}^{(\infty)}(\alpha, \beta) \leq \underline{\Theta}_{\text{LD}}(\alpha, \beta) := \min_{\substack{Q_X, Q_Y: D(Q_X \| P_X) \geq \alpha, \\ D(Q_Y \| P_Y) \geq \beta}} \mathbb{D}(Q_X, Q_Y \| P_{XY}),$$

$$\overline{\Theta}_{\text{LD}}^{(\infty)}(\alpha, \beta) \geq \overline{\Theta}_{\text{LD}}(\alpha, \beta) := \max_{\substack{Q_X, Q_Y: D(Q_X \| P_X) \leq \alpha, \\ D(Q_Y \| P_Y) \leq \beta}} \mathbb{D}(Q_X, Q_Y \| P_{XY}).$$

- attained by **concentric** and **anti-concentric** spheres
- or respectively attained by concentric and anti-concentric **balls**.

# Achievability: Hamming Spheres (LD Regime)

- Optimizing  $\mathbb{D}(Q_X, Q_Y \| P_{XY})$  over feasible  $Q_X := (\lambda, \bar{\lambda}), Q_Y := (\mu, \bar{\mu}) \implies$

$$\underline{\Theta}_{\text{LD}}^{(\infty)}(\alpha, \beta) \leq \underline{\Theta}_{\text{LD}}(\alpha, \beta) := \min_{\substack{Q_X, Q_Y: D(Q_X \| P_X) \geq \alpha, \\ D(Q_Y \| P_Y) \geq \beta}} \mathbb{D}(Q_X, Q_Y \| P_{XY}),$$

$$\overline{\Theta}_{\text{LD}}^{(\infty)}(\alpha, \beta) \geq \overline{\Theta}_{\text{LD}}(\alpha, \beta) := \max_{\substack{Q_X, Q_Y: D(Q_X \| P_X) \leq \alpha, \\ D(Q_Y \| P_Y) \leq \beta}} \mathbb{D}(Q_X, Q_Y \| P_{XY}).$$

- attained by **concentric** and **anti-concentric** spheres
- or respectively attained by concentric and anti-concentric **balls**.

## Conjecture (Orlentlich–Polyanskiy–Shayevitz's (OPS's) Conjecture (2019))

For the DSBS and  $\alpha, \beta \in (0, 1)$ ,

$$\underline{\Theta}_{\text{LD}}^{(\infty)}(\alpha, \beta) = \underline{\Theta}_{\text{LD}}(\alpha, \beta), \quad \overline{\Theta}_{\text{LD}}^{(\infty)}(\alpha, \beta) = \overline{\Theta}_{\text{LD}}(\alpha, \beta).$$

# Achievability: Hamming Spheres (MD Regime)

- MD regime: Choose  $A = \mathbb{S}_{r_n}(\mathbf{0})$ ,  $B = \mathbb{S}_{s_n}(\mathbf{0})$  with  $r_n = \frac{n}{2} + \lambda\sqrt{n\theta_n}$ ,  $s_n = \frac{n}{2} + \mu\sqrt{n\theta_n}$  where  $\theta_n \rightarrow \infty$ ,  $\frac{\theta_n}{n} \rightarrow 0$ , and  $\lambda, \mu \in \mathbb{R}$

# Achievability: Hamming Spheres (MD Regime)

- MD regime: Choose  $A = \mathbb{S}_{r_n}(\mathbf{0})$ ,  $B = \mathbb{S}_{s_n}(\mathbf{0})$  with  $r_n = \frac{n}{2} + \lambda\sqrt{n\theta_n}$ ,  $s_n = \frac{n}{2} + \mu\sqrt{n\theta_n}$  where  $\theta_n \rightarrow \infty$ ,  $\frac{\theta_n}{n} \rightarrow 0$ , and  $\lambda, \mu \in \mathbb{R}$
- By MD theory,

$$\begin{aligned}-\frac{1}{\theta_n} \log P_X^n(A) &\rightarrow \frac{1}{2} \chi^2((\lambda, \bar{\lambda}) \| P_X), & -\frac{1}{\theta_n} \log P_Y^n(B) &\rightarrow \frac{1}{2} \chi^2((\mu, \bar{\mu}) \| P_Y), \\-\frac{1}{\theta_n} \log P_{XY}^n(A \times B) &\rightarrow \frac{1}{2} \mathbb{X}^2((\lambda, \bar{\lambda}), (\mu, \bar{\mu}) \| P_{XY})\end{aligned}$$

# Achievability: Hamming Spheres (MD Regime)

- MD regime: Choose  $A = \mathbb{S}_{r_n}(\mathbf{0})$ ,  $B = \mathbb{S}_{s_n}(\mathbf{0})$  with  $r_n = \frac{n}{2} + \lambda\sqrt{n\theta_n}$ ,  $s_n = \frac{n}{2} + \mu\sqrt{n\theta_n}$  where  $\theta_n \rightarrow \infty$ ,  $\frac{\theta_n}{n} \rightarrow 0$ , and  $\lambda, \mu \in \mathbb{R}$
- By MD theory,

$$\begin{aligned}-\frac{1}{\theta_n} \log P_X^n(A) &\rightarrow \frac{1}{2}\chi^2((\lambda, \bar{\lambda}) \| P_X), & -\frac{1}{\theta_n} \log P_Y^n(B) &\rightarrow \frac{1}{2}\chi^2((\mu, \bar{\mu}) \| P_Y), \\-\frac{1}{\theta_n} \log P_{XY}^n(A \times B) &\rightarrow \frac{1}{2}\mathbb{X}^2((\lambda, \bar{\lambda}), (\mu, \bar{\mu}) \| P_{XY})\end{aligned}$$

- $\chi^2$ -divergence:  $\chi^2(Q \| P) := \sum_x P(x) \left( \frac{Q(x)}{P(x)} - 1 \right)^2$
- minimum- $\chi^2$ -divergence over couplings of  $(Q_X, Q_Y)$ :

$$\mathbb{X}^2(Q_X, Q_Y \| P_{XY}) := \min_{Q_{XY} \in \mathcal{C}(Q_X, Q_Y)} \chi^2(Q_{XY} \| P_{XY})$$

# Achievability: Hamming Spheres (MD Regime)

- MD regime: Choose  $A = \mathbb{S}_{r_n}(\mathbf{0})$ ,  $B = \mathbb{S}_{s_n}(\mathbf{0})$  with  $r_n = \frac{n}{2} + \lambda\sqrt{n\theta_n}$ ,  $s_n = \frac{n}{2} + \mu\sqrt{n\theta_n}$  where  $\theta_n \rightarrow \infty$ ,  $\frac{\theta_n}{n} \rightarrow 0$ , and  $\lambda, \mu \in \mathbb{R}$
- By MD theory,

$$\begin{aligned}-\frac{1}{\theta_n} \log P_X^n(A) &\rightarrow \frac{1}{2}\chi^2((\lambda, \bar{\lambda}) \| P_X), & -\frac{1}{\theta_n} \log P_Y^n(B) &\rightarrow \frac{1}{2}\chi^2((\mu, \bar{\mu}) \| P_Y), \\-\frac{1}{\theta_n} \log P_{XY}^n(A \times B) &\rightarrow \frac{1}{2}\mathbb{X}^2((\lambda, \bar{\lambda}), (\mu, \bar{\mu}) \| P_{XY})\end{aligned}$$

- $\chi^2$ -divergence:  $\chi^2(Q \| P) := \sum_x P(x) \left( \frac{Q(x)}{P(x)} - 1 \right)^2$
- minimum- $\chi^2$ -divergence over couplings of  $(Q_X, Q_Y)$ :

$$\mathbb{X}^2(Q_X, Q_Y \| P_{XY}) := \min_{Q_{XY} \in \mathcal{C}(Q_X, Q_Y)} \chi^2(Q_{XY} \| P_{XY})$$

- Same as the LD case but with  $D(\cdot \| \cdot)$  replaced by  $\frac{1}{2}\chi^2(\cdot \| \cdot)$

# Achievability: Hamming Spheres (MD Regime)

- Optimizing  $\mathbb{X}^2(Q_X, Q_Y \| P_{XY})$  over feasible  $Q_X := (r, \bar{r}), Q_Y := (s, \bar{s}) \implies$

$$\underline{\Theta}_{\text{MD}}^{(\infty)}(\alpha, \beta) \leq \underline{\Theta}_{\text{MD}}(\alpha, \beta) := \min_{\substack{Q_X, Q_Y: \chi^2(Q_X \| P_X) \geq \alpha, \\ \chi^2(Q_Y \| P_Y) \geq \beta}} \mathbb{X}^2(Q_X, Q_Y \| P_{XY}) = \begin{cases} \frac{\alpha + \beta - 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}, & \rho^2\alpha \leq \beta \leq \frac{\alpha}{\rho^2} \\ \alpha, & \beta < \rho^2\alpha \\ \beta, & \alpha < \rho^2\beta \end{cases}$$

$$\overline{\Theta}_{\text{MD}}^{(\infty)}(\alpha, \beta) \geq \overline{\Theta}_{\text{MD}}(\alpha, \beta) := \max_{\substack{Q_X, Q_Y: \chi^2(Q_X \| P_X) \leq \alpha, \\ \chi^2(Q_Y \| P_Y) \leq \beta}} \mathbb{X}^2(Q_X, Q_Y \| P_{XY}) = \frac{\alpha + \beta + 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}$$

# Achievability: Hamming Spheres (MD Regime)

- Optimizing  $\mathbb{X}^2(Q_X, Q_Y \| P_{XY})$  over feasible  $Q_X := (r, \bar{r}), Q_Y := (s, \bar{s}) \implies$

$$\underline{\Theta}_{\text{MD}}^{(\infty)}(\alpha, \beta) \leq \underline{\Theta}_{\text{MD}}(\alpha, \beta) := \min_{\substack{Q_X, Q_Y: \chi^2(Q_X \| P_X) \geq \alpha, \\ \chi^2(Q_Y \| P_Y) \geq \beta}} \mathbb{X}^2(Q_X, Q_Y \| P_{XY}) = \begin{cases} \frac{\alpha + \beta - 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}, & \rho^2\alpha \leq \beta \leq \frac{\alpha}{\rho^2} \\ \alpha, & \beta < \rho^2\alpha \\ \beta, & \alpha < \rho^2\beta \end{cases}$$

$$\overline{\Theta}_{\text{MD}}^{(\infty)}(\alpha, \beta) \geq \overline{\Theta}_{\text{MD}}(\alpha, \beta) := \max_{\substack{Q_X, Q_Y: \chi^2(Q_X \| P_X) \leq \alpha, \\ \chi^2(Q_Y \| P_Y) \leq \beta}} \mathbb{X}^2(Q_X, Q_Y \| P_{XY}) = \frac{\alpha + \beta + 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}$$

- attained by **concentric** and **anti-concentric** spheres
- or respectively attained by concentric and anti-concentric **balls**.

# Achievability: Hamming Spheres (MD Regime)

- Optimizing  $\mathbb{X}^2(Q_X, Q_Y \| P_{XY})$  over feasible  $Q_X := (r, \bar{r}), Q_Y := (s, \bar{s}) \implies$

$$\underline{\Theta}_{\text{MD}}^{(\infty)}(\alpha, \beta) \leq \underline{\Theta}_{\text{MD}}(\alpha, \beta) := \min_{\substack{Q_X, Q_Y: \chi^2(Q_X \| P_X) \geq \alpha, \\ \chi^2(Q_Y \| P_Y) \geq \beta}} \mathbb{X}^2(Q_X, Q_Y \| P_{XY}) = \begin{cases} \frac{\alpha + \beta - 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}, & \rho^2\alpha \leq \beta \leq \frac{\alpha}{\rho^2} \\ \alpha, & \beta < \rho^2\alpha \\ \beta, & \alpha < \rho^2\beta \end{cases}$$

$$\overline{\Theta}_{\text{MD}}^{(\infty)}(\alpha, \beta) \geq \overline{\Theta}_{\text{MD}}(\alpha, \beta) := \max_{\substack{Q_X, Q_Y: \chi^2(Q_X \| P_X) \leq \alpha, \\ \chi^2(Q_Y \| P_Y) \leq \beta}} \mathbb{X}^2(Q_X, Q_Y \| P_{XY}) = \frac{\alpha + \beta + 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}$$

- attained by **concentric** and **anti-concentric** spheres
  - or respectively attained by concentric and anti-concentric **balls**.
- Notice: All  $\frac{1}{2}$ 's have been removed since the resulting exponents  $\frac{1}{k}\underline{\Theta}_{\text{MD}}(k\alpha, k\beta), \frac{1}{k}\overline{\Theta}_{\text{MD}}(k\alpha, k\beta)$  with  $k = 2$  are respectively equal to  $\underline{\Theta}_{\text{MD}}(\alpha, \beta), \overline{\Theta}_{\text{MD}}(\alpha, \beta)$

# Both LD and CL imply MD

- LD implies MD:

- For MD,  $Q_X \rightarrow P_X, Q_Y \rightarrow P_Y$ .
- Fact:  $D(Q\|P) = \frac{1}{2}\chi^2(Q\|P) + o(\chi^2(Q\|P))$  (by Taylor's expansion) and  
 $\mathbb{D}(Q_X, Q_Y \| P_{XY}) = \frac{1}{2}\mathbb{X}^2(Q_X, Q_Y \| P_{XY}) + o(\mathbb{X}^2(Q_X, Q_Y \| P_{XY}))$

# Both LD and CL imply MD

- LD implies MD:

- For MD,  $Q_X \rightarrow P_X, Q_Y \rightarrow P_Y$ .
- Fact:  $D(Q\|P) = \frac{1}{2}\chi^2(Q\|P) + o(\chi^2(Q\|P))$  (by Taylor's expansion) and  
 $\mathbb{D}(Q_X, Q_Y \| P_{XY}) = \frac{1}{2}\mathbb{X}^2(Q_X, Q_Y \| P_{XY}) + o(\mathbb{X}^2(Q_X, Q_Y \| P_{XY}))$
- Replacing  $D \leftarrow \frac{1}{2}\chi^2, \mathbb{D} \leftarrow \frac{1}{2}\mathbb{X}^2$  in LD exponents  $\Rightarrow$  MD exponents

# Both LD and CL imply MD

- LD implies MD:

- For MD,  $Q_X \rightarrow P_X, Q_Y \rightarrow P_Y$ .
- Fact:  $D(Q\|P) = \frac{1}{2}\chi^2(Q\|P) + o(\chi^2(Q\|P))$  (by Taylor's expansion) and  
 $\mathbb{D}(Q_X, Q_Y \| P_{XY}) = \frac{1}{2}\mathbb{X}^2(Q_X, Q_Y \| P_{XY}) + o(\mathbb{X}^2(Q_X, Q_Y \| P_{XY}))$
- Replacing  $D \leftarrow \frac{1}{2}\chi^2, \mathbb{D} \leftarrow \frac{1}{2}\mathbb{X}^2$  in LD exponents  $\Rightarrow$  MD exponents

- Formally,

$$\lim_{\epsilon \downarrow 0} \frac{1}{\epsilon} \underline{\Theta}_{\text{LD}}(\epsilon\alpha, \epsilon\beta) = \underline{\Theta}_{\text{MD}}(\alpha, \beta), \quad \lim_{\epsilon \downarrow 0} \frac{1}{\epsilon} \overline{\Theta}_{\text{LD}}(\epsilon\alpha, \epsilon\beta) = \overline{\Theta}_{\text{MD}}(\alpha, \beta)$$

# Both LD and CL imply MD

- LD implies MD:

- For MD,  $Q_X \rightarrow P_X, Q_Y \rightarrow P_Y$ .
- Fact:  $D(Q||P) = \frac{1}{2}\chi^2(Q||P) + o(\chi^2(Q||P))$  (by Taylor's expansion) and  
 $\mathbb{D}(Q_X, Q_Y || P_{XY}) = \frac{1}{2}\mathbb{X}^2(Q_X, Q_Y || P_{XY}) + o(\mathbb{X}^2(Q_X, Q_Y || P_{XY}))$
- Replacing  $D \leftarrow \frac{1}{2}\chi^2, \mathbb{D} \leftarrow \frac{1}{2}\mathbb{X}^2$  in LD exponents  $\Rightarrow$  MD exponents

- Formally,


$$\lim_{\epsilon \downarrow 0} \frac{1}{\epsilon} \underline{\Theta}_{\text{LD}}(\epsilon\alpha, \epsilon\beta) = \underline{\Theta}_{\text{MD}}(\alpha, \beta), \quad \lim_{\epsilon \downarrow 0} \frac{1}{\epsilon} \overline{\Theta}_{\text{LD}}(\epsilon\alpha, \epsilon\beta) = \overline{\Theta}_{\text{MD}}(\alpha, \beta)$$

- CL implies MD [O'Donnell, 2014]:

$$\lim_{\theta \rightarrow \infty} \frac{1}{\theta} \underline{\Theta}_{\text{CL}}(\theta\alpha, \theta\beta) = \underline{\Theta}_{\text{MD}}(\alpha, \beta), \quad \lim_{\theta \rightarrow \infty} \frac{1}{\theta} \overline{\Theta}_{\text{CL}}(\theta\alpha, \theta\beta) = \overline{\Theta}_{\text{MD}}(\alpha, \beta)$$


- Recall:  $\underline{\Theta}_{\text{CL}}(\alpha, \beta) := -\log \Lambda_\rho(e^{-\alpha}, e^{-\beta})$  with  $\Lambda_\rho(a, b) := \Phi_\rho(\Phi^{-1}(a), \Phi^{-1}(b))$   
(and  $\overline{\Theta}_{\text{CL}}(\alpha, \beta)$  is defined similarly with  $\rho \leftarrow -\rho$ )

# Exponents induced by Hamming Balls/Spheres for $\rho = 0.9$




$\underline{\Theta}_{\text{CL}}(\alpha, \beta)$

Zoom out  
⇒




$\underline{\Theta}_{\text{MD}}(\alpha, \beta)$

Zoom in to  
a neighborhood of the origin  
⇐




$\underline{\Theta}_{\text{LD}}(\alpha, \beta)$

# Exponents induced by Hamming Balls/Spheres for $\rho = 0.9$




$$\underline{\Theta}_{CL}(\alpha, \beta)$$

Zoom out  
⇒



$$\underline{\Theta}_{MD}(\alpha, \beta)$$


Zoom in to  
a neighborhood of the origin  
⇐



$$\underline{\Theta}_{LD}(\alpha, \beta)$$


- $\underline{\Theta}_{MD}(k\alpha, k\beta) = k\underline{\Theta}_{MD}(\alpha, \beta)$

# Exponents induced by Hamming Balls/Spheres for $\rho = 0.9$




- $\underline{\Theta}_{\text{MD}}(k\alpha, k\beta) = k\underline{\Theta}_{\text{MD}}(\alpha, \beta)$
- $\underline{\Theta}_{\text{MD}}$  contains much less information than  $\underline{\Theta}_{\text{CL}}$  and  $\underline{\Theta}_{\text{LD}}$


# Exponents induced by Hamming Balls/Spheres for $\rho = 0.9$



$\underline{\Theta}_{CL}(\alpha, \beta)$        $\xrightarrow{\text{Zoom out}}$




$\underline{\Theta}_{MD}(\alpha, \beta)$        $\xleftarrow[\text{a neighborhood of the origin}]{\text{Zoom in to}}$



$\underline{\Theta}_{LD}(\alpha, \beta)$


- $\underline{\Theta}_{MD}(k\alpha, k\beta) = k\underline{\Theta}_{MD}(\alpha, \beta)$
- $\underline{\Theta}_{MD}$  contains much less information than  $\underline{\Theta}_{CL}$  and  $\underline{\Theta}_{LD}$
- Later we will show: Both  $\underline{\Theta}_{MD}$  and  $\underline{\Theta}_{LD}$  are convex; but  $\underline{\Theta}_{CL}$  is not

# Exponents induced by Hamming Balls/Spheres for $\rho = 0.9$




$\bar{\Theta}_{CL}(\alpha, \beta)$

Zoom out  
 $\Rightarrow$




$\bar{\Theta}_{MD}(\alpha, \beta)$

Zoom in to  
a neighborhood of the origin  
 $\Leftarrow$




$\bar{\Theta}_{LD}(\alpha, \beta)$

# Exponents induced by Hamming Balls/Spheres for $\rho = 0.9$




$$\bar{\Theta}_{CL}(\alpha, \beta)$$

Zoom out  
⇒



$$\bar{\Theta}_{MD}(\alpha, \beta)$$

Zoom in to  
a neighborhood of the origin  
⇐



$$\bar{\Theta}_{LD}(\alpha, \beta)$$

- $\bar{\Theta}_{MD}(k\alpha, k\beta) = k\bar{\Theta}_{MD}(\alpha, \beta)$
- $\bar{\Theta}_{MD}$  contains much less information than  $\bar{\Theta}_{CL}$  and  $\bar{\Theta}_{LD}$
- Later we will show: Both  $\bar{\Theta}_{MD}$  and  $\bar{\Theta}_{LD}$  are **concave**; but  $\bar{\Theta}_{CL}$  is not

# Comparison: Hamming Subcubes vs. Hamming Balls/Spheres

Regime	Central Limit		Moderate Deviation	Large Deviation
$a, b$	fixed and large $a, b$	fixed but small $a, b$	subexp. vanishing $a, b$	exp. vanishing $a, b$
Subcubes	Better			
Balls		Better	Better	Better

# Comparison: Hamming Subcubes vs. Hamming Balls/Spheres


Regime	Central Limit		Moderate Deviation	Large Deviation
$a, b$	fixed and large $a, b$	fixed but small $a, b$	subexp. vanishing $a, b$	exp. vanishing $a, b$
Subcubes	Better			
Balls		Better	Better	Better

- For large  $a, b$ , subcubes are better; for small  $a, b$ , balls are better.

# Comparison: Hamming Subcubes vs. Hamming Balls/Spheres

Regime	Central Limit		Moderate Deviation	Large Deviation
$a, b$	fixed and large $a, b$	fixed but small $a, b$	subexp. vanishing $a, b$	exp. vanishing $a, b$
Subcubes	Better			
Balls		Better	Better	Better

- For large  $a, b$ , subcubes are better; for small  $a, b$ , balls are better.



# Natural Questions on Optimality

Questions: Are Hamming subcubes optimal for large  $a, b$ ? Are Hamming balls optimal for small  $a, b$ ?

# Natural Questions on Optimality

Questions: Are Hamming subcubes optimal for large  $a, b$ ? Are Hamming balls optimal for small  $a, b$ ?

- CL regime:  $a, b$  are fixed

- Are subcubes optimal for  $a = b \in \{\frac{1}{2}, \frac{1}{4}\}$ ? — including Mossel's mean-1/4 stability problem
- Are balls optimal for small but fixed  $a, b$ ?

# Natural Questions on Optimality

Questions: Are Hamming subcubes optimal for large  $a, b$ ? Are Hamming balls optimal for small  $a, b$ ?

- CL regime:  $a, b$  are fixed
  - Are subcubes optimal for  $a = b \in \{\frac{1}{2}, \frac{1}{4}\}$ ? — including Mossel's mean-1/4 stability problem
  - Are balls optimal for small but fixed  $a, b$ ?
- MD regime:  $a, b$  are “subexponentially” small
  - Are balls/spheres optimal?
- LD regime:  $a, b$  are exponentially small
  - Are balls/spheres optimal? — Ordentlich–Polyanskiy–Shayevitz's conjecture

# Natural Questions on Optimality

Questions: Are Hamming subcubes optimal for large  $a, b$ ? Are Hamming balls optimal for small  $a, b$ ?

- CL regime:  $a, b$  are fixed
  - Are subcubes optimal for  $a = b \in \{\frac{1}{2}, \frac{1}{4}\}$ ? — including Mossel's mean-1/4 stability problem
  - Are balls optimal for small but fixed  $a, b$ ?
- MD regime:  $a, b$  are “subexponentially” small
  - Are balls/spheres optimal?
- LD regime:  $a, b$  are exponentially small
  - Are balls/spheres optimal? — Ordentlich–Polyanskiy–Shayevitz's conjecture
- We answer these questions in the following.

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- **Converse Part: CL Regime with Large  $a, b$**
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

## Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

- Confirmed positively by Witsenhausen (1975) using maximal correlation

## Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

- Confirmed positively by Witsenhausen (1975) using maximal correlation
- The (Hirschfeld–Gebelein–Rényi) maximal correlation

$$\rho_m(X; Y) := \sup_{f,g} \rho(f(X); g(Y)),$$

- $\rho(U; V) := \frac{\mathbb{E}[UV]}{\sqrt{\text{var}[U]\text{var}[V]}}$  is the Pearson correlation
- the supremum is taken over all real-valued functions with finite variances

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

- Confirmed positively by Witsenhausen (1975) using maximal correlation
- The (Hirschfeld–Gebelein–Rényi) maximal correlation

$$\rho_m(X; Y) := \sup_{f, g} \rho(f(X); g(Y)),$$

- $\rho(U; V) := \frac{\mathbb{E}[UV]}{\sqrt{\text{var}[U]\text{var}[V]}}$  is the Pearson correlation
- the supremum is taken over all real-valued functions with finite variances
- Tensorization:

$$\rho_m(\mathbf{X}; \mathbf{Y}) = \rho_m(X; Y).$$

## Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

- Confirmed positively by Witsenhausen (1975) using maximal correlation
- The (Hirschfeld–Gebelein–Rényi) maximal correlation

$$\rho_m(X; Y) := \sup_{f,g} \rho(f(X); g(Y)),$$

- $\rho(U; V) := \frac{\mathbb{E}[UV]}{\sqrt{\text{var}[U]\text{var}[V]}}$  is the Pearson correlation
- the supremum is taken over all real-valued functions with finite variances
- Tensorization:  
$$\rho_m(\mathbf{X}; \mathbf{Y}) = \rho_m(X; Y).$$
- Data Processing Inequality (DPI): For a Markov chain  $U - X - Y - V$ ,

$$\rho_m(U; V) \leq \rho_m(X; Y).$$

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

- Confirmed positively by Witsenhausen (1975) using maximal correlation
- The (Hirschfeld–Gebelein–Rényi) maximal correlation

$$\rho_m(X; Y) := \sup_{f,g} \rho(f(X); g(Y)),$$

- $\rho(U; V) := \frac{\mathbb{E}[UV]}{\sqrt{\text{var}[U]\text{var}[V]}}$  is the Pearson correlation
- the supremum is taken over all real-valued functions with finite variances
- Tensorization:  
$$\rho_m(\mathbf{X}; \mathbf{Y}) = \rho_m(X; Y).$$
- Data Processing Inequality (DPI): For a Markov chain  $U - X - Y - V$ ,  
$$\rho_m(U; V) \leq \rho_m(X; Y).$$
- For binary  $X, Y$ ,  $\rho_m(X; Y) = |\rho(X; Y)|$ .

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

Theorem ([Witsenhausen, 1975])

For any  $A, B$  with  $P_X^n(A) = a, P_Y^n(B) = b$ ,

$$ab - \rho\sqrt{a\bar{a}b\bar{b}} \leq P_{XY}^n(A \times B) \leq ab + \rho\sqrt{a\bar{a}b\bar{b}}, \quad \text{where } \bar{x} = 1 - x.$$

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

Theorem ([Witsenhausen, 1975])

For any  $A, B$  with  $P_X^n(A) = a, P_Y^n(B) = b$ ,

$$ab - \rho \sqrt{a\bar{a}b\bar{b}} \leq P_{XY}^n(A \times B) \leq ab + \rho \sqrt{a\bar{a}b\bar{b}}, \quad \text{where } \bar{x} = 1 - x.$$

- Proof: Setting  $U = 1_A(\mathbf{X}), V = 1_B(\mathbf{Y}) \implies U - \mathbf{X} - \mathbf{Y} - V \implies$

$$\frac{|P_{XY}^n(A \times B) - ab|}{\sqrt{a\bar{a}}\sqrt{b\bar{b}}} = |\rho(U; V)| = \rho_m(U; V) \stackrel{\text{DPI}}{\leq} \rho_m(\mathbf{X}; \mathbf{Y}) \stackrel{\text{Tensorization}}{=} \rho_m(X; Y) = \rho.$$

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

Theorem ([Witsenhausen, 1975])

For any  $A, B$  with  $P_X^n(A) = a, P_Y^n(B) = b$ ,

$$ab - \rho\sqrt{a\bar{a}b\bar{b}} \leq P_{XY}^n(A \times B) \leq ab + \rho\sqrt{a\bar{a}b\bar{b}}, \quad \text{where } \bar{x} = 1 - x.$$

- Proof: Setting  $U = 1_A(\mathbf{X}), V = 1_B(\mathbf{Y}) \implies U - \mathbf{X} - \mathbf{Y} - V \implies$

$$\frac{|P_{XY}^n(A \times B) - ab|}{\sqrt{a\bar{a}}\sqrt{b\bar{b}}} = |\rho(U; V)| = \rho_m(U; V) \stackrel{\text{DPI}}{\leq} \rho_m(\mathbf{X}; \mathbf{Y}) \stackrel{\text{Tensorization}}{=} \rho_m(X; Y) = \rho.$$

- Important Consequence:

- For  $a = b = 1/2$ ,  $\frac{1-\rho}{4} \leq P_{XY}^n(A \times B) \leq \frac{1+\rho}{4}$ .

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

Theorem ([Witsenhausen, 1975])

For any  $A, B$  with  $P_X^n(A) = a, P_Y^n(B) = b$ ,

$$ab - \rho\sqrt{a\bar{a}b\bar{b}} \leq P_{XY}^n(A \times B) \leq ab + \rho\sqrt{a\bar{a}b\bar{b}}, \quad \text{where } \bar{x} = 1 - x.$$

- Proof: Setting  $U = 1_A(\mathbf{X}), V = 1_B(\mathbf{Y}) \implies U - \mathbf{X} - \mathbf{Y} - V \implies$

$$\frac{|P_{XY}^n(A \times B) - ab|}{\sqrt{a\bar{a}}\sqrt{b\bar{b}}} = |\rho(U; V)| = \rho_m(U; V) \stackrel{\text{DPI}}{\leq} \rho_m(\mathbf{X}; \mathbf{Y}) \stackrel{\text{Tensorization}}{=} \rho_m(X; Y) = \rho.$$

- Important Consequence:

- For  $a = b = 1/2$ ,  $\frac{1-\rho}{4} \leq P_{XY}^n(A \times B) \leq \frac{1+\rho}{4}$ .
- Upper bound is attained by  $f(\mathbf{x}) = g(\mathbf{x}) = x_i$  and lower bound by  $f(\mathbf{x}) = -g(\mathbf{x}) = x_i$

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

Theorem ([Witsenhausen, 1975])

For any  $A, B$  with  $P_X^n(A) = a, P_Y^n(B) = b$ ,

$$ab - \rho\sqrt{a\bar{a}b\bar{b}} \leq P_{XY}^n(A \times B) \leq ab + \rho\sqrt{a\bar{a}b\bar{b}}, \quad \text{where } \bar{x} = 1 - x.$$

- Proof: Setting  $U = 1_A(\mathbf{X}), V = 1_B(\mathbf{Y}) \implies U - \mathbf{X} - \mathbf{Y} - V \implies$

$$\frac{|P_{XY}^n(A \times B) - ab|}{\sqrt{a\bar{a}}\sqrt{b\bar{b}}} = |\rho(U; V)| = \rho_m(U; V) \stackrel{\text{DPI}}{\leq} \rho_m(\mathbf{X}; \mathbf{Y}) \stackrel{\text{Tensorization}}{=} \rho_m(X; Y) = \rho.$$

- Important Consequence:

- For  $a = b = 1/2$ ,  $\frac{1-\rho}{4} \leq P_{XY}^n(A \times B) \leq \frac{1+\rho}{4}$ .
- Upper bound is attained by  $f(\mathbf{x}) = g(\mathbf{x}) = x_i$  and lower bound by  $f(\mathbf{x}) = -g(\mathbf{x}) = x_i$
- Dictators (subcubes) are optimal for  $a = b = 1/2$ , i.e.,  $\bar{\Gamma}^{(n)}\left(\frac{1}{2}, \frac{1}{2}\right) = \frac{1+\rho}{4}$  and  $\underline{\Gamma}^{(n)}\left(\frac{1}{2}, \frac{1}{2}\right) = \frac{1-\rho}{4}$

# Converse for $a = b = \frac{1}{2}$ : Are subcubes (dictators) optimal?

Theorem ([Witsenhausen, 1975])

For any  $A, B$  with  $P_X^n(A) = a, P_Y^n(B) = b$ ,

$$ab - \rho\sqrt{a\bar{a}b\bar{b}} \leq P_{XY}^n(A \times B) \leq ab + \rho\sqrt{a\bar{a}b\bar{b}}, \quad \text{where } \bar{x} = 1 - x.$$

- Proof: Setting  $U = 1_A(\mathbf{X}), V = 1_B(\mathbf{Y}) \implies U - \mathbf{X} - \mathbf{Y} - V \implies$

$$\frac{|P_{XY}^n(A \times B) - ab|}{\sqrt{a\bar{a}}\sqrt{b\bar{b}}} = |\rho(U; V)| = \rho_m(U; V) \stackrel{\text{DPI}}{\leq} \rho_m(\mathbf{X}; \mathbf{Y}) \stackrel{\text{Tensorization}}{=} \rho_m(X; Y) = \rho.$$

- Important Consequence:

- For  $a = b = 1/2$ ,  $\frac{1-\rho}{4} \leq P_{XY}^n(A \times B) \leq \frac{1+\rho}{4}$ .
- Upper bound is attained by  $f(\mathbf{x}) = g(\mathbf{x}) = x_i$  and lower bound by  $f(\mathbf{x}) = -g(\mathbf{x}) = x_i$
- Dictators (subcubes) are **optimal** for  $a = b = 1/2$ , i.e.,  $\bar{\Gamma}^{(n)}\left(\frac{1}{2}, \frac{1}{2}\right) = \frac{1+\rho}{4}$  and  $\underline{\Gamma}^{(n)}\left(\frac{1}{2}, \frac{1}{2}\right) = \frac{1-\rho}{4}$
- Note: This point also can be proven by **hypercontractivity** and **Fourier analysis**

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal? — Mossel's mean-1/4 stability problem

- Confirmed positively by Yu and Tan (2021) using Fourier analysis

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal? — Mossel's mean-1/4 stability problem

- Confirmed positively by Yu and Tan (2021) using Fourier analysis
- Fourier coefficients of  $f : \{0, 1\}^n \rightarrow \{0, 1\}$  are

$$\hat{f}(\mathbf{y}) := \frac{1}{2^n} \sum_{\mathbf{x}} f(\mathbf{x}) (-1)^{\langle \mathbf{x}, \mathbf{y} \rangle}$$

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal? — Mossel's mean-1/4 stability problem

- Confirmed positively by Yu and Tan (2021) using Fourier analysis
- Fourier coefficients of  $f : \{0, 1\}^n \rightarrow \{0, 1\}$  are

$$\hat{f}(\mathbf{y}) := \frac{1}{2^n} \sum_{\mathbf{x}} f(\mathbf{x}) (-1)^{\langle \mathbf{x}, \mathbf{y} \rangle}$$

- Fourier expansion of  $f$  is

$$f(\mathbf{x}) = \sum_{\mathbf{y}} \hat{f}(\mathbf{y}) (-1)^{\langle \mathbf{x}, \mathbf{y} \rangle}$$

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal? — Mossel's mean-1/4 stability problem

- Confirmed positively by Yu and Tan (2021) using Fourier analysis
- Fourier coefficients of  $f : \{0, 1\}^n \rightarrow \{0, 1\}$  are

$$\hat{f}(\mathbf{y}) := \frac{1}{2^n} \sum_{\mathbf{x}} f(\mathbf{x}) (-1)^{\langle \mathbf{x}, \mathbf{y} \rangle}$$

- Fourier expansion of  $f$  is

$$f(\mathbf{x}) = \sum_{\mathbf{y}} \hat{f}(\mathbf{y}) (-1)^{\langle \mathbf{x}, \mathbf{y} \rangle}$$

- Define the  $k$ -degree Fourier weight as

$$\mathbf{W}_k(f) := \sum_{|\mathbf{y}|=k} \hat{f}(\mathbf{y})^2$$

where  $|\mathbf{y}|$  denotes the Hamming weight of  $\mathbf{y}$ .

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal?

- Properties: For a Boolean  $f$  with mean  $a$ ,

$$\mathbf{W}_0(f) = a^2, \quad \sum_{k=0}^n \mathbf{W}_k(f) = a, \quad \mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1) = \sum_{k=0}^n \mathbf{W}_k(f) \rho^k.$$

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal?

- Properties: For a Boolean  $f$  with mean  $a$ ,

$$\mathbf{W}_0(f) = a^2, \quad \sum_{k=0}^n \mathbf{W}_k(f) = a, \quad \mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1) = \sum_{k=0}^n \mathbf{W}_k(f) \rho^k.$$

- LP bound on  $\mathbf{W}_1$  [Fu et al., 2001, Yu and Tan, 2019]:  $\mathbf{W}_1(f) \leq \varphi(a) := \begin{cases} 2a(\sqrt{a} - a) & 0 \leq a \leq \frac{1}{4} \\ \frac{a}{2} & \frac{1}{4} < a \leq \frac{1}{2} \end{cases}$

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal?

- Properties: For a Boolean  $f$  with mean  $a$ ,

$$\mathbf{W}_0(f) = a^2, \quad \sum_{k=0}^n \mathbf{W}_k(f) = a, \quad \mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1) = \sum_{k=0}^n \mathbf{W}_k(f) \rho^k.$$

- LP bound on  $\mathbf{W}_1$  [Fu et al., 2001, Yu and Tan, 2019]:  $\mathbf{W}_1(f) \leq \varphi(a) := \begin{cases} 2a(\sqrt{a} - a) & 0 \leq a \leq \frac{1}{4} \\ \frac{a}{2} & \frac{1}{4} < a \leq \frac{1}{2} \end{cases}$
- Fact:  $\mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \leq \max \{\mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1), \mathbb{P}(g(\mathbf{X}) = g(\mathbf{Y}) = 1)\}$

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal?

- Properties: For a Boolean  $f$  with mean  $a$ ,

$$\mathbf{W}_0(f) = a^2, \quad \sum_{k=0}^n \mathbf{W}_k(f) = a, \quad \mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1) = \sum_{k=0}^n \mathbf{W}_k(f) \rho^k.$$

- LP bound on  $\mathbf{W}_1$  [Fu et al., 2001, Yu and Tan, 2019]:  $\mathbf{W}_1(f) \leq \varphi(a) := \begin{cases} 2a(\sqrt{a} - a) & 0 \leq a \leq \frac{1}{4} \\ \frac{a}{2} & \frac{1}{4} < a \leq \frac{1}{2} \end{cases}$
- Fact:  $\mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \leq \max \{\mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1), \mathbb{P}(g(\mathbf{X}) = g(\mathbf{Y}) = 1)\}$

Theorem ([Yu and Tan, 2019, Yu and Tan, 2021])

$$\bar{\Gamma}^{(n)}(a, a) \leq a^2 + \rho\varphi(a) + \rho^2(a - a^2 - \varphi(a)).$$

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal?

- Properties: For a Boolean  $f$  with mean  $a$ ,

$$\mathbf{W}_0(f) = a^2, \quad \sum_{k=0}^n \mathbf{W}_k(f) = a, \quad \mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1) = \sum_{k=0}^n \mathbf{W}_k(f) \rho^k.$$

- LP bound on  $\mathbf{W}_1$  [Fu et al., 2001, Yu and Tan, 2019]:  $\mathbf{W}_1(f) \leq \varphi(a) := \begin{cases} 2a(\sqrt{a} - a) & 0 \leq a \leq \frac{1}{4} \\ \frac{a}{2} & \frac{1}{4} < a \leq \frac{1}{2} \end{cases}$
- Fact:  $\mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \leq \max \{\mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1), \mathbb{P}(g(\mathbf{X}) = g(\mathbf{Y}) = 1)\}$

Theorem ([Yu and Tan, 2019, Yu and Tan, 2021])

$$\bar{\Gamma}^{(n)}(a, a) \leq a^2 + \rho\varphi(a) + \rho^2(a - a^2 - \varphi(a)).$$

- Consequence: For  $a = 1/4$ , the upper bound reduces to  $\left(\frac{1+\rho}{4}\right)^2 \Rightarrow \bar{\Gamma}^{(n)}\left(\frac{1}{4}, \frac{1}{4}\right) = \left(\frac{1+\rho}{4}\right)^2$  for  $n \geq 2$ , attained by subcubes

# Converse for $a = b = \frac{1}{4}$ : Are subcubes optimal?

- Properties: For a Boolean  $f$  with mean  $a$ ,

$$\mathbf{W}_0(f) = a^2, \quad \sum_{k=0}^n \mathbf{W}_k(f) = a, \quad \mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1) = \sum_{k=0}^n \mathbf{W}_k(f) \rho^k.$$

- LP bound on  $\mathbf{W}_1$  [Fu et al., 2001, Yu and Tan, 2019]:  $\mathbf{W}_1(f) \leq \varphi(a) := \begin{cases} 2a(\sqrt{a} - a) & 0 \leq a \leq \frac{1}{4} \\ \frac{a}{2} & \frac{1}{4} < a \leq \frac{1}{2} \end{cases}$
- Fact:  $\mathbb{P}(f(\mathbf{X}) = g(\mathbf{Y}) = 1) \leq \max \{\mathbb{P}(f(\mathbf{X}) = f(\mathbf{Y}) = 1), \mathbb{P}(g(\mathbf{X}) = g(\mathbf{Y}) = 1)\}$

Theorem ([Yu and Tan, 2019, Yu and Tan, 2021])

$$\bar{\Gamma}^{(n)}(a, a) \leq a^2 + \rho\varphi(a) + \rho^2(a - a^2 - \varphi(a)).$$

- Consequence: For  $a = 1/4$ , the upper bound reduces to  $\left(\frac{1+\rho}{4}\right)^2 \Rightarrow \bar{\Gamma}^{(n)}\left(\frac{1}{4}, \frac{1}{4}\right) = \left(\frac{1+\rho}{4}\right)^2$  for  $n \geq 2$ , attained by subcubes
- However,  $\underline{\Gamma}^{(n)}\left(\frac{1}{4}, \frac{1}{4}\right)$  is still open!

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- **Converse Part: MD Regime and CL Regime with Small  $a, b$**
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Converse for MD and CL with Small $a, b$ : Small-Set Expansion (SSE) Theorem

Theorem (Small-Set Expansion [Ahlswede and Gács, 1976, Kahn et al., 1988, Mossel et al., 2006, O'Donnell, 2014])

For any  $n \geq 1$  and  $\alpha, \beta > 0$ ,

$$\underline{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) \geq \underline{\Theta}_{\text{MD}}(\alpha, \beta),$$
$$\overline{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) \leq \overline{\Theta}_{\text{MD}}(\alpha, \beta).$$

- Recall:

$$\underline{\Theta}_{\text{MD}}(\alpha, \beta) = \begin{cases} \frac{\alpha + \beta - 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}, & \rho^2\alpha \leq \beta \leq \frac{\alpha}{\rho^2}, \\ \alpha, & \beta < \rho^2\alpha, \\ \beta, & \alpha < \rho^2\beta \end{cases}, \quad \overline{\Theta}_{\text{MD}}(\alpha, \beta) = \frac{\alpha + \beta + 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}.$$

# Converse for MD and CL with Small $a, b$ : Small-Set Expansion (SSE) Theorem

Theorem (Small-Set Expansion [Ahlswede and Gács, 1976, Kahn et al., 1988, Mossel et al., 2006, O'Donnell, 2014])

For any  $n \geq 1$  and  $\alpha, \beta > 0$ ,

$$\underline{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) \geq \underline{\Theta}_{\text{MD}}(\alpha, \beta),$$
$$\overline{\Theta}_{\text{MD}}^{(n)}(\alpha, \beta) \leq \overline{\Theta}_{\text{MD}}(\alpha, \beta).$$

- Recall:

$$\underline{\Theta}_{\text{MD}}(\alpha, \beta) = \begin{cases} \frac{\alpha + \beta - 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}, & \rho^2\alpha \leq \beta \leq \frac{\alpha}{\rho^2}, \\ \alpha, & \beta < \rho^2\alpha, \\ \beta, & \alpha < \rho^2\beta \end{cases}, \quad \overline{\Theta}_{\text{MD}}(\alpha, \beta) = \frac{\alpha + \beta + 2\rho\sqrt{\alpha\beta}}{1 - \rho^2}.$$

- Recall:  $\underline{\Theta}_{\text{MD}}(\alpha, \beta)$ ,  $\overline{\Theta}_{\text{MD}}(\alpha, \beta)$  can be achieved by spheres/balls

# Converse for MD and CL with Small $a, b$ : SSE Theorem

Balls/spheres are optimal in MD regime!

# Converse for MD and CL with Small $a, b$ : SSE Theorem

Balls/spheres are optimal in MD regime!

- By the equivalence of CL, MD, and LD exponents, SSE Theorem can be rewritten as

$$\begin{aligned}\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) &\geq \underline{\Theta}_{\text{MD}}(\alpha, \beta), \\ \overline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) &\leq \overline{\Theta}_{\text{MD}}(\alpha, \beta).\end{aligned}$$

# Converse for MD and CL with Small $a, b$ : SSE Theorem

Balls/spheres are optimal in MD regime!

- By the equivalence of CL, MD, and LD exponents, SSE Theorem can be rewritten as

$$\begin{aligned}\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) &\geq \underline{\Theta}_{\text{MD}}(\alpha, \beta), \\ \overline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) &\leq \overline{\Theta}_{\text{MD}}(\alpha, \beta).\end{aligned}$$

- Proof of SSE Theorem: Based on the classic [hypercontractivity](#) inequalities [Bonami, 1968, Gross, 1975, Borell, 1982]:

# Converse for MD and CL with Small $a, b$ : SSE Theorem

Balls/spheres are optimal in MD regime!

- By the equivalence of CL, MD, and LD exponents, SSE Theorem can be rewritten as

$$\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \geq \underline{\Theta}_{\text{MD}}(\alpha, \beta),$$
$$\overline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \leq \overline{\Theta}_{\text{MD}}(\alpha, \beta).$$

- Proof of SSE Theorem: Based on the classic **hypercontractivity** inequalities [Bonami, 1968, Gross, 1975, Borell, 1982]:
  - Forward part: For  $p, q \geq 1$ , s.t.  $(p - 1)(q - 1) \geq \rho^2$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q, \quad \forall f : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}, g : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$$

where  $\|f(\mathbf{X})\|_p := (\mathbb{E}[f^p(\mathbf{X})])^{1/p}$  and  $\|g(\mathbf{Y})\|_q := (\mathbb{E}[g^q(\mathbf{Y})])^{1/q}$ .

# Converse for MD and CL with Small $a, b$ : SSE Theorem

Balls/spheres are optimal in MD regime!

- By the equivalence of CL, MD, and LD exponents, SSE Theorem can be rewritten as

$$\underline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \geq \underline{\Theta}_{\text{MD}}(\alpha, \beta),$$
$$\overline{\Theta}_{\text{CL}}^{(n)}(\alpha, \beta), \overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \leq \overline{\Theta}_{\text{MD}}(\alpha, \beta).$$

- Proof of SSE Theorem: Based on the classic **hypercontractivity** inequalities [Bonami, 1968, Gross, 1975, Borell, 1982]:

- Forward part: For  $p, q \geq 1$ , s.t.  $(p - 1)(q - 1) \geq \rho^2$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q, \quad \forall f : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}, g : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$$

where  $\|f(\mathbf{X})\|_p := (\mathbb{E}[f^p(\mathbf{X})])^{1/p}$  and  $\|g(\mathbf{Y})\|_q := (\mathbb{E}[g^q(\mathbf{Y})])^{1/q}$ .

- Substituting  $f \leftarrow 1_A, g \leftarrow 1_B$  and optimizing  $p, q$ , SSE Theorem follows.

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- **Converse Part: LD Regime**
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Converse for LD: Strong SSE Theorem

Theorem (Strong SSE [Yu et al., 2021, Yu, 2021c])

For any  $n \geq 1$  and  $\alpha, \beta \in (0, 1]$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \geq \mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta),$$
$$\overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \leq \mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta),$$

where  $\mathfrak{R}[f], \mathfrak{C}[f]$  respectively denote the lower convex and upper concave envelopes of a function  $f$ .

# Converse for LD: Strong SSE Theorem

Theorem (Strong SSE [Yu et al., 2021, Yu, 2021c])

For any  $n \geq 1$  and  $\alpha, \beta \in (0, 1]$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \geq \mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta),$$
$$\overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \leq \mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta),$$

where  $\mathfrak{R}[f]$ ,  $\mathfrak{C}[f]$  respectively denote the lower convex and upper concave envelopes of a function  $f$ .

- Why is it “Strong”?

# Converse for LD: Strong SSE Theorem

Theorem (Strong SSE [Yu et al., 2021, Yu, 2021c])

For any  $n \geq 1$  and  $\alpha, \beta \in (0, 1]$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \geq \mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta),$$
$$\overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \leq \mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta),$$

where  $\mathfrak{R}[f], \mathfrak{C}[f]$  respectively denote the lower convex and upper concave envelopes of a function  $f$ .

- Why is it “Strong”?
  - $\mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta) \geq \underline{\Theta}_{\text{MD}}(\alpha, \beta)$  and  $\mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta) \leq \overline{\Theta}_{\text{MD}}(\alpha, \beta) \implies$  stronger than SSE Theorem

# Converse for LD: Strong SSE Theorem

Theorem (Strong SSE [Yu et al., 2021, Yu, 2021c])

For any  $n \geq 1$  and  $\alpha, \beta \in (0, 1]$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \geq \mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta),$$
$$\overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \leq \mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta),$$

where  $\mathfrak{R}[f], \mathfrak{C}[f]$  respectively denote the lower convex and upper concave envelopes of a function  $f$ .

- Why is it “Strong”?
  - $\mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta) \geq \underline{\Theta}_{\text{MD}}(\alpha, \beta)$  and  $\mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta) \leq \overline{\Theta}_{\text{MD}}(\alpha, \beta) \implies$  stronger than SSE Theorem
- Recall:  $\underline{\Theta}_{\text{LD}}(\alpha, \beta), \overline{\Theta}_{\text{LD}}(\alpha, \beta)$  can be achieved by spheres/balls

# Converse for LD: Strong SSE Theorem

Theorem (Strong SSE [Yu et al., 2021, Yu, 2021c])

For any  $n \geq 1$  and  $\alpha, \beta \in (0, 1]$ ,

$$\underline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \geq \mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta),$$
$$\overline{\Theta}_{\text{LD}}^{(n)}(\alpha, \beta) \leq \mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta),$$

where  $\mathfrak{R}[f], \mathfrak{C}[f]$  respectively denote the lower convex and upper concave envelopes of a function  $f$ .

- Why is it “Strong”?
  - $\mathfrak{R}[\underline{\Theta}_{\text{LD}}](\alpha, \beta) \geq \underline{\Theta}_{\text{MD}}(\alpha, \beta)$  and  $\mathfrak{C}[\overline{\Theta}_{\text{LD}}](\alpha, \beta) \leq \overline{\Theta}_{\text{MD}}(\alpha, \beta) \implies$  stronger than SSE Theorem
- Recall:  $\underline{\Theta}_{\text{LD}}(\alpha, \beta), \overline{\Theta}_{\text{LD}}(\alpha, \beta)$  can be achieved by spheres/balls
- Consequence: Time-sharing certain spheres/balls is optimal in LD regime! — A weaker version of OPS’s conjecture

# Converse for LD: Strong SSE Theorem

Lemma ([Yu, 2021a])

$\underline{\Theta}_{\text{LD}}$  is convex, and  $\overline{\Theta}_{\text{LD}}$  is concave.

# Converse for LD: Strong SSE Theorem

Lemma ([Yu, 2021a])

$\underline{\Theta}_{LD}$  is convex, and  $\bar{\Theta}_{LD}$  is concave.

- $\implies \mathfrak{R}[\underline{\Theta}_{LD}] = \underline{\Theta}_{LD}$  and  $\mathfrak{C}[\bar{\Theta}_{LD}] = \bar{\Theta}_{LD}$ .

# Converse for LD: Strong SSE Theorem

Lemma ([Yu, 2021a])

$\underline{\Theta}_{LD}$  is convex, and  $\bar{\Theta}_{LD}$  is concave.

- $\Rightarrow \mathfrak{R}[\underline{\Theta}_{LD}] = \underline{\Theta}_{LD}$  and  $\mathfrak{C}[\bar{\Theta}_{LD}] = \bar{\Theta}_{LD}$ .
- Substituting these to Strong SSE Theorem  $\Rightarrow$

# Converse for LD: Strong SSE Theorem

Lemma ([Yu, 2021a])

$\underline{\Theta}_{\text{LD}}$  is convex, and  $\bar{\Theta}_{\text{LD}}$  is concave.

- $\Rightarrow \mathfrak{R}[\underline{\Theta}_{\text{LD}}] = \underline{\Theta}_{\text{LD}}$  and  $\mathfrak{C}[\bar{\Theta}_{\text{LD}}] = \bar{\Theta}_{\text{LD}}$ .
- Substituting these to Strong SSE Theorem  $\Rightarrow$

OPS's conjecture is true: Balls/spheres are optimal in LD regime!

# Converse for LD: Strong SSE Theorem

Lemma ([Yu, 2021a])

$\underline{\Theta}_{\text{LD}}$  is convex, and  $\bar{\Theta}_{\text{LD}}$  is concave.

- $\Rightarrow \mathfrak{R}[\underline{\Theta}_{\text{LD}}] = \underline{\Theta}_{\text{LD}}$  and  $\mathfrak{C}[\bar{\Theta}_{\text{LD}}] = \bar{\Theta}_{\text{LD}}$ .
- Substituting these to Strong SSE Theorem  $\Rightarrow$

OPS's conjecture is true: Balls/spheres are optimal in LD regime!

- Notice:
  - The limiting cases as  $\rho \rightarrow 0$  or  $1$  were previously proven in [Ordentlich et al., 2020].
  - The special case with  $\alpha = \beta$  was previously proven in [Kirshner and Samorodnitsky, 2021].

# Summary of Converse Parts

Bounds	Central Limit		Moderate Deviation	Large Deviation
	fixed and large $a, b$	fixed but small $a, b$	subexp. vanishing $a, b$	exp. vanishing $a, b$
Maximal Correlation	Sharp for $a = b = 1/2$ (Subcubes)			
Fourier Analysis	Sharp for $a = b = 1/2$ and $a = b = 1/4$ (Subcubes)			
SSE		Almost sharp (Balls)	Sharp (Balls/Spheres)	
Strong SSE				Sharp (Balls/Spheres)

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- **Connection to Hypercontractivity**
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Hypercontractivity (HC) $\implies$ SSE

- Forward HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

# Hypercontractivity (HC) $\implies$ SSE

- Forward HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the Hölder inequality for which  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

# Hypercontractivity (HC) $\implies$ SSE

- Forward HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the Hölder inequality for which  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

- Reverse HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \geq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \leq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

# Hypercontractivity (HC) $\implies$ SSE

- Forward HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the Hölder inequality for which  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

- Reverse HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \geq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \leq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the reverse Hölder inequality for which  $p, q \leq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

# Hypercontractivity (HC) $\implies$ SSE

- Forward HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the Hölder inequality for which  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

- Reverse HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \geq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \leq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the reverse Hölder inequality for which  $p, q \leq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

- HC  $\implies$  SSE: By setting  $f \leftarrow 1_A, g \leftarrow 1_B$ ,

$$P_{XY}^n(A \times B) \leq P_X^n(A)^{1/p} P_Y^n(B)^{1/q},$$
$$P_{XY}^n(A \times B) \geq P_X^n(A)^{1/p} P_Y^n(B)^{1/q}.$$

# Hypercontractivity (HC) $\implies$ SSE

- Forward HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \leq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the Hölder inequality for which  $p, q \geq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

- Reverse HC: For any nonnegative  $f, g$ ,

$$\mathbb{E}[f(\mathbf{X})g(\mathbf{Y})] \geq \|f(\mathbf{X})\|_p \|g(\mathbf{Y})\|_q$$

where  $p, q \leq 1$  s.t.  $(p - 1)(q - 1) \geq \rho^2$

- strengthening the reverse Hölder inequality for which  $p, q \leq 1$  s.t.  $(p - 1)(q - 1) \geq 1$

- HC  $\implies$  SSE: By setting  $f \leftarrow 1_A, g \leftarrow 1_B$ ,

$$P_{XY}^n(A \times B) \leq P_X^n(A)^{1/p} P_Y^n(B)^{1/q},$$
$$P_{XY}^n(A \times B) \geq P_X^n(A)^{1/p} P_Y^n(B)^{1/q}.$$

- SSE  $\implies$  HC?

# SSE $\Rightarrow$ HC

- Yes!

# SSE $\implies$ HC

- Yes!
- The “level-partition” technique [Kirshner and Samorodnitsky, 2021]:

$$f \approx \sum_{i=1}^M c_i 1_{A_i}, \quad g \approx \sum_{j=1}^N d_j 1_{B_j}$$

# SSE $\implies$ HC

- Yes!
- The “level-partition” technique [Kirshner and Samorodnitsky, 2021]:

$$f \approx \sum_{i=1}^M c_i 1_{A_i}, \quad g \approx \sum_{j=1}^N d_j 1_{B_j}$$

- Roughly speaking, applying SSE to each pair of  $(A_i, B_j) \implies \text{HC}$

# SSE $\implies$ HC

- Yes!
- The “level-partition” technique [Kirshner and Samorodnitsky, 2021]:

$$f \approx \sum_{i=1}^M c_i 1_{A_i}, \quad g \approx \sum_{j=1}^N d_j 1_{B_j}$$

- Roughly speaking, applying SSE to each pair of  $(A_i, B_j) \implies \text{HC}$
- Strong SSE + “level-partition” technique  $\implies$  Strong HC [Kirshner and Samorodnitsky, 2021, Yu et al., 2021]
  - Strong HC: A class of HC inequalities with improved  $(p, q)$  such that they are exponentially sharp for nonnegative functions with exponentially small supports

# SSE $\implies$ HC

- Yes!
- The “level-partition” technique [Kirshner and Samorodnitsky, 2021]:

$$f \approx \sum_{i=1}^M c_i 1_{A_i}, \quad g \approx \sum_{j=1}^N d_j 1_{B_j}$$

- Roughly speaking, applying SSE to each pair of  $(A_i, B_j) \implies \text{HC}$
- Strong SSE + “level-partition” technique  $\implies$  Strong HC [Kirshner and Samorodnitsky, 2021, Yu et al., 2021]
  - Strong HC: A class of HC inequalities with improved  $(p, q)$  such that they are exponentially sharp for nonnegative functions with exponentially small supports

(Strong) SSE  $\iff$  (Strong) HC (in some sense)!

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Extension to Gaussian Sources

- How about Gaussian sources?

# Extension to Gaussian Sources

- How about Gaussian sources?
- Completely solved

# Extension to Gaussian Sources

- How about Gaussian sources?
- Completely solved

Theorem (Borell's Isoperimetric Theorem [Borell, 1985, Mossel and Neeman, 2015])

Let  $(\mathbf{X}, \mathbf{Y})$  be a bivariate Gaussian source with corr.  $\rho \in (0, 1)$ . Then, for any  $a, b \in [0, 1]$ ,

$$\bar{\Gamma}^{(n)}(a, b) = \Lambda_{\rho}(a, b)$$

$$\underline{\Gamma}^{(n)}(a, b) = \Lambda_{-\rho}(a, b).$$

# Extension to Gaussian Sources

- How about Gaussian sources?
- Completely solved

Theorem (Borell's Isoperimetric Theorem [Borell, 1985, Mossel and Neeman, 2015])

Let  $(\mathbf{X}, \mathbf{Y})$  be a bivariate Gaussian source with corr.  $\rho \in (0, 1)$ . Then, for any  $a, b \in [0, 1]$ ,

$$\bar{\Gamma}^{(n)}(a, b) = \Lambda_{\rho}(a, b)$$

$$\underline{\Gamma}^{(n)}(a, b) = \Lambda_{-\rho}(a, b).$$

- Recall: Bivariate normal copula:

$$\Lambda_{\rho}(a, b) := \Phi_{\rho}(\Phi^{-1}(a), \Phi^{-1}(b))$$

# Extension to Gaussian Sources

- How about Gaussian sources?
- Completely solved

Theorem (Borell's Isoperimetric Theorem [Borell, 1985, Mossel and Neeman, 2015])

Let  $(\mathbf{X}, \mathbf{Y})$  be a bivariate Gaussian source with corr.  $\rho \in (0, 1)$ . Then, for any  $a, b \in [0, 1]$ ,

$$\bar{\Gamma}^{(n)}(a, b) = \Lambda_{\rho}(a, b)$$

$$\underline{\Gamma}^{(n)}(a, b) = \Lambda_{-\rho}(a, b).$$

- Recall: Bivariate normal copula:

$$\Lambda_{\rho}(a, b) := \Phi_{\rho}(\Phi^{-1}(a), \Phi^{-1}(b))$$

- Parallel half-spaces are optimal (e.g.,  $f = 1_{\{x_1 \leq r\}}$ ,  $g = 1_{\{y_1 \leq s\}}$ )

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users


- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# NICD with $k$ Users

$$\mathbf{Y} \sim \text{Bern}^n\left(\frac{1}{2}\right)$$



Asymmetric Version:  $\max \mathbb{P}(U_1 = U_2 = \dots = U_k = 1)$

Symmetric Version:  $\max \mathbb{P}(U_1 = U_2 = \dots = U_k)$

# $k$ -User NICD and $q$ -Stability: Asymmetric Version

- $k$ -User NICD: symmetric version in [Mossel and O'Donnell, 2005] and asymmetric version in [Li and Médard, 2021]

## $k$ -User NICD and $q$ -Stability: Asymmetric Version

- $k$ -User NICD: symmetric version in [Mossel and O'Donnell, 2005] and asymmetric version in [Li and Médard, 2021]
- Asymmetric NICD: For  $a = \frac{M}{2^n}$  (with integer  $M$ ), define **(asymmetric) max  $k$ -stability** (or forward joint probability) as

$$\begin{aligned}\bar{\Gamma}_k(a) &:= \max_{\text{Boolean } f_i, 1 \leq i \leq k : \mathbb{E}[f_i(\mathbf{X}_i)] = a} \mathbb{P}(f_1(\mathbf{X}_1) = f_2(\mathbf{X}_2) = \dots = f_k(\mathbf{X}_k) = 1) \\ &= \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X}) = a} \mathbb{E}_{\mathbf{Y}} \left[ (T_\rho f(\mathbf{Y}))^k \right] \\ &= \max_{A : P_X^n(A) = a} \mathbb{E}_{\mathbf{Y}} \left[ P_{X|Y}^n(A|\mathbf{Y})^k \right]\end{aligned}$$

where the noise operator  $T_\rho$  is given by  $T_\rho f(\mathbf{y}) := \mathbb{E} [f(\mathbf{X}) | \mathbf{Y} = \mathbf{y}]$

# $k$ -User NICD and $q$ -Stability: Asymmetric Version

- $k$ -User NICD: symmetric version in [Mossel and O'Donnell, 2005] and asymmetric version in [Li and Médard, 2021]
- Asymmetric NICD: For  $a = \frac{M}{2^n}$  (with integer  $M$ ), define **(asymmetric) max  $k$ -stability** (or forward joint probability) as

$$\begin{aligned}\bar{\Gamma}_k(a) &:= \max_{\text{Boolean } f_i, 1 \leq i \leq k : \mathbb{E}[f_i(\mathbf{X}_i)] = a} \mathbb{P}(f_1(\mathbf{X}_1) = f_2(\mathbf{X}_2) = \dots = f_k(\mathbf{X}_k) = 1) \\ &= \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X}) = a} \mathbb{E}_{\mathbf{Y}} \left[ (T_\rho f(\mathbf{Y}))^k \right] \\ &= \max_{A : P_X^n(A) = a} \mathbb{E}_{\mathbf{Y}} \left[ P_{X|Y}^n(A|\mathbf{Y})^k \right]\end{aligned}$$

where the noise operator  $T_\rho$  is given by  $T_\rho f(\mathbf{y}) := \mathbb{E} [f(\mathbf{X}) | \mathbf{Y} = \mathbf{y}]$

- Generalization to **(asymmetric) max  $q$ -stability** (or max  $\alpha$ -stability) [Li and Médard, 2021]: For  $q \in [1, \infty)$ ,

$$\bar{\Gamma}_q(a) := \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X}) = a} \mathbb{E}_{\mathbf{Y}} \left[ T_\rho f(\mathbf{Y})^q \right] = \max_{A : P_X^n(A) = a} \mathbb{E}_{\mathbf{Y}} \left[ P_{X|Y}^n(A|\mathbf{Y})^q \right]$$

# $k$ -User NICD and $q$ -Stability: Symmetric Version

- Symmetric NICD: Define **symmetric max  $k$ -stability** as (removing “= 1”)

$$\begin{aligned}\bar{\Gamma}_k^{\text{sym}}(a) &:= \max_{\text{Boolean } f_i, 1 \leq i \leq k : \mathbb{P}(f_i(\mathbf{X}_i)=1)=a} \mathbb{P}(f_1(\mathbf{X}_1) = f_2(\mathbf{X}_2) = \dots = f_k(\mathbf{X}_k)) \\ &= \max_{\text{Boolean } f : \mathbb{E} f(\mathbf{X})=a} \mathbb{E}_{\mathbf{Y}} \left[ (T_\rho f(\mathbf{Y}))^k + (1 - T_\rho f(\mathbf{Y}))^k \right]\end{aligned}$$

# $k$ -User NICD and $q$ -Stability: Symmetric Version

- Symmetric NICD: Define **symmetric max  $k$ -stability** as (removing “= 1”)

$$\begin{aligned}\bar{\Gamma}_k^{\text{sym}}(a) &:= \max_{\text{Boolean } f_i, 1 \leq i \leq k : \mathbb{P}(f_i(\mathbf{X}_i)=1)=a} \mathbb{P}(f_1(\mathbf{X}_1) = f_2(\mathbf{X}_2) = \dots = f_k(\mathbf{X}_k)) \\ &= \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E}_{\mathbf{Y}} \left[ (T_{\rho}f(\mathbf{Y}))^k + (1 - T_{\rho}f(\mathbf{Y}))^k \right]\end{aligned}$$

- Generalization to **symmetric max  $q$ -stability**: For  $q \in [1, \infty)$ ,

$$\bar{\Gamma}_q^{\text{sym}}(a) := \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E}_{\mathbf{Y}} \left[ (T_{\rho}f(\mathbf{Y}))^q + (1 - T_{\rho}f(\mathbf{Y}))^q \right]$$

# $k$ -User NICD and $q$ -Stability: Symmetric Version

- Symmetric NICD: Define **symmetric max  $k$ -stability** as (removing “= 1”)

$$\begin{aligned}\bar{\Gamma}_k^{\text{sym}}(a) &:= \max_{\text{Boolean } f_i, 1 \leq i \leq k : \mathbb{P}(f_i(\mathbf{X}_i)=1)=a} \mathbb{P}(f_1(\mathbf{X}_1) = f_2(\mathbf{X}_2) = \dots = f_k(\mathbf{X}_k)) \\ &= \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E}_{\mathbf{Y}} \left[ (T_{\rho}f(\mathbf{Y}))^k + (1 - T_{\rho}f(\mathbf{Y}))^k \right]\end{aligned}$$

- Generalization to **symmetric max  $q$ -stability**: For  $q \in [1, \infty)$ ,

$$\bar{\Gamma}_q^{\text{sym}}(a) := \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E}_{\mathbf{Y}} \left[ (T_{\rho}f(\mathbf{Y}))^q + (1 - T_{\rho}f(\mathbf{Y}))^q \right]$$

- For  $q = 1$ , trivial: For any Boolean  $f$  s.t.  $\mathbb{E}f(\mathbf{X}) = a$ ,

$$\mathbb{E}_{\mathbf{Y}} [T_{\rho}f(\mathbf{Y})] = a, \quad \mathbb{E}_{\mathbf{Y}} [(T_{\rho}f(\mathbf{Y})) + (1 - T_{\rho}f(\mathbf{Y}))] = 1$$

# $k$ -User NICD and $q$ -Stability: Symmetric Version

- Symmetric NICD: Define **symmetric max  $k$ -stability** as (removing “= 1”)

$$\begin{aligned}\bar{\Gamma}_k^{\text{sym}}(a) &:= \max_{\text{Boolean } f_i, 1 \leq i \leq k : \mathbb{P}(f_i(\mathbf{X}_i)=1)=a} \mathbb{P}(f_1(\mathbf{X}_1) = f_2(\mathbf{X}_2) = \dots = f_k(\mathbf{X}_k)) \\ &= \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E}_{\mathbf{Y}} \left[ (T_{\rho}f(\mathbf{Y}))^k + (1 - T_{\rho}f(\mathbf{Y}))^k \right]\end{aligned}$$

- Generalization to **symmetric max  $q$ -stability**: For  $q \in [1, \infty)$ ,

$$\bar{\Gamma}_q^{\text{sym}}(a) := \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E}_{\mathbf{Y}} \left[ (T_{\rho}f(\mathbf{Y}))^q + (1 - T_{\rho}f(\mathbf{Y}))^q \right]$$

- For  $q = 1$ , trivial: For any Boolean  $f$  s.t.  $\mathbb{E}f(\mathbf{X}) = a$ ,

$$\mathbb{E}_{\mathbf{Y}} [T_{\rho}f(\mathbf{Y})] = a, \quad \mathbb{E}_{\mathbf{Y}} [(T_{\rho}f(\mathbf{Y})) + (1 - T_{\rho}f(\mathbf{Y}))] = 1$$

- Better definitions of asymmetric and symmetric max  $q$ -stability?

# Variants of max $q$ -Stability

- Define

$$\Phi_q(t) := \begin{cases} \frac{t^q - t}{q-1} & q > 1 \\ t \log t & q = 1 \end{cases} \quad \text{and} \quad \Phi_q^{\text{sym}}(t) := \Phi_q(t) + \Phi_q(1-t) = \begin{cases} \frac{t^q + (1-t)^q - 1}{q-1} & q > 1 \\ t \log t + (1-t) \log (1-t) & q = 1 \end{cases}$$

# Variants of max $q$ -Stability

- Define

$$\Phi_q(t) := \begin{cases} \frac{t^q - t}{q-1} & q > 1 \\ t \log t & q = 1 \end{cases} \quad \text{and} \quad \Phi_q^{\text{sym}}(t) := \Phi_q(t) + \Phi_q(1-t) = \begin{cases} \frac{t^q + (1-t)^q - 1}{q-1} & q > 1 \\ t \log t + (1-t) \log (1-t) & q = 1 \end{cases}$$

- Variants of max  $q$ -stability:

$$\overline{\Pi}_q(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q(T_\rho f(\mathbf{Y}))], \quad \overline{\Pi}_q^{\text{sym}}(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q^{\text{sym}}(T_\rho f(\mathbf{Y}))]$$

# Variants of max $q$ -Stability

- Define

$$\Phi_q(t) := \begin{cases} \frac{t^q - t}{q-1} & q > 1 \\ t \log t & q = 1 \end{cases} \quad \text{and} \quad \Phi_q^{\text{sym}}(t) := \Phi_q(t) + \Phi_q(1-t) = \begin{cases} \frac{t^q + (1-t)^q - 1}{q-1} & q > 1 \\ t \log t + (1-t) \log (1-t) & q = 1 \end{cases}$$

- Variants of max  $q$ -stability:

$$\overline{\Pi}_q(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q(T_\rho f(\mathbf{Y}))], \quad \overline{\Pi}_q^{\text{sym}}(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q^{\text{sym}}(T_\rho f(\mathbf{Y}))]$$

- In contrast, for original max  $q$ -stability,  $\Phi_q(t) \leftarrow t^q$  and  $\Phi_q^{\text{sym}}(t) \leftarrow t^q + (1-t)^q$

# Variants of max $q$ -Stability

- Define

$$\Phi_q(t) := \begin{cases} \frac{t^q - t}{q-1} & q > 1 \\ t \log t & q = 1 \end{cases} \quad \text{and} \quad \Phi_q^{\text{sym}}(t) := \Phi_q(t) + \Phi_q(1-t) = \begin{cases} \frac{t^q + (1-t)^q - 1}{q-1} & q > 1 \\ t \log t + (1-t) \log (1-t) & q = 1 \end{cases}$$

- Variants of max  $q$ -stability:

$$\bar{\Pi}_q(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q(T_\rho f(\mathbf{Y}))], \quad \bar{\Pi}_q^{\text{sym}}(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q^{\text{sym}}(T_\rho f(\mathbf{Y}))]$$

- In contrast, for original max  $q$ -stability,  $\Phi_q(t) \leftarrow t^q$  and  $\Phi_q^{\text{sym}}(t) \leftarrow t^q + (1-t)^q$
- For  $q > 1$ ,  $\bar{\Pi}_q(a) \iff \bar{\Gamma}_q(a)$  and  $\bar{\Pi}_q^{\text{sym}}(a) \iff \bar{\Gamma}_q^{\text{sym}}(a)$

# Variants of max $q$ -Stability

- Define

$$\Phi_q(t) := \begin{cases} \frac{t^q - t}{q-1} & q > 1 \\ t \log t & q = 1 \end{cases} \quad \text{and} \quad \Phi_q^{\text{sym}}(t) := \Phi_q(t) + \Phi_q(1-t) = \begin{cases} \frac{t^q + (1-t)^q - 1}{q-1} & q > 1 \\ t \log t + (1-t) \log (1-t) & q = 1 \end{cases}$$

- Variants of max  $q$ -stability:

$$\bar{\Pi}_q(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q(T_\rho f(\mathbf{Y}))], \quad \bar{\Pi}_q^{\text{sym}}(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q^{\text{sym}}(T_\rho f(\mathbf{Y}))]$$

- In contrast, for original max  $q$ -stability,  $\Phi_q(t) \leftarrow t^q$  and  $\Phi_q^{\text{sym}}(t) \leftarrow t^q + (1-t)^q$
- For  $q > 1$ ,  $\bar{\Pi}_q(a) \iff \bar{\Gamma}_q(a)$  and  $\bar{\Pi}_q^{\text{sym}}(a) \iff \bar{\Gamma}_q^{\text{sym}}(a)$
- For  $q = 1$ ,

$$\bar{\Pi}_q^{\text{sym}}(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} -H(f(\mathbf{X}) | \mathbf{Y}) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} I(f(\mathbf{X}) ; \mathbf{Y}) - H_2(a)$$

# Variants of max $q$ -Stability

- Define

$$\Phi_q(t) := \begin{cases} \frac{t^q - t}{q-1} & q > 1 \\ t \log t & q = 1 \end{cases} \quad \text{and} \quad \Phi_q^{\text{sym}}(t) := \Phi_q(t) + \Phi_q(1-t) = \begin{cases} \frac{t^q + (1-t)^q - 1}{q-1} & q > 1 \\ t \log t + (1-t) \log(1-t) & q = 1 \end{cases}$$

- Variants of max  $q$ -stability:

$$\bar{\Pi}_q(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q(T_\rho f(\mathbf{Y}))], \quad \bar{\Pi}_q^{\text{sym}}(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} \mathbb{E} [\Phi_q^{\text{sym}}(T_\rho f(\mathbf{Y}))]$$

- In contrast, for original max  $q$ -stability,  $\Phi_q(t) \leftarrow t^q$  and  $\Phi_q^{\text{sym}}(t) \leftarrow t^q + (1-t)^q$
- For  $q > 1$ ,  $\bar{\Pi}_q(a) \iff \bar{\Gamma}_q(a)$  and  $\bar{\Pi}_q^{\text{sym}}(a) \iff \bar{\Gamma}_q^{\text{sym}}(a)$
- For  $q = 1$ ,

$$\bar{\Pi}_q^{\text{sym}}(a) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} -H(f(\mathbf{X}) | \mathbf{Y}) = \max_{\text{Boolean } f : \mathbb{E}f(\mathbf{X})=a} I(f(\mathbf{X}); \mathbf{Y}) - H_2(a)$$

- Attained by dictators for  $a = 1/2$ ? — Courtade–Kumar Conjecture (Balanced Function Version) (2013)

# Max $q$ -Stability: Interesting Cases

- Balanced case (CL regime):  $a = 1/2$
- LD regime:  $a = 2^{-n\alpha}$  is exponentially small
- MD regime:  $a = 2^{-\theta_n \alpha}$  is “subexponentially” small with  $\theta_n \rightarrow \infty$ ,  $\frac{\theta_n}{n} \rightarrow 0$

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- **Balanced Case**
- MD and LD Regimes

# Related Conjectures for the Balanced Case

$q$	Are <b>dictators</b> optimal for $a = 1/2$ ?
$q = 1$	<a href="#">Courtade–Kumar Conjecture (2013) (balanced version)</a>
$1 < q < 2$	<a href="#">Li–Médard Conjecture (2019)</a>
$q = 2$ (2-User NICD)	True [Witsenhausen, 1975]
$q > 2$	<a href="#">Mossel–O’Donnell Conjecture for <math>2 &lt; q \leq 9</math> (2006)</a>

# $q$ -Stability: Balanced Case


Lemma ([Barnes and Özgür, 2020])

For  $a = 1/2$ , there are two **thresholds**  $1 \leq q_{\min} < 2 < q_{\max}$  s.t. dictators are optimal if and only if  $q \in [q_{\min}, q_{\max}]$ .

# $q$ -Stability: Balanced Case

Lemma ([Barnes and Özgür, 2020])

For  $a = 1/2$ , there are two **thresholds**  $1 \leq q_{\min} < 2 < q_{\max}$  s.t. dictators are optimal if and only if  $q \in [q_{\min}, q_{\max}]$ .



# $q$ -Stability: Balanced Case

- What are  $q_{\min}, q_{\max}$ ?

# $q$ -Stability: Balanced Case

- What are  $q_{\min}, q_{\max}$ ?
  - Courtade–Kumar Conjecture  $\iff q_{\min} = 1 \iff$  Li–Médard Conjecture

# $q$ -Stability: Balanced Case


- What are  $q_{\min}, q_{\max}$ ?
  - Courtade–Kumar Conjecture  $\iff q_{\min} = 1 \iff$  Li–Médard Conjecture
  - Mossel–O’Donnell Conjecture  $\iff q_{\max} \geq 9$

# $q$ -Stability: Balanced Case

- What are  $q_{\min}, q_{\max}$ ?
  - Courtade–Kumar Conjecture  $\iff q_{\min} = 1 \iff$  Li–Médard Conjecture
  - Mossel–O’Donnell Conjecture  $\iff q_{\max} \geq 9$
  - $q_{\max} < 10$  [Mossel and O’Donnell, 2005]

# $q$ -Stability: Balanced Case

- What are  $q_{\min}, q_{\max}$ ?
  - Courtade–Kumar Conjecture  $\iff q_{\min} = 1 \iff$  Li–Médard Conjecture
  - Mossel–O’Donnell Conjecture  $\iff q_{\max} \geq 9$
  - $q_{\max} < 10$  [Mossel and O’Donnell, 2005]



# Progress on Courtade–Kumar Conjecture (Li–Médard Conjecture)

Related Works	Upper Bounds on $\max_{\text{Boolean } f : \mathbb{E}f = 1/2} I(f(\mathbf{X}) ; \mathbf{Y})$	Tools
Courtade–Kumar Conjecture	$1 - H_2\left(\frac{1-\rho}{2}\right)$	
[Witsenhausen and Wyner, 1975]	$\rho^2$	Mrs. Gerber's lemma (or HC)
[Ordentlich et al., 2016]	$\frac{\log_2 e}{2} \rho^2 + 9\left(1 - \frac{\log_2 e}{2}\right) \rho^4$ for $0 \leq \rho \leq \frac{1}{\sqrt{3}}$ (asymptotically sharp as $\rho \rightarrow 0$ )	Fourier analysis + HC
[Samorodnitsky, 2016]	Sharp bound for $\rho \in [0, \rho_0]$ with some $0 < \rho_0 < 1$	Fourier analysis + Random restrictions + ...
[Yu, 2021b]	Sharp bound for $\rho \in [0, \rho_1]$ with $\rho_1 \approx 0.46$ (explicitly given)	Fourier analysis + KKT conditions
[Pichler et al., 2018]	A weaker version: $\max_{\text{Boolean } f, g} I(f(\mathbf{X}) ; g(\mathbf{Y})) =$ $1 - H_2\left(\frac{1-\rho}{2}\right)$	Fourier analysis + Partition technique

# Progress on Mossel–O’Donnell Conjecture

Related Works	Dictators are optimal for ...	Tools
<b>Mossel–O’Donnell Conjecture</b>	$2 < q \leq 9$ <b>(both symmetric and asymmetric max <math>q</math>-stability)</b>	
[Mossel and O’Donnell, 2005]	$2 < q \leq 3$ (symmetric)	Reducing $q = 3$ to $q = 2$
[Witsenhausen, 1975]	$q = 2$ (asymmetric)	Maximal correlation
[Yu, 2021b]	$2 < q \leq 5$ (symmetric); $2 < q \leq 3$ (asymmetric)	Fourier analysis + KKT conditions
[Mossel and O’Donnell, 2005, Li and Médard, 2021]	$\rho \rightarrow 0$ or $1$ (symmetric and asymmetric)	Fourier analysis

# Outline

## 1 Gács–Körner–Witsenhausen's CI

## 2 NICD with 2 Users

- Formulation of NICD
- Achievability Part: Subcubes, Balls, and Spheres
- Converse Part: CL Regime with Large  $a, b$
- Converse Part: MD Regime and CL Regime with Small  $a, b$
- Converse Part: LD Regime
- Connection to Hypercontractivity
- Extension to Gaussian Sources

## 3 NICD with Multiple Users and $q$ -Stability

- Formulation
- Balanced Case
- MD and LD Regimes

# MD and LD Regimes

- Recall: For  $a = \frac{M}{2^n}$  (with integer  $M$ ), (asymmetric) **max  $q$ -stability** with  $q \in [1, \infty)$  is defined as

$$\begin{aligned}\bar{\Gamma}_q(a) &:= \max_{A: P_X^n(A)=a} \mathbb{E}_{\mathbf{Y}} \left[ P_{X|Y}^n(A|\mathbf{Y})^q \right] \\ &= \left( \max_{A: P_X^n(A)=a} \left\| P_{X|Y}^n(A|\cdot) \right\|_q \right)^q\end{aligned}$$

# MD and LD Regimes

- Recall: For  $a = \frac{M}{2^n}$  (with integer  $M$ ), (asymmetric) **max  $q$ -stability** with  $q \in [1, \infty)$  is defined as

$$\begin{aligned}\bar{\Gamma}_q(a) &:= \max_{A: P_X^n(A)=a} \mathbb{E}_{\mathbf{Y}} \left[ P_{X|Y}^n(A|\mathbf{Y})^q \right] \\ &= \left( \max_{A: P_X^n(A)=a} \left\| P_{X|Y}^n(A|\cdot) \right\|_q \right)^q\end{aligned}$$

- Counterpart: **min  $q$ -stability** with  $q \in (-\infty, 1) \setminus \{0\}$ ,

$$\underline{\Gamma}_q(a) := \left( \min_{A: P_X^n(A)=a} \left\| P_{X|Y}^n(A|\cdot) \right\|_q \right)^q$$

# MD and LD Regimes

- Recall: For  $a = \frac{M}{2^n}$  (with integer  $M$ ), (asymmetric) **max  $q$ -stability** with  $q \in [1, \infty)$  is defined as

$$\begin{aligned}\bar{\Gamma}_q(a) &:= \max_{A: P_X^n(A)=a} \mathbb{E}_{\mathbf{Y}} \left[ P_{X|Y}^n(A|\mathbf{Y})^q \right] \\ &= \left( \max_{A: P_X^n(A)=a} \left\| P_{X|Y}^n(A|\cdot) \right\|_q \right)^q\end{aligned}$$

- Counterpart: **min  $q$ -stability** with  $q \in (-\infty, 1) \setminus \{0\}$ ,

$$\underline{\Gamma}_q(a) := \left( \min_{A: P_X^n(A)=a} \left\| P_{X|Y}^n(A|\cdot) \right\|_q \right)^q$$

- Limit cases as  $n \rightarrow \infty$ :

- LD regime:**  $a = 2^{-n\alpha}$  is exponentially small

# MD and LD Regimes

- Recall: For  $a = \frac{M}{2^n}$  (with integer  $M$ ), (asymmetric) **max  $q$ -stability** with  $q \in [1, \infty)$  is defined as

$$\begin{aligned}\bar{\Gamma}_q(a) &:= \max_{A: P_X^n(A)=a} \mathbb{E}_{\mathbf{Y}} \left[ P_{X|Y}^n(A|\mathbf{Y})^q \right] \\ &= \left( \max_{A: P_X^n(A)=a} \left\| P_{X|Y}^n(A|\cdot) \right\|_q \right)^q\end{aligned}$$

- Counterpart: **min  $q$ -stability** with  $q \in (-\infty, 1) \setminus \{0\}$ ,

$$\underline{\Gamma}_q(a) := \left( \min_{A: P_X^n(A)=a} \left\| P_{X|Y}^n(A|\cdot) \right\|_q \right)^q$$

- Limit cases as  $n \rightarrow \infty$ :

- LD regime:**  $a = 2^{-n\alpha}$  is exponentially small
- MD regime:**  $a = 2^{-\theta_n \alpha}$  is “subexponentially” small with  $\theta_n \rightarrow \infty$ ,  $\frac{\theta_n}{n} \rightarrow 0$

## Theorem ( $q$ -Stability Theorem)

*Hamming balls/spheres are optimal in attaining max  $q$ -stability for  $q \in [1, \infty)$  and min  $q$ -stability for  $q \in (-\infty, 1) \setminus \{0\}$  in the MD regime.*

## Theorem ( $q$ -Stability Theorem)

*Hamming balls/spheres are optimal in attaining max  $q$ -stability for  $q \in [1, \infty)$  and min  $q$ -stability for  $q \in (-\infty, 1) \setminus \{0\}$  in the MD regime.*

Proof:

- Single-Function Version of Forward HC: For any nonnegative  $f$ ,

$$\|T_\rho f(\mathbf{Y})\|_{q'} \leq \|f(\mathbf{X})\|_p$$

where  $q' := \frac{q}{q-1}$  denote the Hölder conjugate of  $q$ , and  $p, q \geq 1$  s.t.  $(p-1)(q-1) \geq \rho^2$

## Theorem ( $q$ -Stability Theorem)

*Hamming balls/spheres are optimal in attaining max  $q$ -stability for  $q \in [1, \infty)$  and min  $q$ -stability for  $q \in (-\infty, 1) \setminus \{0\}$  in the MD regime.*

Proof:

- Single-Function Version of Forward HC: For any nonnegative  $f$ ,

$$\|T_\rho f(\mathbf{Y})\|_{q'} \leq \|f(\mathbf{X})\|_p$$

where  $q' := \frac{q}{q-1}$  denote the Hölder conjugate of  $q$ , and  $p, q \geq 1$  s.t.  $(p-1)(q-1) \geq \rho^2$

- Single-Function Version of Reverse HC: For any nonnegative  $f$ ,

$$\|T_\rho f(\mathbf{Y})\|_{q'} \geq \|f(\mathbf{X})\|_p$$

where  $p, q \leq 1$  s.t.  $(p-1)(q-1) \geq \rho^2$

## Theorem ( $q$ -Stability Theorem)

*Hamming balls/spheres are optimal in attaining max  $q$ -stability for  $q \in [1, \infty)$  and min  $q$ -stability for  $q \in (-\infty, 1) \setminus \{0\}$  in the MD regime.*

Proof:

- Single-Function Version of Forward HC: For any nonnegative  $f$ ,

$$\|T_\rho f(\mathbf{Y})\|_{q'} \leq \|f(\mathbf{X})\|_p$$

where  $q' := \frac{q}{q-1}$  denote the Hölder conjugate of  $q$ , and  $p, q \geq 1$  s.t.  $(p-1)(q-1) \geq \rho^2$

- Single-Function Version of Reverse HC: For any nonnegative  $f$ ,

$$\|T_\rho f(\mathbf{Y})\|_{q'} \geq \|f(\mathbf{X})\|_p$$

where  $p, q \leq 1$  s.t.  $(p-1)(q-1) \geq \rho^2$

- Substituting  $f \leftarrow 1_A$  to the Single-Function Version of HC  $\implies q$ -Stability Theorem

# LD Regime

- Similarly, by Single-Function Version of Strong HC [Yu, 2021c], we have:

- Similarly, by Single-Function Version of Strong HC [Yu, 2021c], we have:

## Theorem (Strong $q$ -Stability Theorem [Yu, 2021c, Yu, 2021a])

*Hamming balls/spheres* are optimal in attaining max  $q$ -stability for  $q \in [1, \infty)$  and min  $q$ -stability for  $q \in (-\infty, 1) \setminus \{0\}$  in the LD regime.

# LD Regime

- Similarly, by Single-Function Version of Strong HC [Yu, 2021c], we have:

## Theorem (Strong $q$ -Stability Theorem [Yu, 2021c, Yu, 2021a])

*Hamming balls/spheres* are optimal in attaining max  $q$ -stability for  $q \in [1, \infty)$  and min  $q$ -stability for  $q \in (-\infty, 1) \setminus \{0\}$  in the LD regime.

Bounds	Moderate Deviation	Large Deviation
$q$ -Stability Theorem	subexp. vanishing $a$	exp. vanishing $a$
Strong $q$ -Stability Theorem [Yu, 2021c]	Sharp (Balls/Spheres Optimal)	Sharp (Balls/Spheres Optimal)

# Summary of Tools used in NICD and $q$ -Stability

Methods		Central Limit		Moderate Deviation	Large Deviation
		fixed and large $a$ (and $b$ )	fixed but small $a$ (and $b$ )	subexp. vanishing $a$ (and $b$ )	exp. vanishing $a$ (and $b$ )
Information-Theoretic Methods	Maximal Correlation	<b>Sharp</b> for 2-user NICD with $a = b = 1/2$	Almost sharp	<b>Sharp</b>	
	HC/SSE (stronger than MC)				
	Strong HC/SSE (stronger than HC/SSE)				<b>Sharp</b>
Fourier Analysis	Combined with Optimization Theory (LP or KKT)	<b>Sharp</b> for 2-user NICD with $a = b = 1/2, 1/4$ ; <b>Sharp</b> for $q$ -Stability with $a = 1/2$ and certain $(q, \rho)$			

# References I



Ahlswede, R. and Gács, P. (1976).

Spreading of sets in product spaces and hypercontraction of the markov operator.

*The Annals of Probability*, pages 925–939.



Barnes, L. P. and Özgür, A. (2020).

The Courtade-Kumar most informative Boolean function conjecture and a symmetrized Li-Médard conjecture are equivalent.

In *2020 IEEE International Symposium on Information Theory (ISIT)*, pages 2205–2209. IEEE.



Beigi, S. and Nair, C. (2016).

Equivalent characterization of reverse Brascamp-Lieb-type inequalities using information measures.

In *2016 IEEE International Symposium on Information Theory (ISIT)*, pages 1038–1042. IEEE.



Bonami, A. (1968).

Ensembles  $\Lambda(p)$  dans le dual de  $D^\infty$ .

In *Annales de l'institut Fourier*, volume 18, pages 193–204.



Borell, C. (1982).

Positivity improving operators and hypercontractivity.

*Mathematische Zeitschrift*, 180(3):225–234.



Borell, C. (1985).

Geometric bounds on the Ornstein-Uhlenbeck velocity process.

*Probability Theory and Related Fields*, 70(1):1–13.

# References II



Carlen, E. A. and Cordero-Erausquin, D. (2009).  
Subadditivity of the entropy and its relation to Brascamp–Lieb type inequalities.  
*Geometric and Functional Analysis*, 19(2):373–405.



Csiszár, I. and Narayan, P. (2000).  
Common randomness and secret key generation with a helper.  
*IEEE Trans. Inf. Theory*, 46(2):344–366.



Fu, F.-W., Wei, V. K., and Yeung, R. W. (2001).  
On the minimum average distance of binary codes: Linear programming approach.  
*Discrete Applied Mathematics*, 111(3):263–281.



Gács, P. and Körner, J. (1973).  
Common information is far less than mutual information.  
*Problems of Control and Information Theory*, 2(2):149–162.



Gross, L. (1975).  
Logarithmic sobolev inequalities.  
*American Journal of Mathematics*, 97(4):1061–1083.



Kahn, J., Kalai, G., and Linial, N. (1988).  
The influence of variables on Boolean functions.  
In *29th Annual Symposium on Foundations of Computer Science*, pages 68–80. IEEE.

# References III

-  Kamath, S. and Anantharam, V. (2016).  
On non-interactive simulation of joint distributions.  
*IEEE Trans. Inf. Theory*, 62(6):3419–3435.
-  Kirshner, N. and Samorodnitsky, A. (2021).  
A moment ratio bound for polynomials and some extremal properties of Krawchouk polynomials and Hamming spheres.  
*IEEE Trans. Inf. Theory*, 67(6):3509–3541.
-  Li, J. and Médard, M. (2021).  
Boolean functions: noise stability, non-interactive correlation distillation, and mutual information.  
*IEEE Trans. Inf. Theory*, 67(2):778–789.
-  Mossel, E. and Neeman, J. (2015).  
Robust optimality of Gaussian noise stability.  
*Journal of the European Mathematical Society*, 17(2):433–482.
-  Mossel, E. and O'Donnell, R. (2005).  
Coin flipping from a cosmic source: On error correction of truly random bits.  
*Random Structures & Algorithms*, 26(4):418–436.
-  Mossel, E., O'Donnell, R., Regev, O., Steif, J. E., and Sudakov, B. (2006).  
Non-interactive correlation distillation, inhomogeneous Markov chains, and the reverse Bonami-Beckner inequality.  
*Israel Journal of Mathematics*, 154(1):299–336.

# References IV



O'Donnell, R. (2014).  
*Analysis of Boolean Functions*.  
Cambridge University Press.



Ordentlich, O., Polyanskiy, Y., and Shayevitz, O. (2020).  
A note on the probability of rectangles for correlated binary strings.  
*IEEE Trans. Inf. Theory*.



Ordentlich, O., Shayevitz, O., and Weinstein, O. (2016).  
An improved upper bound for the most informative Boolean function conjecture.  
In *2016 IEEE International Symposium on Information Theory (ISIT)*, pages 500–504. IEEE.



Pichler, G., Piantanida, P., and Matz, G. (2018).  
Dictator functions maximize mutual information.  
*The Annals of Applied Probability*, 28(5):3094–3101.



Samorodnitsky, A. (2016).  
On the entropy of a noisy function.  
*IEEE Trans. Inf. Theory*, 62(10):5446–5464.



Witsenhausen, H. and Wyner, A. (1975).  
A conditional entropy bound for a pair of discrete random variables.  
*IEEE Trans. Inf. Theory*, 21(5):493–501.

# References V



Witsenhausen, H. S. (1975).

On sequences of pairs of dependent random variables.

*SIAM Journal on Applied Mathematics*, 28(1):100–113.



Yu, L. (2021a).

The convexity and concavity of envelopes of the minimum-relative-entropy region for the DSBS.

*arXiv preprint arXiv:2106.03654*.



Yu, L. (2021b).

On the  $\Phi$ -stability and related conjectures.

*arXiv preprint arXiv:2104.08740*.



Yu, L. (2021c).

Strong Brascamp–Lieb inequalities.

*arXiv preprint arXiv:2102.06935*.



Yu, L., Anantharam, V., and Chen, J. (2021).

Graphs of joint types, noninteractive simulation, and stronger hypercontractivity.

*arXiv preprint arXiv:2102.00668*.



Yu, L. and Tan, V. Y. F. (2019).

An improved linear programming bound on the average distance of a binary code.

*arXiv preprint arXiv:1910.09416*.

# References VI

- 
- Yu, L. and Tan, V. Y. F. (2021).  
On non-interactive simulation of binary random variables.  
*IEEE Trans. Inf. Theory*, 67(4):2528–2538.

