

Necessary and Sufficient Conditions for High-Dimensional Salient Feature Subset Recovery

Vincent Tan, Matt Johnson, Alan S. Willsky

Stochastic Systems Group,
Laboratory for Information and Decision Systems,
Massachusetts Institute of Technology


ISIT (Jun 14, 2010)

Motivation: A Real-Life Example

- The Manchester Asthma and Allergy Study (MAAS) is a birth-cohort study of more than $n \approx 1000$ children.

Motivation: A Real-Life Example

- The Manchester Asthma and Allergy Study (MAAS) is a birth-cohort study of more than $n \approx 1000$ children.



www.maas.org.uk

Motivation: A Real-Life Example

- The Manchester Asthma and Allergy Study (MAAS) is a birth-cohort study of more than $n \approx 1000$ children.




www.maas.org.uk

- Number of variables $d \approx 10^6$.

Motivation: A Real-Life Example

- The Manchester Asthma and Allergy Study (MAAS) is a birth-cohort study of more than $n \approx 1000$ children.



www.maas.org.uk

- Number of variables $d \approx 10^6$.
- But only $k \approx 30$ are **salient** for assessing susceptibility to asthma.

Motivation: A Real-Life Example

- The Manchester Asthma and Allergy Study (MAAS) is a birth-cohort study of more than $n \approx 1000$ children.



www.maas.org.uk

- Number of variables $d \approx 10^6$.
- But only $k \approx 30$ are **salient** for assessing susceptibility to asthma.
- Identification of these salient features is important.

Some Natural Questions

Natural questions:

Some Natural Questions

Natural questions:

- What is the meaning of **saliency**?
- How to define it **information-theoretically**?

Some Natural Questions

Natural questions:

- What is the meaning of **saliency**?
- How to define it **information-theoretically**?
- How many samples n are required to **identify** the k salient features out of the d features?

Num features $d \gg$ Num salient features k

- What are the **fundamental limits** for recovery of the **salient set**?

Some Natural Questions

Natural questions:

- What is the meaning of **saliency**?
- How to define it **information-theoretically**?
- How many samples n are required to **identify** the k salient features out of the d features?

Num features $d \gg$ Num salient features k

- What are the **fundamental limits** for recovery of the **salient set**?
- Are there any **efficient algorithms** for special classes of features?

Main Contributions

- We define the **salient set** by appealing to the **error exponents** in hypothesis testing.

Main Contributions

- We define the **salient set** by appealing to the **error exponents** in **hypothesis testing**.
- **Sufficiency**: We show that if for all n sufficiently large,

$$n > \max \left\{ C_1 \cdot k \cdot \log \left(\frac{d-k}{k} \right), \exp(C_2 \cdot k) \right\},$$

then the error probability is arbitrarily small.

Main Contributions

- We define the **salient set** by appealing to the **error exponents** in **hypothesis testing**.
- **Sufficiency**: We show that if for all n sufficiently large,

$$n > \max \left\{ C_1 \cdot k \cdot \log \left(\frac{d-k}{k} \right), \exp(C_2 \cdot k) \right\},$$

then the error probability is arbitrarily small.

- **Necessity**: Under certain conditions, for $\lambda \in (0, 1)$, if

$$n < \lambda \cdot C_3 \cdot \log \left(\frac{d}{k} \right)$$

then the error probability $\geq 1 - \lambda$.

System Model

- Let the alphabet for each variable be \mathcal{X} , a finite set.
- **High-dimensional** setting where d and k grow with n .
- **Two sequences** of unknown d -dimensional distributions

$$P^{(d)}, Q^{(d)} \in \mathcal{P}(\mathcal{X}^d), \quad d \in \mathbb{N}.$$

System Model

- Let the alphabet for each variable be \mathcal{X} , a finite set.
- **High-dimensional** setting where d and k grow with n .
- **Two sequences** of unknown d -dimensional distributions

$$P^{(d)}, Q^{(d)} \in \mathcal{P}(\mathcal{X}^d), \quad d \in \mathbb{N}.$$

- IID samples $(\mathbf{x}^n, \mathbf{y}^n) := (\{\mathbf{x}^{(l)}\}_{l=1}^n, \{\mathbf{y}^{(l)}\}_{l=1}^n)$ drawn from $P^{(d)} \times Q^{(d)}$.
- Each pair of samples $\mathbf{x}^{(l)}, \mathbf{y}^{(l)} \in \mathcal{X}^d$.

Definition of Saliency

Motivated by **asymptotics** of binary hypothesis testing

$$H_0 : \mathbf{z}^n \sim P^{(d)}, \quad H_1 : \mathbf{z}^n \sim Q^{(d)}$$

Definition of Saliency

Motivated by **asymptotics** of binary hypothesis testing

$$H_0 : \mathbf{z}^n \sim P^{(d)}, \quad H_1 : \mathbf{z}^n \sim Q^{(d)}$$

Chernoff-Stein Lemma: If $\Pr(\hat{H}_1|H_0) < \alpha$, then

$$\Pr(\hat{H}_0|H_1) \doteq \exp(-nD(P^{(d)} \parallel Q^{(d)}))$$

Chernoff Information:

$$\Pr(\text{err}) = \pi_0 \Pr(\hat{H}_1|H_0) + \pi_1 \Pr(\hat{H}_0|H_1) \doteq \exp(-nC(P^{(d)}, Q^{(d)}))$$

where

$$C(P, Q) := - \min_{t \in [0,1]} \log \sum_{\mathbf{z}} P(\mathbf{z})^t Q(\mathbf{z})^{1-t}$$

Definition of Saliency

$P_A^{(d)}$ is the **marginal** of $P^{(d)}$ on the subset $A \subset \{1, \dots, d\}$.

Definition of Saliency

$P_A^{(d)}$ is the **marginal** of $P^{(d)}$ on the subset $A \subset \{1, \dots, d\}$.

- **Definition:** A set $S_d \subset \{1, \dots, d\}$ is **KL-divergence-salient** if

$$D(P^{(d)} \parallel Q^{(d)}) = D(P_{S_d}^{(d)} \parallel Q_{S_d}^{(d)}).$$

Definition of Saliency

$P_A^{(d)}$ is the **marginal** of $P^{(d)}$ on the subset $A \subset \{1, \dots, d\}$.

- **Definition:** A set $S_d \subset \{1, \dots, d\}$ is **KL-divergence-salient** if

$$D(P^{(d)} \parallel Q^{(d)}) = D(P_{S_d}^{(d)} \parallel Q_{S_d}^{(d)}).$$

- **Definition:** A set $S_d \subset \{1, \dots, d\}$ is **Chernoff Information-salient** if

$$C(P^{(d)}, Q^{(d)}) = C(P_{S_d}^{(d)}, Q_{S_d}^{(d)}).$$

Characterization of Saliency

Lemma

The following are equivalent:

- S_d is KL-divergence-salient.
- S_d is Chernoff Information-salient.

Characterization of Saliency

Lemma

The following are equivalent:

- S_d is KL-divergence-salient.
- S_d is Chernoff Information-salient.
- Conditionals are identical

$$P^{(d)} = P_{S_d}^{(d)} \cdot W_{S_d^c|S_d} \quad Q^{(d)} = Q_{S_d}^{(d)} \cdot W_{S_d^c|S_d}.$$

Characterization of Saliency

Lemma

The following are equivalent:

- S_d is KL-divergence-salient.
- S_d is Chernoff Information-salient.
- Conditionals are identical

$$P^{(d)} = P_{S_d}^{(d)} \cdot W_{S_d^c|S_d} \quad Q^{(d)} = Q_{S_d}^{(d)} \cdot W_{S_d^c|S_d}.$$

What are the **scaling laws** on (n, d, k) so that the error probability can be made arbitrarily small?

Definition of Achievability

- A **decoder** is a set-valued function that maps samples to subsets of size k , i.e.,

$$\psi_n : (\mathcal{X}^d)^n \times (\mathcal{X}^d)^n \rightarrow \binom{\{1, \dots, d\}}{k}.$$

- The decoder is **given** the true value of k , the number of salient features.
- We are working on relaxing this assumption.

Definition of Achievability

- A **decoder** is a set-valued function that maps samples to subsets of size k , i.e.,

$$\psi_n : (\mathcal{X}^d)^n \times (\mathcal{X}^d)^n \rightarrow \binom{\{1, \dots, d\}}{k}.$$

- The decoder is **given** the true value of k , the number of salient features.
- We are working on relaxing this assumption.

Definition: The tuple of model parameters (n, d, k) is **achievable** for $\{P^{(d)}, Q^{(d)}\}_{d \in \mathbb{N}}$ if **there exists a sequence of decoders** $\{\psi_n\}$ such that

$$q_n(\psi_n) := \Pr(\psi_n(\mathbf{x}^n, \mathbf{y}^n) \neq S_d) < \epsilon, \quad \forall n > N_\epsilon.$$

Three Assumptions on Distributions $P^{(d)}, Q^{(d)}$

- **Saliency**: For every $P^{(d)}, Q^{(d)}$ there exists a salient set S_d of known size k , i.e.,

$$D(P^{(d)} || Q^{(d)}) = D(P_{S_d}^{(d)} || Q_{S_d}^{(d)}).$$

Three Assumptions on Distributions $P^{(d)}, Q^{(d)}$

- **Saliency**: For every $P^{(d)}, Q^{(d)}$ there exists a salient set S_d of known size k , i.e.,

$$D(P^{(d)} || Q^{(d)}) = D(P_{S_d}^{(d)} || Q_{S_d}^{(d)}).$$

- **η -Distinguishability**: There exists a constant $\eta > 0$ such that

$$D(P_{S_d}^{(d)} || Q_{S_d}^{(d)}) - D(P_{T_d}^{(d)} || Q_{T_d}^{(d)}) \geq \eta$$

for all $T_d \neq S_d$ such that $|T_d| = k$.

Three Assumptions on Distributions $P^{(d)}, Q^{(d)}$

- **Saliency**: For every $P^{(d)}, Q^{(d)}$ there exists a salient set S_d of known size k , i.e.,

$$D(P^{(d)} || Q^{(d)}) = D(P_{S_d}^{(d)} || Q_{S_d}^{(d)}).$$

- **η -Distinguishability**: There exists a constant $\eta > 0$ such that

$$D(P_{S_d}^{(d)} || Q_{S_d}^{(d)}) - D(P_{T_d}^{(d)} || Q_{T_d}^{(d)}) \geq \eta$$

for all $T_d \neq S_d$ such that $|T_d| = k$.

- **L -Boundedness**: There exists a constant $L \in (0, \infty)$ such that

$$\log \left[\frac{P_{S_d}^{(d)}(\mathbf{z}_{S_d})}{Q_{S_d}^{(d)}(\mathbf{z}_{S_d})} \right] \in [-L, L]$$

for all states $\mathbf{z}_{S_d} \in \mathcal{X}^k$.

Achievability Result

Theorem

If there exists an $\delta > 0$ such that for some $B > 0$

$$n > \max \left\{ \frac{k}{B} \log \left(\frac{d-k}{k} \right), \exp \left(\frac{2k \log |\mathcal{X}|}{1-\delta} \right) \right\},$$

then there exists a sequence of decoders ψ_n^ that satisfies*

$$q_n(\psi_n^*) = O(\exp(-nE)),$$

for some exponent $E > 0$.

Proof Idea and a Corollary

- Use the **exhaustive search decoder**. Search for the size- k set with the **largest empirical KL-divergence**.
- Large deviation bounds, e.g., Sanov's theorem.

Proof Idea and a Corollary

- Use the **exhaustive search decoder**. Search for the size- k set with the **largest empirical KL-divergence**.
- Large deviation bounds, e.g., Sanov's theorem.
- Positivity of **error exponents** under the specified conditions.
- Similar to main result in [Ng, UAI 1998] but we focus on exact subset recovery and not generalization error.

Proof Idea and a Corollary

- Use the **exhaustive search decoder**. Search for the size- k set with the **largest empirical KL-divergence**.
- Large deviation bounds, e.g., Sanov's theorem.
- Positivity of **error exponents** under the specified conditions.
- Similar to main result in [Ng, UAI 1998] but we focus on exact subset recovery and not generalization error.

Corollary

Let $k = k_0$ be a constant and $R \in (0, B/k_0)$. Then if

$$n > \frac{\log d}{R}$$

Proof Idea and a Corollary

- Use the **exhaustive search decoder**. Search for the size- k set with the **largest empirical KL-divergence**.
- Large deviation bounds, e.g., Sanov's theorem.
- Positivity of **error exponents** under the specified conditions.
- Similar to main result in [Ng, UAI 1998] but we focus on exact subset recovery and not generalization error.

Corollary

Let $k = k_0$ be a constant and $R \in (0, B/k_0)$. Then if

$$n > \frac{\log d}{R} \quad \Rightarrow \quad q_n(\psi_n^*) = O(\exp(-nE)).$$

Converse Result

We assume that the salient set S_d is chosen **uniformly at random** over all subsets of size k .

Converse Result

We assume that the salient set S_d is chosen **uniformly at random** over all subsets of size k .

Theorem

If for some $\lambda \in (0, 1)$,

$$n < \frac{\lambda \cdot k \cdot \log\left(\frac{d}{k}\right)}{H(P^{(d)}) + H(Q^{(d)})}$$

then

$$q_n(\psi_n) \geq 1 - \lambda$$

for all decoders ψ_n .

Remarks

- Proof is a consequence of **Fano's inequality**.

Remarks

- Proof is a consequence of **Fano's inequality**.
- Bound **never satisfied** if variables in S_d^c are uniform and independent of S_d .

$$n < \underbrace{\frac{\lambda}{H(P^{(d)}) + H(Q^{(d)})}}_{O(d)} \cdot k \cdot \log \left(\frac{d}{k} \right)$$

- However, converse is interesting if distributions have additional structure on their **entropies**.

Remarks

- Proof is a consequence of **Fano's inequality**.
- Bound **never satisfied** if variables in S_d^c are uniform and independent of S_d .

$$n < \underbrace{\frac{\lambda}{H(P^{(d)}) + H(Q^{(d)})}}_{O(d)} \cdot k \cdot \log \left(\frac{d}{k} \right)$$

- However, converse is interesting if distributions have additional structure on their **entropies**.
- Assume most of the features are **processed** or **redundant**.
- Example: there could be two features, “BMI” and “isObese”. One is a **processed** version of the other.

Converse Result

Corollary

Assume that there exists a $M < \infty$ such that the conditional entropies satisfy

$$\max \left\{ H(P_{S_d^c|S_d}^{(d)}), H(Q_{S_d^c|S_d}^{(d)}) \right\} \leq M \cdot k.$$

Converse Result

Corollary

Assume that there exists a $M < \infty$ such that the conditional entropies satisfy

$$\max \left\{ H(P_{S_d^c|S_d}^{(d)}), H(Q_{S_d^c|S_d}^{(d)}) \right\} \leq M \cdot k.$$

If the number of samples satisfies

$$n < \frac{\lambda}{2(M + \log |\mathcal{X}|)} \cdot \log \left(\frac{d}{k} \right)$$

then


$$q_n(\psi_n) \geq 1 - \lambda.$$

Comparison to Achievability Result

- Assume $d = \exp(nR)$ and k is constant.


Comparison to Achievability Result

- Assume $d = \exp(nR)$ and k is constant.
- There is a rate R_{ach} so that if $R < R_{\text{ach}}$, then (n, d, k) is **achievable**.
- **Conversely**, there is another rate R_{conv} so that if $R > R_{\text{conv}}$, then recovery is not possible.




Comparison to Achievability Result

- Assume $d = \exp(nR)$ and k is constant.
- There is a rate R_{ach} so that if $R < R_{\text{ach}}$, then (n, d, k) is **achievable**.
- Conversely**, there is another rate R_{conv} so that if $R > R_{\text{conv}}$, then recovery is not possible.



Comparison to Achievability Result

- Assume $d = \exp(nR)$ and k is constant.
- There is a rate R_{ach} so that if $R < R_{\text{ach}}$, then (n, d, k) is **achievable**.
- Conversely**, there is another rate R_{conv} so that if $R > R_{\text{conv}}$, then recovery is not possible.



Conclusions

- Provided an information-theoretic definition of saliency motivated by **error exponents** in **hypothesis testing**.

Conclusions

- Provided an information-theoretic definition of saliency motivated by **error exponents** in **hypothesis testing**.
- Provided **necessary** and **sufficient** conditions for salient set recovery.
- Number of samples n can be **much smaller** than d , total number of variables.

Conclusions

- Provided an information-theoretic definition of saliency motivated by **error exponents** in **hypothesis testing**.
- Provided **necessary** and **sufficient** conditions for salient set recovery.
- Number of samples n can be **much smaller** than d , total number of variables.
- In the paper, we provide a **computationally efficient** and **consistent** algorithm to search for S_d when $P^{(d)}$ and $Q^{(d)}$ are Markov on trees.