

1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 4



Chapter 5 · Original slide 5



1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 6

Examples

This means we can **substitute** the values of a coordinate into our equation whenever we know the point lies on the line.

The point $(5, a)$ lies on the line with equation $y = 3x + 2$. Determine the value of a.

Find the coordinate of the point where the line $2x + y = 5$ cuts the x-axis.

Chapter 5 · Original slide 7

Test Your Understanding

Determine where the line $x + 2y = 3$ crosses the:

a) y-axis:

b) x-axis:

What mistakes do you think it's easy to make?


-
- **S**

1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 8

Recap of gradient

The steepness of a line is known as the **gradient**.
It tells us what y changes by as x increases by 1.



So if the y value increased by 6 as the x value increased by 2, what is y increasing by for each unit increase of x?
How would that give us a suitable formula for the gradient m?

$m =$ ☐

Chapter 5 · Original slide 9

Examples

Find the gradient of the line that goes through the points:

1 $(1, 4)$ $(3, 10)$ $m =$ ☐

2 $(5, 7)$ $(8, 1)$ $m =$ ☐

3 $(2, 2)$ $(-1, 10)$ $m =$ ☐

4 Show that the points
$\mathrm{A}\,(3, 4), B\,(5, 5), C\,(11, 8)$ all lie on a straight

1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 10

Further Example

The line joining $(2, -5)$ to $(4, a)$ has gradient -1. Work out the value of a.

$\frac{a}{}$

$\frac{a}{}$

a

Chapter 5 · Original slide 11

$y = mx + c$

One form we can put a straight line equation in is:

$$y = mx + c$$

Gradient

y-intercept

Why does it work?

1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 12

$y = mx + c$

Determine the gradient and y-intercept of the line with equation $4x - 3y + 5 = 0$

$3y$

$y =$

Make y the subject so we have the form $y = mx + c$
Put y on the side it's positive.

Chapter 5 · Original slide 13

$ax + by + c = 0$

At GCSE, $\mathbf{y=mx+c}$ was the main form you would express a straight line equation, sometimes known as the '**slope-intercept form**'.
But another common form is $\mathbf{ax+by+c=0}$, where a, b, c are integers.
This is known as the '**standard**' form.

Express $y = \frac{1}{3}x - \frac{2}{3}$ in the form $ax + by + c = 0$, where a, b, c are integers.

We'll see on the next slide WHY we might want to put an equation in this form over $y = mx + c$...

1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 14

Just for your interest…

Why might we want to put a straight line equation in the form $ax + by + c = 0$?

$y = mx + c$
"Slope-Intercept Form"

$ax + by + c = 0$
"Standard Form"

Coverage

$y = mx + c$ doesn't allow you to represent vertical lines. Standard form allows us to do this by just making b zero.

$x + 4 = 0$

Symmetry

In general, the '**linear combination**' of two variables x and y is $ax + by$, i.e. "some amount of x and some amount of y". There is a greater elegance and symmetry to this form over $y = mx + c$ because x and y appear similarly within the expression.

Usefulness

This more 'elegant' form also means it ties in with vectors and matrices. In FM, you will learn about the '**dot product**' of two vectors:

$$\begin{pmatrix} a \\ b \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = ax + by$$

thus since $ax + by + c = 0$, we can represent a straight line using:


$$\begin{pmatrix} a \\ b \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} + c = 0 \qquad (1)$$

We can extend to 3D points to get the equation of a **plane**:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} + d = 0 \qquad (2)$$

Conveniently, in equation (1), the vector $\begin{pmatrix} a \\ b \end{pmatrix}$ is **perpendicular to the line**. And in equation (2), the vector $\begin{pmatrix} a \\ b \\ c \end{pmatrix}$ is perpendicular to the plane. Nice!

$2x + y = 4$ ➡ $\begin{pmatrix} 2 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = 4$ ➡



Chapter 5 · Original slide 15

Test Your Understanding

Express $y = \frac{2}{5}x + \frac{3}{5}$ in the form $ax + by + c = 0$, where a, b, c are integers.

$5y$
$2x$

1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 16

Exercise 5A – Pure Y1 Page 91

9 Show that the points $A(2, 3)$, $B(4, 4)$ and $C(10, 7)$ can be joined by a straight line.

10 Show that the points $A(-2a, 5a)$, $B(0, 4a)$ and points $C(6a, a)$ are collinear. $(3\ \mathrm{ma}\ldots$

Chapter 5 · Original slide 17

Exercise 5B – Pure Y1 Page 92-93

8 A line l cuts the x-axis at $(5, 0)$ and the y-axis at $(0, 2)$.

a Find the gradient of the line.

b Find an equation of the line in the form $ax + by + c = 0$.

9 Show that the line with equation $ax + by + c = 0$ has gradient $-\frac{a}{b}$ and cuts the y-axis at $-\frac{c}{b}$.

10 The line l with gradient 3 and y-intercept $(0, 5)$ has the equation $ax - 2y + c = 0$.


Find the values of a and c. (2 marks)

1.04 Graph Sketching: Linear Graphs

Chapter 5 · Original slide 18

Exercise 5B – Pure Y1 Page 92-93

11 The straight line l passes through $(0, 6)$ and has gradient -2. It intersects the line with equation $5x - 8y - 15 = 0$ at point P. Find the coordinates of P.



12 The straight line l_1 with equation $y = 3x - 7$ intersects the straight line l_2 with equation $ax + 4y - 17 = 0$ at the point $P(-3, b)$.

a Find the value of b.

b Find the value of a.