
1.14 Polynomials

Chapter 7 · Original slide 6

Terminology

6


$$11 \div 4 = 2rem3$$



Chapter 7 · Original slide 7

Normal Long Division

7



1. We found how many whole number of times (i.e. the quotient) the divisor went into the dividend.

2. We multiplied the quotient by the dividend.

3. …in order to find the remainder.

4. Find we ‘brought down’ the next number.

1.14 Polynomials

Chapter 7 · Original slide 8

Example

8

How many times does x go into $6x^3$?

$$6x^2 - 2x + 3$$

Now repeat! How many times does x go into $-2x^2$?

$$x + 5 \;\overline{)\; 6x^3 + 28x^2 - 7x + 15}$$

Multiply $6x^2$ by $(x + 5)$. The first term should match with above.

$$6x^3 + 30x^2$$

Subtract and carry down next term.

$$-2x^2 - 7x$$

$$-2x^2 - 10x$$

$$3x + 15$$

$$3x + 15$$

$$0$$

This is the remainder.

Tip:
You can check your solution by expanding:
$(x + 5)(6x^2 - 2x + 3)$
But if you know you should get no remainder, ending with 0 at the bottom is a good sign!

Tip:
Be **very** careful subtracting negatives.
$-7x - (-10x) = 3x$

Chapter 7 · Original slide 9

Further Example

9

Find the remainder when $3x^3 - 2x + 4$ is divided by $x - 1$.

$$x - 1 \;\overline{)\; 3x^3 + 0x^2 - 2x + 4}$$

Note: Ensure that any missing terms in the polynomial are filled in, so that the powers decrease by 1 each time.

1.14 Polynomials

Chapter 7 · Original slide 10

Test Your Understanding

10

Find the remainder when $2x^3 - 5x^2 - 16x + 10$ is divided by $x - 4$.

Chapter 7 · Original slide 11

Further Test Your Understanding

11

Divide $8x^3 - 1$ by $2x - 1$.

1.14 Polynomials

Chapter 7 · Original slide 12

Exercise 7B – Y1 Pages 142

12

15 $\mathrm{f}(x) = 10x^3 + 43x^2 - 2x - 10$.
Find the remainder when $\mathrm{f}(x)$ is divided by $(5x + 4)$.

16 $\mathrm{f}(x) = 3x^3 - 14x^2 - 47x - 14$.
a Find the remainder when $\mathrm{f}(x)$ is divided by $(x - 3)$.
b Given that $(x + 2)$ is a factor of $\mathrm{f}(x)$, factorise $\mathrm{f}(x)$ completely.

Chapter 7 · Original slide 13

Exercise 7B – Y1 Pages 142

13

17 a Find the remainder when $x^3 + 6x^2 + 5x - 12$ is divided by
i $x - 2$,
ii $x + 3$.
b Hence, or otherwise, find all the solutions to the equation $x^3 + 6x^2 + 5x - 12 = 0$.

1.14 Polynomials

Chapter 7 · Original slide 14

Exercise 7B – Y1 Pages 142

14

18 $\mathrm{f}(x) = 2x^3 + 3x^2 - 8x + 3$.
a Show that $\mathrm{f}(x) = (2x - 1)(ax^2 + bx + c)$ where a, b and c are constants to be found.
b Hence factorise $\mathrm{f}(x)$ completely.
c Write down all the real roots of the equation $\mathrm{f}(x) = 0$.

Chapter 7 · Original slide 15

Exercise 7B – Y1 Pages 142

15

19 $\mathrm{f}(x) = 12x^3 + 5x^2 + 2x - 1$.
a Show that $(4x - 1)$ is a factor of $\mathrm{f}(x)$ and write $\mathrm{f}(x)$ in the form $(4x - 1)(ax^2 + bx + c)$.
b Hence, show that the equation $12x^3 + 5x^2 + 2x - 1 = 0$ has exactly 1 real solution.

1.14 Polynomials

Chapter 7 · Original slide 16

3: The Factor Theorem

16

$$x^3 + x^2 - 4x - 4 = (x - 2)\left(x^2 + 3x + 2\right)$$

We can see that $(x - 2)$ is a factor of $x^3 + x^2 - 4x - 4$.
What would happen if x is 2?

Chapter 7 · Original slide 17

Examples (in textbook)

17

Show that $(x - 2)$ is a factor of $x^3 + x^2 - 4x - 4$.

Fully factorise $2x^3 + x^2 - 18x - 9$.

1.14 Polynomials

Chapter 7 · Original slide 18

Using Factor Theorem to find unknown coefficients

18

Given that $2x + 1$ is a factor of $6x^3 + ax^2 + 1$, determine the value of a.

Chapter 7 · Original slide 19

Test Your Understanding

19

Edexcel C2 May 2016 Q2

$\mathrm{f}(x) = 6x^3 + 13x^2 - 4$

No long in spec.

(a) Use the remainder theorem to find the remainder when $\mathrm{f}(x)$ is divided by $(2x + 3)$. \hfill(2)

~~(b) Use the factor theorem to show that $(x + 2)$ is a factor of $\mathrm{f}(x)$. \hfill(2)~~

(c) Factorise $\mathrm{f}(x)$ completely. \hfill(4)

Given that $3x - 1$ is a factor of $3x^3 + 11x^2 + ax + 1$, determine the value of a.

$\mathbf{f}\left(\frac{1}{3}\right.$
$\frac{7}{3}+$

1.14 Polynomials

Chapter 7 · Original slide 20

Exercise 7C – Y1 Pages 146

20

13 $\mathrm{f}(x) = 3x^3 - 12x^2 + 6x - 24$.
a Use the factor theorem to show that $(x - 4)$ is a factor of $\mathrm{f}(x)$.
b Hence, show that 4 is the only real root of the equation $\mathrm{f}(x) = 0$.

Chapter 7 · Original slide 21

Exercise 7C – Y1 Pages 145-146

21

14 $\mathrm{f}(x) = 4x^3 + 4x^2 - 11x - 6$.
a Use the factor theorem to show that $(x + 2)$ is a factor of $\mathrm{f}(x)$.
b Factorise $\mathrm{f}(x)$ completely.

1.14 Polynomials

Chapter 7 · Original slide 22

Exercise 7C – Y1 Pages 145-146

22

15 a Show that $(x-2)$ is a factor of $9x^4-18x^3-x^2+2x$.
b Hence, find four real solutions to the equation $9x^4-18x^3-x^2+2x=0$.

Chapter 7 · Original slide 23

Exercise Extension

23

1 *[MAT 2006 1E]* The cubic

$$x^3+ax+b$$

Has both $(x-1)$ and $(x-2)$ as factors. Determine the values of a and b.

$\mathbf{a=-7, b=6}$

2 [MAT 2009 1I]

$$n^2x^{2n+3}-25nx^{n+1}+150x^7$$

Has x^2-1 as a factor
A) for no values of n;
B) for $n=10$ only;
C) for $n=15$ only;
D) for $n=10$ and $n=15$ only.

$\mathbf{x^2-1=(x+1)(x-1)}$ **so both factors.**

$$\mathbf{f(1)=n^2-25n+150=0}$$

If n is odd: $\mathbf{f(-1)=-n^2-25n-150=0}$

The **remainder theorem** states that if $f(x)$ is divided by $(x-a)$, the remainder is $f(a)$. This similarly works whenever a makes the divisor 0.

3 [MAT 2013 1G] Let $n \geq 2$ be an integer and $p_n(x)$ be the polynomial

$$p_n(x)=(x-1)+(x-2)+\ldots+(x-n)$$

What is the remainder, in terms of n, when $p_n(x)$ is divided by $p_{n-1}(x)$?