1.23 e and Natural Logarithms

Chapter 14 · Original slide 8

$y = e^x$

8

Click ↓

Function	Gradient	
$y = 1^x$		>
$y = 1.5^x$		>
$y = 2^x$		>
$y = 2.5^x$		>
$y = 3^x$		>
$y = 3.5^x$		>

Compare each exponential function against its respective gradient function. What do you notice?

You won't yet be able to differentiate general exponential functions $y = a^x$ until Year 2.
But I've calculated the gradient functions for you – click the black arrow to reveal the graph and gradient function.

Chapter 14 · Original slide 9

$y = e^x$

9

Function	Gradient	
$y = 1^x$	$\frac{dy}{dx} = 0$	>
$y = 1.5^x$	$\frac{dy}{dx} = 0.41 \times 1.5^x$	>
$y = 2^x$	$\frac{dy}{dx} = 0.69 \times 2^x$	>
$y = 2.5^x$	$\frac{dy}{dx} = 0.92 \times 2.5^x$	>
$y = 3^x$	$\frac{dy}{dx} = 1.10 \times 3^x$	>
$y = 3.5^x$	$\frac{dy}{dx} = 1.25 \times 3.5^x$	>

$y = 2.5^x$ and $y = 3^x$ seem to be similar to their respective gradient functions. So is there a base between 2.5 and 3 where the **function is equal to its gradient function**?

$e = 2.71828...$ is known as **Euler's Number.**

It is one of the five most fundamental constants in mathematics (0, 1, i, e, π).

It has the property that:

$$y = e^x \quad \rightarrow \quad \frac{dy}{dx} = e^x$$

Although any function of the form $y = a^x$ is known as **an** exponential function, e^x is known as "**the**" exponential function.

You can find the exponential function on your calculator, to the right (above the "ln" key)

1.23 e and Natural Logarithms

Chapter 14 · Original slide 10

Differentiating $y = ae^{kx}$

10

If $y = e^{kx}$, where k is a constant, then $\frac{dy}{dx} = ke^{kx}$

Note: This is not a standalone rule but an application of something called the '*chain rule*', which you will encounter in Year 2.

Different e^{5x} with respect to x.

Different e^{-x} with respect to x.

Different $4e^{3x}$ with respect to x.

Chapter 14 · Original slide 41

Natural Logarithms

41

We have previously seen that $y = \log_a x$ is the inverse function of $y = a^x$.
We also saw that e^x is "**the**" exponential function.
The inverse of e^x is $\log_e x$, but because of its special importance, it has its own function name!

The inverse of $y = e^x$ is $y = \ln x$

$$\ln e^x = \mathbf{x}$$
$$e^{\ln x} = \mathbf{x}$$

Solve $e^x = 5$

Solve $2\ln x + 1 = 5$

1.23 e and Natural Logarithms

Chapter 14 · Original slide 42

Quadratics in e^x

42

In previous chapters we've already dealt with quadratics in disguise, e.g. "quadratic in sin". We therefore just apply our usual approach: either make a suitable substitution so the equation **is** then quadratic, or (strongly recommended!) **go straight for the factorisation**.

Solve $e^{2x} + 2e^x - 15 = 0$

Solve $e^x - 2e^{-x} = 1$

Chapter 14 · Original slide 43

Test Your Understanding

43

Solve $\ln(3x + 1) = 2$

Solve $e^{2x} + 5e^x = 6$

e
$($
e
x

Solve $2^x e^{x+1} = 3$ giving your answer as an exact value.

1.23 e and Natural Logarithms

Chapter 14 · Original slide 44

Exercise 14G – Y1 Page 327

44

4 $Find the exact solutions to the$ equation $e^x + 12e^{-x} = 7$.

6 Solve $3^x e^{4x-1} = 5$, giving your answer in the form $\dfrac{a + \ln b}{c + \ln d}$

Chapter 14 · Original slide 45


Exercise 14G – Y1 Page 328

45

8 The graph of $y = 3 + \ln(4 - x)$ is shown to the right.

a State the exact coordinates of point A. \hfill (1

b Calculate the exact coordinates of point B. \hfill (3 m



Challenge: The graph of the function passes through (0,5) and (6,10). Given that the line $y = 2$ is an asymptote to the graph, show that $B = \frac{1}{6}\ln\left(\frac{8}{3}\right)$