1.31 Differentiation: Introduction

Chapter 12 · Original slide 14

Differentiating x^n

14

Thankfully, there's a quick way to differentiate terms of the form x^n (where n is a constant) with having to use first principles every time:

! If $y = ax^n$ then $\frac{dy}{dx} = nax^{n-1}$ (where a, n are constants)
i.e. multiply by the power and reduce the power by 1

Examples:

$y = x^5 \quad \rightarrow \quad \frac{dy}{dx} =$

$f(x) = x^{\frac{1}{2}} \quad \rightarrow$

$y = 2x^6 \quad \rightarrow$

$f(x) = \frac{x}{x^4} =$ $\rightarrow$

$y = \sqrt{x^6} =$ $\rightarrow$

Why would it be incorrect to say that $y=2^x$ differentiates to $\frac{dy}{dx}=x2^{x-1}$?

Chapter 12 · Original slide 15

Test Your Understanding

15

1. $y = x^7 \quad \rightarrow \quad \frac{dy}{dx} =$
2. $y = 3x^{10} \quad \rightarrow$
3. $f(x) = \frac{x^{\frac{1}{2}}}{x^2} =$ $\rightarrow$
4. $y = ax^a \quad \rightarrow$
5. $f(x) = \sqrt{49x^7} =$ $\rightarrow \quad f'(x) =$

1.31 Differentiation: Introduction

Chapter 12 · Original slide 16

Differentiating Multiple Terms

16

Differentiate $y = x^2 + 4x + 3$

First thing to note:
If $y = f(x) + g(x)$ then $\frac{dy}{dx} = f'(x) + g'(x)$
i.e. **differentiate each term individually in a sum/subtraction**.

$\frac{dy}{dx} =$

Chapter 12 · Original slide 17

Quickfire Questions

17

1. $y = 2x^2 - 3x \rightarrow \frac{dy}{dx}$
2. $y = 4 - 9x^3 \rightarrow \frac{dy}{dx}$
3. $y = 5x + 1 \rightarrow$
4. $y = ax \rightarrow$ (where a is a constant)
5. $y = 6x - 3 + px^2 \rightarrow \frac{dy}{dx}$ (where p is a constant)

1.31 Differentiation: Introduction

Chapter 12 · Original slide 18

Harder Example

18

Let $f(x) = 4x^2 - 8x + 3$
a) Find the gradient of $y = f(x)$ at the point $\left(\frac{1}{2}, 0\right)$
b) Find the coordinates of the point on the graph of $y = f(x)$ where the gradient is 8.
c) Find the gradient of $y = f(x)$ at the points where the curve meets the line $y = 4x - 5$.

a

b

c

Chapter 12 · Original slide 19

Test Your Understanding

19

Let $f(x) = x^2 - 4x + 2$
a) Find the gradient of $y = f(x)$ at the point $(1, -1)$
b) Find the coordinates of the point on the graph of $y = f(x)$ where the gradient is 5.
c) Find the gradient of $y = f(x)$ at the points where the curve meets the line $y = 2 - x$.

a

b

c

1.31 Differentiation: Introduction

Chapter 12 · Original slide 20

Exercise 12D – Pearson Pure Y1 page 265-266

20

5 Find the gradients of the curve $y = x^2 - 5x + 10$ at the points A and B where the curve meets the line $y = 4$.

Chapter 12 · Original slide 21

Exercise 12D – Pearson Pure Y1 page 265-266

21

6 Find the gradients of the curve $y = 2x^2$ at the points C and D where the curve meets the line $y = x + 3$.


1.31 Differentiation: Introduction

Chapter 12 · Original slide 22

Exercise 12D – Pearson Pure Y1 page 265-266

22

7\quad $\mathrm{f}(x) = x^2 - 2x - 8$.
a Sketch the graph of $y = \mathrm{f}(x)$.
b On the same set of axes, sketch the graph of $y = \mathrm{f}'(x)$.
c Explain why the x-coordinate of the turning point of $y = \mathrm{f}(x)$ is the same as the x-coordinate of the point where the graph of $y = \mathrm{f}'(x)$ crosses the x-axis.



Chapter 12 · Original slide 23

Differentiating Harder Expressions

23

If your expression isn't a sum of x^n terms, simply manipulate it until it is!

1. Turn roots into powers:

$y = \sqrt{x} = x$ $\rightarrow$ $\frac{dy}{dx} = \frac{1}{2}$

$y = \frac{1}{\sqrt[3]{x}} = x$ $\rightarrow$ $\frac{dy}{dx} =$

2. Split up fractions.

$y = \frac{x^2+3}{\sqrt{x}} = \frac{x^2+}{x^{\frac{1}{2}}}$

$\rightarrow$ $\frac{dy}{dx} = \frac{3}{2}x^{\frac{1}{2}}$

3. Expand out brackets.

$y = x^2(x-3) =$

$\rightarrow$ $\frac{dy}{dx}$

4. Beware of numbers in denominators!

$y = \frac{1}{3x} =$ $\rightarrow$ $\frac{d}{d}$ x^{-2}

1.31 Differentiation: Introduction

Chapter 12 · Original slide 24

Test Your Understanding

24

Differentiate the following.

Chapter 12 · Original slide 25

Exercise 12E – Pearson Pure Y1 page 267-268

25

6\quad $\mathrm{f}(x) = \frac{12}{p\sqrt{x}} + x$, where p is a real constant and $x > 0$. Given that $\mathrm{f}'(2) = 3$, find p, giving your answer in the form $a\sqrt{2}$ where a is a rational number.

1.31 Differentiation: Introduction

Chapter 12 · Original slide 26

Exercise 12E – Pearson Pure Y1 page 267-268

26

7\quad $\mathrm{f}(x) = (2 - x)^9$.
a Find the first 3 terms, in ascending powers of x, of the binomial expansion of $\mathrm{f}(x)$, giving each term in its simplest form.
b If x is small, so that x^2 and higher powers can be ignored, show that $\mathrm{f}'(x) \approx 9216x - 2304$.

Chapter 12 · Original slide 27

Extension

27

[MAT 2013 1E]
The expression $\frac{d^2}{dx^2}\left[(2x-1)^4(1-x)^5\right] + \frac{d}{dx}\left[(2x+1)^4(3x^2-2)^2\right]$ is a polynomial of degree:
A) 9
B) 8
C) 7
D) less than 7
E)

This just means “differentiate twice”. The ‘second derivative’ comes later in this chapter.

Full expansion is not needed. The highest power term in the first

1.31 Differentiation: Introduction

Chapter 12 · Original slide 42

Second Order Derivatives

42

When you differentiate once, the expression you get is known as the **first derivative**. Unsurprisingly, when we differentiate a second time, the resulting expression is known as the **second derivative**. And so on…

Original Function	First Derivative	Second Derivative
$y = x^4$ (Leibniz's)	$\frac{dy}{dx} = 4x^3$	$\frac{d^2y}{dx^2} = 12x^2$
$y = x^4$	$y\prime = 4x^3$	$y\prime\prime = 12x^2$
$y = x^4$ (Newton's)	$\dot{y} = 4x^3$	$\ddot{y} = 12x^2$
$f(x) = x^4$ (Lagrange's)	$f\prime(x) = 4x^3$	$f\prime\prime(x) = 12x^2$

You can similarly have the third derivative ($\frac{d^3y}{dx^3}$), although this is no longer in the A Level syllabus. We'll see why might use the second derivative soon…

Chapter 12 · Original slide 43

Just for your interest…

43

How does the notation $\frac{d^2y}{dx^2}$ work?
Why are the squareds where they are?

Suppose that $y = x^3 + 1$.
Then when we write $\frac{dy}{dx}$, we're effectively doing $\frac{d(x^3+1)}{dx}$ (by substitution), although this would typically be written:

$$\frac{d}{dx}\left(x^3 + 1\right)$$

The $\frac{d}{dx}(\dots)$ notation is quite handy, because it behaves as a function and allows us to write the original expression and the derivative within a single equation:

$$\frac{d}{dx}\left(x^3 + 1\right) = 3x^2$$

Therefore, if we wanted to differentiate y twice, we'd do:

$$\frac{d}{dx}\left(\frac{d}{dx}(y)\right)$$
$$= \frac{d^2}{dx^2}(y)$$
$$= \frac{d^2y}{dx^2}$$

1.31 Differentiation: Introduction

Chapter 12 · Original slide 44

Examples

44

If $y = 3x^5 + \frac{4}{x^2}$, find $\frac{d^2y}{dx^2}$.

If $f(x) = 3\sqrt{x} + \frac{1}{2\sqrt{x}}$, find $f''(x)$.

Chapter 12 · Original slide 45

Test Your Understanding

45

If $y = 5x^3 - \frac{x}{3\sqrt{x}}$, find $\frac{d^2y}{dx^2}$.

1.31 Differentiation: Introduction

Chapter 12 · Original slide 46

Exercise 12H – Pearson Pure Y1 page 272

46

3 Given that $y = (2x - 3)^3$, find the value of x when $\frac{\mathrm{d}^2y}{\mathrm{d}x^2} = 0$.

Chapter 12 · Original slide 47

Exercise 12H – Pearson Pure Y1 page 272

47

4\quad $\mathrm{f}(x) = px^3 - 3px^2 + x^2 - 4$. When $x = 2$, $\mathrm{f}''(x) = -1$. Find the value of p.