1.27 Reduction to Linear Form

Chapter 14 · Original slide 46


Graphs for Exponential Data

46

In Science and Economics, **experimental data often has exponential growth**, e.g. bacteria in a sample, rabbit populations, energy produced by earthquakes, my Twitter followers over time, etc.

Because exponential functions increase rapidly, it tends to look a bit rubbish if we tried to draw a suitable graph:

Take for example "Moore's Law", which hypothesised that the processing power of computers would double every 2 years. Suppose we tried to plot this for computers we sampled over time:




If we tried to force all the data onto the graph, we would end up making most of the data close to the horizontal axis. This is not ideal.

But suppose we **took the log** of the number of transistors for each computer. Suppose the number of transistors one year was y, then doubled 2 years later to get $2y$.

When we log (base 2) these:

$$\begin{aligned} y &\rightarrow \log_2 y \\ 2y &\rightarrow \log_2(2y) = \log_2 2 + \log_2 y \\ &\qquad = 1 + \log_2 y \end{aligned}$$


The logged value only increased by 1! Thus **taking the log of the values turns exponential growth into linear growth (because each time we would have doubled, we're now just adding 1), and the resulting graph is a straight line.**



Chapter 14 · Original slide 47

Graphs for Exponential Data

47



Richter Scale

Because the energy involved in **earthquakes** decreases exponentially from the epicentre of the earthquake, such energy values recorded from different earthquakes would **vary wildly**.

The **Richter Scale** is a **logarithmic scale**, and takes the log (base 10) of the amplitude of the waves, giving a more even spread of values in a more sensible range.

(The largest recorded value on the Richter Scale is 9.5 in Chile in 1960, and 15 would destroy the Earth completely – evil scientists take note)


The result is that an earthquake just 1 greater on the Richter scale would in fact be 10 times as powerful.

1.27 Reduction to Linear Form

Chapter 14 · Original slide 48

Other Non-Linear Growth

48



We would also have similar graphing problems if we tried to plot data that followed some **polynomial function** such as a quadratic or cubic.

We will therefore look at the process to convert a **polynomial graph into a linear one**, as well as a **exponential graph into a linear one…**

Chapter 14 · Original slide 49

Turning non-linear graphs into linear ones

49

Case 1: Polynomial $\rightarrow$ Linear

Suppose our original model was a polynomial one*:

$$y = ax^n$$

Then taking logs of both sides:

$\log y = \log ax^n$


$\log y = \log a + n \log x$

We can compare this against a straight line:

$$Y = mX + c$$

! If $y = ax^n$, then the graph of $\log y$ against $\log x$ will be a straight line wih gradient n and vertical intercept $\log a$.

!
!
!
!
!



* We could also allow non-integer n; the term would then not strictly be polynomial, but we'd still say the function had "polynomial growth".

Case 2: Exponential $\rightarrow$ Linear

Suppose our original model was an exponential one:

$$y = ab^x$$

Then taking logs of both sides:

$\log y = \log ab^x$


$\log y = \log a + x \log b$

Again we can compare this against a straight line:

$$Y = mX + c$$

! If $y = ab^x$, then the graph of $\log y$ against x will be a straight line with gradient $\log b$ and vertical intercept $\log a$.

!
!
!
!
!



The key difference compared to Case 1 is that we're **only logging the y values** (e.g. number of transistors), not the x values (e.g. years elapsed). **Note that you do not need to memorise the contents of these boxes and we will work out from scratch each time…**

In summary, logging the y-axis **turns an exponential graph into a linear one**.
Logging **both** the x and y-axis turns a polynomial graph into a linear one.


1.27 Reduction to Linear Form

Chapter 14 · Original slide 50

Example

50

[Textbook] The graph represents the growth of a population of bacteria, P, over t hours. The graph has a gradient of 0.6 and meets the vertical axis at $(0, 2)$ as shown.
A scientist suggest that this growth can be modelled by the equation $P = ab^t$, where a and b are constants to be found.
a. Write down an equation for the line.
b. Using your answer to part (a) or otherwise, find the values of a and b, giving them to 3 sf where necessary.
c. Interpret the meaning of the constant a in this model.



a

Equation of straight line is $y = mx + c$
where here:

b Just like on previous slide, start with the model then log it:

Recall that $\log a$ means $\log_{10} a$

c Comparing with our straight line equation in (a):

Recall that the coefficient of an exponential term gives the 'initial value'.

Chapter 14 · Original slide 51

Example

51

[Textbook] The table below gives the rank (by size) and population of the UK's largest cities and districts (London is number 1 but has been excluded as an outlier).

City	B'ham	Leeds	Glasgow	Sheffield	Bradford
Rank, R	2	3	4	5	6
Population, P	1 000 000	730 000	620 000	530 000	480 000


The relationship between the rank and population can be modelled by the formula:
$P = aR^n$ where a and n are constants.
a) Draw a table giving values of $\log R$ and $\log P$ to 2dp.
b) Plot a graph of $\log R$ against $\log P$ using the values from your table and draw the line of best fit.
c) Use your graph to estimate the values of a and n to two significant figures.

Textbook Error: They use $R = aP^n$ but then plot $\log P$ against $\log R$.

a

logR					
logP					

b



c First get equation of straight line:

As with previous example, let's log the original model so we can compare against our straight line:

Comparing this with our straight line:

1.27 Reduction to Linear Form

Chapter 14 · Original slide 52

Test Your Understanding

52

A website wants to predict his number of Twitter followers P, t years from the start 2015. He predicts that his followers will increase exponentially according to the model $P = ab^t$, where a, b are constants that he wishes to find.

He records his followers at certain times. Here is the data:

Years t after 2015: 0.7 1.3 2.2
Followers P: 2353 3673 7162

a) Draw a table giving values of t and $\log P$ (to 3dp).
b) A line of best fit is drawn for the data in your new table, and it happens to go through the first data point above (where $t = 0.7$) and last (where $t = 2.2$). Determine the equation of this line of best fit. (The y-intercept is 3.147)
c) Hence, determine the values of a and b in the model.
d) Estimate how many followers the website will have at the start of 2020 (when $t = 5$).

a

t	0.7	1.3	2.2
logP			

b

c

d

Notes

1. We want to find the **parameters of a model**, e.g. $P = ab^t$ that **best fits the data** (in this case the parameters we want to find are a and b).
2. If the data had a linear trend, then this would be easy! We know from KS3 that we'd just plot the data, find the line of best fit, then use the gradient and y-intercept to work out the m and c in our linear model.
3. But the original data wasn't linear, and it would be harder to draw an 'exponential curve of best fit'.
4. We therefore log the model so that the plotted data then roughly forms a straight line, and then we can then draw a (straight) line of best fit.
5. The gradient and y-intercept of this line then allows us to estimate the parameters a and b in the original model that best fit the data.

The process of finding parameters in a model, that best fits the data, is known as **regression**.

Chapter 14 · Original slide 53

Exercise 14H Page 331-333

53

5 Kleiber's law is an empirical law in biology which connects the mass of an animal, m, to its resting metabolic rate, R. The law follows the form $R = am^b$, where a and b are constants.

The table below contains data on five animals.

Animal	Mouse	Guinea pig	Rabbit	Goat	Cow
Mass, m (kg)	0.030	0.408	4.19	34.6	650
Metabolic rate R (kcal per day)	4.2	32.3	195	760	7637

a Copy and complete this table giving values of $\log R$ and $\log m$ to 2 decimal places. \hfill (1 m

$\log m$	-1.52				
$\log R$	0.62	1.51	2.29	2.88	3.88

b Plot a graph of $\log R$ against $\log m$ using the values from your table and draw in a line of best fit.

c Use your graph to estimate the values of a and b to two significant figures.

d Using your values of a and b, estimate the resting metabolic rate of a human male with a mass of 80 kg.

1.27 Reduction to Linear Form

Chapter 14 · Original slide 54

Exercise 14H Page 331-333

54

8 A scientist is modelling the number of people, N, who have fallen sick with a virus after t days.

From looking at this graph, the scientist suggests that the number of sick people can be modelled by the equation $N = ab^t$, where a and b are constants to be found.

The graph passes through the points $(0, 1.6)$ and $(10, 2.55)$.

a Write down the equation of the line. \hfill (2

b Using your answer to part **a** or otherwise, find the values of a and b, giving them to 2 significant figures. \hfill (4

c Interpret the meaning of the constant a in this model. \hfill (

d Use your model to predict the number of sick people after 30 days.

Give one reason why this might be an overestimate. \hfill (2

