Applications of Integration

Questions 1–117

2.34 — Question 1. Find the exact mean value of each of the following on the given intervals:

a) $f(x) = \pi x \cos(\pi x^2)$ on the interval $\left[0, \frac{\sqrt{2}}{2}\right]$

b) $f(x) = \operatorname{cosec}^2 x$ on the interval $\left[\frac{\pi}{4}, \frac{\pi}{3}\right]$

c) $f(x) = \dfrac{x^2}{5x^3 + 1}$ on the interval $[0, 1]$

d) $f(x) = \sin^2 4x$ on the interval $\left[0, \frac{\pi}{4}\right]$

e) $f(x) = 2^x$ on the interval $[2, 5]$

f) $f(x) = \sin 2x \cos^3 2x$ on the interval $\left[0, \frac{\pi}{12}\right]$

2.34 — Question 2. Find the mean value of the function $f(x) = \dfrac{x}{e^{-3x}}$ in the interval $0 \leq x \leq a$, giving your answer in terms of a.

2.34 — Question 3. The mean value of the function $f(x) = \dfrac{2}{\sqrt{3x+1}} + k$ in the interval $[0, 8]$ where k is a positive constant is 2. Find the value of k.

2.34 — Question 4. The mean value of the function $f(x) = \sqrt{x+1}$ in the interval $[0, 3]$ is half the mean value of the function $g(x) = \frac{3}{7}x^2$ for the interval $[0, k]$ where k is a positive number. Find the exact value of k.


2.34 — Question 5. The velocity of a car after t seconds is given by $v = \dfrac{30t}{t+5}$ metres per second.

a) Show that the mean velocity for $0 \leq t \leq 20$ is 17.9 m s^{-1} to 3 significant figures.

b) Use the mean value theorem to calculate the mean acceleration of the car for the same time interval.

2.34 — Question 6. Given that the function f is defined by $f(x) = \ln x$ for $1 \leq x \leq 4$, **a)** Find the mean value of f. Another function g is defined by $g(x) = f(x) + c$ for $1 \leq x \leq 4$. **b)** Given that the mean value of g is $\frac{4}{3}\ln 4$, state the value of c.

2.34 — Question 7. The diagram below shows the curve $f(x) = \sin x$ with shaded regions A and B enclosed by the y-axis, the line $y = k$ and $x = \frac{\pi}{2}$. The areas A and B are equal.



Determine the value of k.

2.34 — Question 8. The root mean square (RMS) of a function $f(x)$ defined for $a \leq x \leq b$ over this interval is

$$\sqrt{\frac{1}{b-a}\int_a^b [f(x)]^2\,dx}.$$

Find the RMS of $f(x) = 4\cos x$ from $x = 0$ to $x = 2\pi$ giving your answer in exact form.

2.34 — Question 9. a) Show that $\cos^4\theta = \frac{1}{8}(3 + 4\cos 2\theta + \cos 4\theta)$. In a mechanical system, the velocity v m s^{-1} of a vibrating component at time t seconds is given by $v = a\cos^2(kt)$, where a and k are positive constants. The effective speed of the component over one complete cycle of $\frac{\pi}{k}$ seconds is defined to be the square root of the mean value of v^2 over this time interval. **b)** Show that the effective speed of the component is $\frac{\sqrt{3}a}{2\sqrt{2}}$ m s^{-1}. **Prerequisite Knowledge for Question 10: Hyperbolic Functions**

2.34 — Question 10. For each of the following functions, defined over the domain $-k\pi \leq x \leq k\pi$ for positive integer k, find the mean value of $[f(x)]^n$

a) $\sinh(x)$, n is an odd positive integer.
b) $\cosh(ix)$, n is an odd positive integer.
c) $\sinh(ix)$, n is an odd positive integer.

The following questions require the use of A-Level Further Maths Integration techniques

2.34 — Question 11. Find the exact mean value of each of the following for the given intervals:

a) $f(x) = \dfrac{1}{9+x^2}$ in the interval $[0, 3]$

b) $f(x) = \dfrac{1}{\sqrt{1-4x^2}}$ in the interval $\left[0, \frac{1}{4}\right]$

c) $f(x) = \dfrac{1}{x^2+7}$ in the interval $[0, a]$ for positive constant a

Prerequisite Knowledge for Questions 12–15: Calculus of Hyperbolic Functions

2.34 — Question 12. Find the exact mean value of the function $f(x) = \dfrac{1}{\sqrt{x^2-6x+16}}$ between $x = 0$ and $x = 6$.

2.34 — Question 13. Find the mean value of the function $f(x) = \dfrac{1}{\sqrt{x^2 - 9}}$ in the interval $3 \leq x \leq a$, giving your answer in exact logarithmic form.

2.34 — Question 14. The intensity of a signal, I watts per square metre, at time t hours after a transmitter is switched on is modelled by

$$I = 12 + 8\,\text{sech}^2(0.5t)$$

for $t \geq 0$. Determine the mean signal intensity during the first four hours after the transmitter is switched on, giving your answer correct to three significant figures.

2.34 — Question 15. The brightness of a light-emitting diode, B in lumens, t minutes after it is switched on, is modelled by

$$B = 50 + 20\,\text{cosech}^2(0.2t)$$

for $t > 0$. Determine the mean brightness of the LED during the second minute after it is switched on, giving your answer to three significant figures.

2.34 — Question 16. The velocity of a cyclist after t seconds is given by

$$v = \frac{18}{\sqrt{t^2 + 9}}$$

metres per second for $t \geq 0$.

a) Show that the mean velocity of the cyclist for $0 \leq t \leq 12$ is 3.14 m s^{-1} to three significant figures.

b) Use the mean value theorem to calculate the mean acceleration of the cyclist for the same time interval giving your answer to three significant figures.

2.34 — Question 17. Given that $\cos\theta = \frac{a}{5}$, find the mean value of the function


$$f(x) = \frac{1}{\sqrt{25 - x^2}}$$

in the interval $0 \leq x \leq a$, giving your answer in the form $p + q\theta$ where p and q are constants to be found in terms of a. **Prerequisite Knowledge for Question 18: A-Level Further Maths Partial Fractions**

2.34 — Question 18. The diagram below shows part of the curve

$$f(x) = \frac{x + 2}{(x^2 + 2)(x - 1)}$$

with shaded regions A and B enclosed by the line $y = k$ and the lines $x = \frac{3}{2}$ and $x = \frac{5}{2}$.




The areas A and B are equal. Determine the exact value of k.

2.35 — Question 1. Find the exact volume created by rotating the curve through 2π radians around the x-axis for each of the following:


a) $y = (2x-1)^{1/3}$ for $1 \leq x \leq 14$
b) $y = \sec x$ for $\frac{\pi}{6} \leq x \leq \frac{\pi}{3}$
c) $y = \sqrt{\ln x}$ for $1 \leq x \leq 2$
d) $y = \sqrt{\frac{2\cos x}{1+\sin x}}$ for $0 \leq x \leq \frac{\pi}{2}$
e) $y = \sqrt{x}\ \operatorname{cosec} x$ for $\frac{\pi}{4} \leq x \leq \frac{\pi}{2}$
f) $y = \sec x + \tan x$ for $0 \leq x \leq \frac{\pi}{4}$

2.35 — Question 2. For each of the following, the shaded region is bounded by the given lines, the curve and the x-axis. In each case, find the exact volume of the solid generated when the region is rotated 2π radians about the x-axis:

a) $y = \sqrt{\cos x}$, $x = \frac{\pi}{4}$



b) $y = \sqrt{x}\ln x$, $x = 4$



2.35 — Question 3. A region R is bounded by the curve $y = \sin x$, the x-axis, the y-axis and the line $x = \pi$. The region is rotated through 2π radians about the x-axis to form a solid of revolution.

a) Sketch the region R.

b) Find the exact volume of the solid formed.

Prerequisite Knowledge for Questions 4–5: Calculus of Hyperbolic Functions

2.35 — Question 4. A region R is bounded by the curve $y = \sinh x$, the x-axis, the y-axis and the line $x = \ln 5$. The region is rotated through 2π radians about the x-axis to form a solid of revolution.

a) Sketch the region R.

b) Find the exact volume of the solid formed.

2.35 — Question 5. A region R is bounded by the curve $y = \cosh x$, the x-axis, the y-axis and the line $x = \ln 3$. The region is rotated through 2π radians about the x-axis to form a solid of revolution.

a) Sketch the region R.


b) Find the exact volume of the solid formed.

2.35 — Question 6. A shaded region is bounded by the curve $y^2 = 2x\cos x$, the y-axis and the line $x = P$ where P is the point shown in the diagram. The region is rotated around the x-axis until a solid of revolution is generated.

a) Find the value of P.

b) Determine the volume of the solid generated giving your answer in exact form.

t shown in the diagram.



2.35 — Question 7. A designer is designing a candle holder which is 12 cm long, with flat circular ends. The candle holder is symmetrical about its midpoint and should have the following diameters:

- 5 cm at each end.
- 2 cm at the midpoint.

The shape is modelled by rotating a curve about the x-axis. The origin O is at the centre of the candle holder and the ends are at $x = \pm 6$. Two possible equations for the curve are:

$$\text{Model A: } y = a + kx^2 \qquad \text{Model B: } y = a + m\left(1 - \cos\frac{\pi x}{6}\right)$$

for constants a, k and m.

a) State the value of a which is the radius of the candle holder at the midpoint.

b) Show that $k = \frac{1}{24}$ and $m = \frac{3}{4}$.

c) Use integration to find the volume predicted by both models giving your answer to three significant figures.

d) The designer wanted the model to change gradually near the midpoint and more rapidly towards the ends. Suggest which of the two models is more appropriate.

Prerequisite Knowledge for Questions 8–10: A-Level Maths Partial Fractions

2.35 — Question 8.

a) Write $\dfrac{6x+11}{(x+3)(2x-1)}$ as partial fractions.

A region R is bounded by the curve $y = \sqrt{\dfrac{6x+11}{(x+3)(2x-1)}}$, the x-axis, the line $x = 1$ and the line $x = 4$. The region is rotated through 2π radians about the x-axis to form a solid of revolution.

b) Find the exact volume of the solid formed.


2.35 — Question 9. A region R is bounded by the curve $y = \dfrac{3x+4}{(x+2)\sqrt{x}}$, the x-axis, the line $x = 1$ and the line $x = 3$. The region is rotated through 2π radians about the x-axis to form a solid of revolution. Show that the volume is given by

$$\pi\left(\ln\left(\frac{5^a}{3^b}\right) - \frac{4}{15}\right)$$

where a and b are integers to be found.

2.35 — Question 10. A shaded region R is bounded by the curve $y = \dfrac{1}{\sqrt{k-x^2}}$, the x-axis, the y-axis and the line $x = \frac{k}{2}$, where $k < 4$ is a positive constant. The region is rotated around the x-axis until a solid of revolution is generated.

< 4 is a positive constant.



a) Show that the exact volume of the solid generated is given by

$$V = \frac{\pi}{2\sqrt{k}}\ln\left(\frac{2\sqrt{k}+k}{2\sqrt{k}-k}\right).$$

b) Given that the region lies entirely between $x = 0$ and $x = \frac{1}{2}$, find the exact volume.

2.35 — Question 11. A region R is bounded by the curves $y = 1 + \sin x$ and $y = 1 + \cos x$, the line $x = \frac{\pi}{4}$ and the line $x = \frac{5\pi}{4}$. The region is rotated through 2π radians about the x-axis.

a) Sketch the region R.

b) Show that the exact volume of the solid formed is given by $k\sqrt{2}\pi$, where k is a constant to be found.

Prerequisite Knowledge for Question 12: De Moivre's Theorem

2.35 — Question 12.

a) Use De Moivre's theorem to determine the constants a, b and c such that $\sin^4\theta = a\cos 4\theta + b\cos 2\theta + c$.


A region is bounded by the curve $y = \sin^2 x + 1$, the x-axis, the y-axis and the line $x = \pi$. The region is rotated through 2π radians about the x-axis to form a solid of revolution.

b) Find the exact volume of the solid formed.

2.35 — Question 13.

a) Using the substitution $u = 2 - \cos x$, find the exact value of $\displaystyle\int_0^{\pi} \frac{\sin x}{\sqrt{2 - \cos x}}\, dx$.

A region R is bounded by the curve $y = 2(\sin x)^{1/2}(2 - \cos x)^{-1/4}$, the x-axis, the y-axis and the line $x = \pi$. The region is rotated through 2π radians about the x-axis.



b) Find the exact volume of the solid formed.

2.35 — Question 14. A region is bounded by the curve $y = \sqrt{xe^x}$, the x-axis and the line $x = \ln k$ for constant $k > 0$. The region is rotated through 2π radians about the x-axis to form a solid of revolution of volume $\pi(81 \ln 3 - 20)$ units^3. Determine the value of k. **Prerequisite Knowledge for Question 15: Harmonic Form**

2.35 — Question 15. The region R is bounded by the curve with equation

$$y = \frac{8}{2\cos x + \sqrt{5}\sin x}$$

the x-axis, the y-axis and $x = \frac{\pi}{4}$. Show that when the region R is rotated around the x-axis until a solid of revolution is generated, its volume is $32\pi(\sqrt{5}-2)$ units^3. **Prerequisite Knowledge for Question 16: Applications of De Moivre's Theorem**

2.35 — Question 16.

a) Given that $z - \frac{1}{z} = 2i\sin\theta$, show that

$$\sin^6\theta = \frac{5}{16} - \frac{15}{32}\cos 2\theta + \frac{3}{16}\cos 4\theta - \frac{1}{32}\cos 6\theta.$$

A decorative glass ornament with a flat base is symmetrical about its horizontal axis. The curve of the glass ornament may be modelled by the curve with equation

$$y = H\sin^3\left(\frac{\pi x}{8}\right) \qquad \text{for } 0 \le x \le 4$$

where H is a positive constant and x is in centimetres.


b) Given that the maximum height of the dome is 3 cm, write down the value of H.

c) Using algebraic integration and the result from part (a), determine the exact volume of the glass dome, in cm^3.

d) State one limitation of the model.

2.35 — Question 17. The shaded region is bounded by the line $y = x$ and the curve $y = \dfrac{1}{2\sqrt{2} - x}$. The region is

line $y = x$ and the curve $y = \frac{1}{2\sqrt{2}-x}$



rotated around the x-axis until a solid of revolution is generated. Determine the volume of the solid generated giving your answer in exact form. **Prerequisite Knowledge for Question 18: Polar Curves**

2.35 — Question 18. A curve is given by the polar equation $r = \dfrac{3}{2 - \cos\theta}$.


a) Show that the Cartesian equation of the curve can be written $y^2 = \dfrac{(x+3)^2}{4} - x^2$.

b) Explain why the curve is symmetrical in the x-axis.

c) A shaded region is bounded by the curve above, the x-axis and the y-axis as shown in the diagram below.

$y^2 = \frac{(x+3)^2}{4} - x^2$

A shaded region is bounded by the curve above, the x-axis and the y-axis as shown the diagram below.



The region is rotated around the x-axis until a solid of revolution is generated. Find the volume of the solid generated giving your answer in exact form.

Prerequisite Knowledge for Question 19: Polar Curves, A-Level Maths Partial Fractions and Reciprocal Trigonometric Functions

2.35 — Question 19. A curve is given by polar equation $r = 3 - 2\sec\theta$.

a) Show that the Cartesian equation of the curve can be written in the form $y^2 = \dfrac{9x^2}{(x+2)^2} - x^2$.

b) Explain why the curve is symmetrical in the x-axis.

c) Show that $y^2 = 9 + \dfrac{36}{(x+2)^2} - \dfrac{36}{x+2} - x^2$.

A shaded region is bounded by the curve above, the x-axis, the y-axis and the line $x = 1$. The region is rotated around the x-axis until a solid of revolution is generated.

d) Show that the volume is given by $\pi\left(\frac{44}{3} + 36\ln\frac{2}{3}\right)$ units^3.

The following questions require the use of A-Level Further Maths Integration techniques

2.35 — Question 20. For each of the following, the region is bounded by the x-axis, the y-axis and the given line. The region is rotated around the x-axis until a solid of revolution is generated. In each case, determine the volume of the solid generated giving your answer in exact form:

a) $y = (1 - x^2)^{-1/4}$ with line $x = \frac{1}{2}$

b) $y = \dfrac{3}{\sqrt{x^2+9}}$ with line $x = 4$

c) $y = (8 - 4x - 4x^2)^{-1/4}$ with line $x = \frac{1}{4}$

d) $y = \dfrac{1}{\sqrt{x^2+2}}$ with line $x = \sqrt{6}$

e) $y = \dfrac{1}{9+x^2}$ with line $x = 1$

Prerequisite Knowledge for Question 21: A-Level Maths Partial Fractions or Hyperbolic Functions

2.35 — Question 21. A shaded region is bounded by the curve $y = \dfrac{1}{\sqrt{4-x^2}}$ for $0 \leq x < 1$, the x-axis and the y-axis. The region is rotated around the x-axis until a solid of revolution is generated. Determine the exact volume of the volume generated. **Prerequisite Knowledge for Question 22: Hyperbolic Functions**

2.35 — Question 22. A shaded region is bounded by the curve $y = \dfrac{1}{(x^2+3)^{1/4}}$, the x-axis, the y-axis and the line $x = 2$. The region is rotated around the x-axis until a solid of revolution is generated. Show that the volume of the solid generated is given by


$$\pi \ln\left(\frac{p+\sqrt{q}}{\sqrt{r}}\right)$$

where p, q and r are integers to be found.

2.35 — Question 23. A designer is creating a fountain spout which is 2 m long, with flat circular ends. The spout is symmetrical about its midpoint and should have diameters of 4 m at the top of the spout and 0.8 m at the bottom of the spout. The shape is modelled by rotating a curve about the x-axis. The origin O is at the top of the spout. Two possible equations for the curve are:

$$\text{Model A: } y = 0.2x^2 + ax + b \qquad \text{Model B: } y = \frac{p}{x^2 + q}$$

e spout and 0.8 m at the bottom of the
out.



for constants a, b, p and q.

a) Show that $a = -1.2$ and $b = 2$.
b) Find the values of p and q.
c) Use integration to find the volume predicted by model A giving your answer to three significant figures.
d) Use integration to find the volume predicted by model B giving your answer to three significant figures.

Prerequisite Knowledge for Question 24: Integration of Inverse Hyperbolic Functions

2.35 — Question 24. A region R is bounded by the curve $y = \sqrt{\text{arcosh}(x)}$, the x-axis and the lines $x = 1$ and $x = 2$. The region is rotated through 2π radians about the x-axis to form a solid of revolution. Show that the exact volume formed is $\pi(2\ln(2 + \sqrt{3}) - \sqrt{3})$.

2.35 — Question 25. A shaded region is bounded by the curve $y = \dfrac{1}{(k-x^2)^{1/4}}$, the x-axis, the y-axis and the line $x = k$ where k is a positive constant. The region is rotated around the x-axis until a solid of revolution is generated.

a) Determine the exact volume of the solid generated giving your answer in exact form in terms of k.


b) If the region lies entirely between $x = 0$ and $x = \frac{1}{2}$, find the volume.

Prerequisite Knowledge for Question 26: Calculus with Hyperbolic Functions

2.35 — Question 26. The shaded region below is bounded by the line $y = \sqrt{2}x + \sqrt{2}$, the x-axis and the curve given by

$$y = -\frac{x-2}{\sqrt{x^2+2}}.$$

$\sqrt{x^2+2}$




The region is rotated around the x-axis until a solid of revolution is generated. Determine the volume of the solid generated giving your answer to three significant figures.

2.35 — Question 27. Find the exact volume created by rotating the curve through 2π radians around the y-axis for each of the following:

a) $x = (2y+1)^{1/3}$ for $0 \leq y \leq \frac{7}{2}$
b) $x = \operatorname{cosec} y$ for $\frac{\pi}{6} \leq y \leq \frac{\pi}{3}$
c) $x = \sqrt{\frac{2\sin y}{1+\cos y}}$ for $0 \leq y \leq \frac{\pi}{2}$
d) $x = \sqrt{y}\sec y$ for $0 \leq y \leq \frac{\pi}{4}$
e) $x = \sec y - \tan y$ for $0 \leq y \leq \frac{\pi}{4}$

2.35 — Question 28. Zaynab is designing 50 small decorative beads which she is making out of resin. The cross-section of each bead is modelled by the curve

$$9x^2 + 2y^2 - \cos\left(\frac{\pi y}{3}\right) = 5$$



for $x \geq 0$ where 1 unit represents 1 centimetre. The curve is rotated about the y-axis through 2π radians to form the bead. The top and bottom of the bead correspond to the points $(0, 1.570)$ and $(0, -1.570)$ rounded to four significant figures.

a) Determine, according to the model, the volume of one bead.
b) Zaynab has 215 cm^3 of resin available. Use your answer to part (a) to decide whether she has enough material to make all 50 beads.

2.35 — Question 29. The inner wall of a conical flask is modelled by $y = \ln x$ where $x \geq 1$. The region bounded by the curve, the y-axis and the lines $y = 0$ and $y = \ln 4$ is rotated about the y-axis. Determine the exact volume of the solid formed.

2.35 — Question 30. A tent has a smooth, curved canopy. A vertical cross-section through the centre of the tent is modelled by


$$x^2 + \sin\left(\frac{y}{h}\right) = 1$$

for $0 \leq y \leq \frac{\pi h}{2}$, where x is the horizontal distance from the centre line in metres, y is the height above the ground in metres and h is the height of the tent. Given that this model finds the exact volume of air enclosed by the tent to be $(2\pi^2 - 4\pi)$ m^3, determine the height of the tent.

2.35 — Question 31. The diagram below shows a cross-section of a design for a perfume bottle, which is formed by rotating it about the y-axis. The base of the bottle lies on the x-axis. It may be modelled using the following:

$$x = 4\sin\left(\frac{1}{2}y + \frac{\pi}{6}\right) \quad \text{for } 0 \leq y \leq k, \qquad x = \sqrt{6} - \sqrt{2} \quad \text{for } k \leq y \leq 6,$$


where units are in centimetres.



a) Show that the value of k is $\frac{3\pi}{2}$.
b) Find the diameter of the bottle at its base.
c) Determine the total volume of the perfume bottle giving your answer to three significant figures.
d) State one limitation of the model.

Prerequisite Knowledge for Question 32: Connected Rates of Change

2.35 — Question 32. A bowl has a circular horizontal cross-section. The bottom diameter of the bowl is 12 cm and the top diameter is 16 cm.



The curve along one side of the bowl, from bottom to top, may be modelled by

$$x = k + 2\sin\left(\frac{\pi y}{12}\right)$$


where x and y are in cm, k is a positive constant and $y = 0$ corresponds to the base of the vase.

a) Find the value of k.
b) Find the height of the bowl according to the model.

The bowl is being filled with water from a tap.

c) Find, in terms of h, the volume of water in the bowl when the water reaches a height of h cm.
d) If water flows in at a constant rate of 0.5 litres per minute, find the rate at which the water level is rising when the depth of water is 4 cm giving your answer to three significant figures.

2.35 — Question 33. A designer is creating a solid paperweight. The central vertical cross-section of the paperweight is described by the curve $20x^2 + \ln(y+1) = 0.1$ for $y \geq 0$.



TLMATHS.com THE FM CORE PU

A region R bounded by this curve and the y-axis is rotated through 2π radians about the y-axis to form the solid paperweight. Units are in metres.

a) Determine the volume of the paperweight giving your answer to three significant figures.
b) Given the paperweight is made from material of density 4.5 g cm^{-3}, determine its mass in kilograms.
c) Comment on the suitability of the model.


Prerequisite Knowledge for Question 34: Calculus of Hyperbolic Functions

2.35 — Question 34. A region R is bounded by the curve $y = \sinh\left(\frac{1}{2}x^2\right)$, the x-axis, the y-axis and the line $y = \sinh 1$. The region is rotated around the y-axis through 2π radians. Show that the exact volume of the solid formed is $2\pi\left(1 - \frac{1}{e}\right)$.

2.35 — Question 35. The curves C_1 and C_2 are given by equations

$$y = \frac{k-10}{k^2}x^2 + 10, \quad 0 \le x \le k, \qquad x = k\left(\frac{y}{k}\left(2 - \frac{y}{k}\right)\right)^{1/4}, \quad 0 \le y \le k,$$

respectively, where $0 < k < 10$.



A region R is bounded by C_1, C_2 and the y-axis. The region is rotated 2π radians about the y-axis. Show that the expression for the volume of the solid as a definite integral in terms of k is

$$\frac{1}{4}k^2\pi\big(k\pi - 2(k-10)\big) \text{ units}^3.$$

The following questions require the use of A-Level Further Maths Integration techniques

2.35 — Question 36. The region R is bounded by the curve $y = \sqrt{\frac{1}{x^2} - 1}$ for $x > 0$, the x-axis, the y-axis and the line $y = 1$. The region R is rotated about the y-axis to form a solid of revolution. Find the exact volume of the solid formed.

2.35 — Question 37. The equation of a curve is given by $x = \frac{1}{\sqrt[4]{k-y^2}}$ where k is a positive constant and $k > 1$. The region between the y-axis, the x-axis and the line $y = 1$ is rotated through 2π radians about the y-axis. Determine the exact volume generated giving your answer in terms of k. **Prerequisite Knowledge for Questions 38–39: Calculus with Hyperbolic Functions**

2.35 — Question 38. The equation of a curve is given by $x = \frac{1}{\sqrt[4]{k+y^2}}$ where k is a positive constant. The region between the y-axis, the x-axis and the line $y = 1$ is rotated through 2π radians about the y-axis. Given that the volume of the solid of revolution formed is $\pi \ln\left(\frac{1+\sqrt{10}}{3}\right)$ units^3, find the exact value of k.

2.35 — Question 39. A region R is bounded by the curve $y = \operatorname{arcosh} x$, the y-axis and the line $y = \frac{1}{2}\ln 5$. The line and the curve intersect at the point P.

a) Sketch the region R.
b) Find the exact coordinates of the point P.
c) The region is rotated through 2π radians about the y-axis to form a solid of revolution. Determine the exact volume of the solid formed.

Prerequisite Knowledge for Question 40: Linear Rational Functions

2.35 — Question 40. It is given that $f(x) = \dfrac{ax+7}{x+b}$ where a and b are constants. It is known that $y = f(x)$ has asymptotes $x = -3$ and $y = 4$.


a) Write down the values of a and b.
b) Sketch $y = f(x)$.

A finite region R is defined to be enclosed by the curve $y = f(x)$, the negative x-axis and the positive y-axis. Find the exact volume formed when rotating 2π radians around the:

c) x-axis
d) y-axis

Prerequisite Knowledge for Question 41: Further Maths Integration Methods

2.35 — Question 41. The equation of a curve is given by $y = \sin\left(\frac{\pi x}{2}\right)$ where $0 \leq x \leq 1$. The point P is the point of intersection of the curve with the lines $x = 1$ and $y = 1$.



The region R_1 is bounded by the curve, the line $y = 1$ and the y-axis. The region R_2 is bounded by the curve, the line $x = 1$ and the x-axis.

a) Show that the exact volume of the solid formed when rotating region R_2 about the x-axis is $\frac{\pi}{2}$ units^3.

b) Show that the exact volume of the solid formed when rotating region R_1 about the y-axis is $\left(\pi - \frac{8}{\pi}\right)$ units^3.

2.36 — Question 1. For each of the following pairs of parametric equations, the region is bounded by the given curve, the x-axis, the y-axis and any line if given. The region is rotated around the x-axis until a solid of revolution is generated. In each case, determine the volume of the solid generated giving your answer in exact form:

a) $x = t^2$, $y = t^3$ for $t \geq 0$ with line $x = 4$

b) $x = 2t^3$, $y = t^2$ for $t \geq 0$ with line $x = 16$

c) $x = t^3 + 1$, $y = t^2$ for $t \geq 0$ with line $x = 28$

d) $x = \sin t$, $y = \sqrt{\cos t}$ for $0 \leq t \leq \frac{\pi}{2}$

2.36 — Question 2. For each of the following pairs of parametric equations, the region is bounded by the given curve, the x-axis, the y-axis and any line if given. The region is rotated around the y-axis until a solid of revolution is generated. In each case, determine the volume of the solid generated giving your answer in exact form:

a) $x = \sqrt{1-t^2}$, $y = 1 + t^2$ for $0 \leq t \leq 1$

b) $x = 2\cos t$, $y = \tan t$ for $0 \leq t \leq \frac{\pi}{2}$ with line $y = 1$

c) $x = \sin^2 t\sqrt{\cos t}$, $y = -\cos t$ for $0 \leq t \leq \frac{\pi}{2}$

2.36 — Question 3. A curve is defined by parametric equations

$$x = e^{\frac{1}{2}t} \quad \text{and} \quad y = \sqrt{t-2}.$$

A region R is enclosed by the curve, the line $x = e^2$ and $x = e^3$. Show that the volume of the solid generated when R is rotated about the x-axis by 2π radians is given by $2\pi e^3$.

2.36 — Question 4. A curve is defined by parametric equations

$$x = \sqrt{\ln|t+1|} \quad \text{and} \quad y = t^2.$$

A region R is enclosed by the curve, the line $y = 1$ and $y = 9$. Show that the volume of the solid generated when R is rotated about the y-axis by 2π radians is given by $2\pi(8\ln 2 - 1)$.

2.36 — Question 5. A funnel may be modelled by a curve C defined parametrically by

$$x = \frac{4}{t} \quad \text{and} \quad y = \ln t$$

for $t \geq 1$. The region bounded by the curve C, the y-axis, and the lines $y = 0$ and $y = \ln 4$ is rotated through 2π radians about the y-axis to form a solid. All lengths are measured in centimetres.


a) Find the exact volume of the funnel.

b) State a limitation of the model.

2.36 — Question 6. A curve is defined by parametric equations

$$x = 1 - \cos t \quad \text{and} \quad y = \sqrt{\sin t}$$

for $0 \leq t \leq k$ for constant k and is shown in the diagram below. A region R is enclosed by the curve and the x-axis.



a) State the value of k.
b) Determine the volume of the solid generated when R is rotated about the x-axis by 2π radians.

2.36 — Question 7. A curve is defined by parametric equations

$$x = \sqrt{t} \quad \text{and} \quad y = e^t$$

for $0 \leq t \leq k$ for constant k. Given that the volume of the solid generated when R is rotated about the y-axis by 2π radians is $\pi(3e^4 + 1)$, determine the value of k.

2.36 — Question 8.

a) By using the substitution $u = \sin\theta$, show that

$$\int \cos^3\theta \, d\theta = \sin\theta - \frac{1}{3}\sin^3\theta + c.$$

b) Find the volume formed by the region enclosed by the curve defined by $x = \sin\theta$ and $y = \cos\theta$ for $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ when rotated 2π radians across the x-axis.

Prerequisite Knowledge for Question 9: Calculus of Hyperbolic Functions and Improper Integrals

2.36 — Question 9. A curve C is defined by parametric equations

$$x = \cosh t - \sinh t \quad \text{and} \quad y = \cosh t + \sinh t$$

for $t \leq 0$. An unbounded region is formed when the region between the curve C, the x-axis and the line corresponding to $t = 0$ is rotated through 2π about the x-axis.

a) Show that the curve can be expressed as $y = \dfrac{1}{x}$ for $x \geq 1$.

b) Hence, determine the exact volume of the solid formed.

2.36 — Question 10. The inside surface of a nozzle is modelled by the parametric curve C

$$x = ae^{-t} \quad \text{and} \quad y = bt^2$$

for $0 \leq t \leq 2$ where a and b are positive constants. The curve is rotated 2π radians about the y-axis. The bottom of the nozzle sits on the x-axis. All lengths are measured in centimetres. It is required that the nozzle has diameter 4 cm and a height of 3 cm.

a) Determine the values of a and b according to the model.
b) Find the volume of the nozzle giving your answer to three significant figures.
c) State a limitation of the model.

2.36 — Question 11. A glass ornament is made by rotating a smooth curve about a vertical axis. The profile of a glass ornament is modelled by a curve C defined parametrically by

$$x = \sqrt{t^2 - 1} \quad \text{and} \quad y = t + \ln t$$


for $1 \leq t \leq 3$. All lengths are measured in centimetres. The region between the curve C and the y-axis is rotated through 2π radians about the y-axis to form a solid.

a) Find the maximum diameter of the ornament, giving your answer to the nearest mm.
b) Find the maximum height of the ornament, giving your answer to the nearest mm.
c) Find the volume of the glass ornament to three significant figures.
d) The ornament is made from glass with density 2.5 g cm^{-3}. Find the mass of the ornament, giving your answer to two significant figures.

2.36 — Question 12. A curve is defined by parametric equations

$$x = \frac{2}{t} \quad \text{and} \quad y = \frac{t^2}{e^t}$$

for $t > 0$. The maximum of the curve is at the point P with $x = a$ where a is a constant to be found.



A region R is enclosed by the curve, the line $x = a$ and $x = 3$.

a) Find the coordinates of P.


b) Show that the volume of the solid generated when R is rotated about the x-axis by 2π radians is given by

$$\pi\left(\frac{29}{18e^{4/3}} - \frac{13}{2e^4}\right).$$

2.36 — Question 13. A curve is defined by parametric equations

$$x = \ln(t-3) \quad \text{and} \quad y = \frac{8}{\sqrt{t}}$$

for $t \geq 3$. The point P has x-coordinate of $\ln 13$.




The shaded region is enclosed by the curve, the y-axis and the line which passes through the point P and the origin. Determine the exact volume of the solid generated when the shaded region is rotated about the x-axis by 2π radians.

2.37 — Question 1. Two points P and Q lie on the curve given. In each case the x-coordinate of P and Q are given along with the curve. Find the exact length of the arc PQ.

a) $x_P = 8$, $x_Q = 35$, $y = \frac{2}{3}x^{3/2}$

b) $x_P = 1$, $x_Q = 4$, $y = \frac{4}{3}x^{3/2}$

2.37 — Question 2. The diagram below shows the curve with equation $y = \cosh x$.



The points A and B have x-coordinates 2 and 3 respectively. Show that the arc length of the curve from A to B is equal to $\frac{7}{12}$.

2.37 — Question 3.

a) Show that $\int \sec x \, dx = \ln|\sec x + \tan x|$.


A curve is given for $\frac{\pi}{4} \leq x \leq \frac{\pi}{3}$ by the equation $y = \ln(\cos x)$.

b) Find the length of the arc on this curve, s, giving your answer in the form $\ln\left(\frac{p}{q}\right)$ where p and q are exact irrational numbers.

2.37 — Question 4. A curve is given by the equation $y = \frac{1}{2}x^2$. Find the exact length of the arc on the curve, s, between the origin and the point where $x = 2$.

2.37 — Question 5. A curve is given by the equation $x = 2\cosh\left(\frac{y}{2}\right)$. Find the exact length of the arc on the curve, s, between the points where $y = 0$ and $y = \ln 4$.

2.37 — Question 6. Part of a suspension bridge is shown in the diagram below, where P and Q are directly beneath A and C respectively.




It is given that $PO = OQ = 70$ metres, and $AP = CQ = 40$ metres. Point B is 15 metres above O. The cable that runs from A to B to C can be modelled by the equation $y = k\cosh\frac{x}{k} - p$, where k and p are constants.

a) Show that $k - p = 15$.
b) Use part (a) to verify that $k = 101.91$ to two decimal places.
c) Find the length of the cable, from A to C, to the nearest metre.

2.37 — Question 7. Two masts, A and B, are built 500 metres apart on horizontal ground. A wire is hung between the masts so that its shape can be modelled by the equation


$$y = p + \cosh\left(\frac{x}{q} - r\right)$$

where y is the height in metres of the wire above the ground, x is the distance in metres from mast A, and p, q and r are positive constants. The diagram below shows the model as being hung from the very top of the mast, which is at a height of 80 metres, and that the minimum distance between the ground and the wire is 60 metres.



a) Find the value of p, and hence find q and r to three significant figures.
b) Find the length of the wire hung between A and B, giving your answer to the nearest metre.

2.37 — Question 8. Two poles AB and CD are placed in the ground, 3 metres apart from each other, as shown in the diagram below. The height of the pole AB is 1.5 metres.



A rope is attached to B and D, the tops of the two poles, and hangs so that it touches the ground at point P, which is 1.6 metres from A. The curve is modelled using the equation

$$y = p\cosh(qx - r) - 2$$

where p, q and r are positive constants.
a) Find the value of p, and hence find q and r to four significant figures.
b) Find the height of pole CD, giving your answer to the nearest centimetre.
c) Find the length of the rope between B and D, giving your answer to the nearest centimetre.

2.37 — Question 9. For each of the following, find the length of the arc, s, on the given curve between the origin and the point at $x = 1$, giving your answer in terms of the constant, $k > 0$:

a) $y = \frac{1}{k}\cosh^2(kx)$

b) $y = \frac{1}{k}\sinh^2(kx)$

2.37 — Question 10.

a) By using the substitution $x = \frac{1}{6}\sinh u$, show that

$$\int \sqrt{1+36x^2}\,dx = \frac{1}{12}\operatorname{arsinh}(6x) + \frac{1}{2}x\sqrt{1+36x^2} + c.$$

Hence calculate the exact length of the arc on the curve $y = 3x^2$ between the points $x = 0$ and $x = \frac{1}{3}$.

2.37 — Question 11.

a) By using substitution $x = \sinh^2 u$ show that

$$\int \sqrt{\frac{x+1}{x}}\, dx = \sqrt{x(x+1)} + \operatorname{arsinh} \sqrt{x} + c.$$

Hence calculate the exact length of the arc, s, on the curve $y = \frac{2}{3}\sqrt{x}$ between the points $x = 1$ and $x = 4$, giving your answer in logarithmic form.

2.37 — Question 12. A curve is given by the equation $y = 5 - \ln(1 - x^2)$ for $x \geq 0$.

a) Show that the exact length of the arc on the curve y is given by

$$s = \int \left(\frac{1+x^2}{1-x^2}\right) dx.$$

b) Hence the exact length of the arc on the curve between the points where $x = 0$ and $x = \frac{1}{2}$.

2.37 — Question 13. A landscape gardener is designing a curved path which will be exactly 39 metres long. The edge of the path follows the curve $y = \frac{1}{3}x^{3/2}$ where x and y are measured in metres. By considering the origin as the start point of the curve, find the value of x which gives a length of 39 metres.

2.37 — Question 14. Engineers are designing the shape of a suspension bridge cable. The cable hangs between two supports and its height above the road can be modelled by the equation

$$h = a + \cosh\left(\frac{x}{b} - c\right)$$

where h is the height of the cable above the road (in metres), x is the horizontal distance from one end of the bridge (in metres), and a, b and c are constants. The bridge span is 60 m long (horizontally), so $0 \leq x \leq 60$. The lowest point of the cable occurs at $x = 30$, where the height is 12 metres. The cable height at each end of the bridge is 20 metres.

a) Find the values of a, b and c giving your answer in exact form.

b) Hence, using your calculator, find the length of the cable used giving your answer to the nearest metre.

2.37 — Question 15. A curve for $t \geq 0$ has parametric equations

$$x = t^2, \quad y = \frac{1}{3}t^3 - t.$$

a) Show that $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (t^2+1)^2$.

b) Hence find the exact length of the curve between the points that occur when $t = 0$ and $t = 4$.

2.37 — Question 16. A curve is defined by the parametric equations

$$x = \frac{2}{3}t^3 + 2, \quad y = 2\sqrt{2}\,t^2 + 1$$

for $0 \leq t \leq 2$.

a) Show that the arc length between $t = 0$ and $t = 2$ is given by $s = \int_0^2 2t\sqrt{t^2 + k}\,dt$ for a constant value k.

b) Hence find the exact length of the arc of the curve between the points where $t = 0$ and $t = 2$.

2.37 — Question 17. A curve is defined by the parametric equations

$$x = 5(1+t)^{1/2}, \quad y = 5(1-t)^{1/2}$$

for $0 \leq t \leq 1$. Find the exact length of the arc of the curve between the points $(5, 5)$ and $(5\sqrt{2}, 0)$.

2.37 — Question 18. A circle has parametric equations

$$x = r\cos\theta \quad \text{and} \quad y = r\sin\theta.$$

By finding the arc length of the curve from $\theta = 0$ to $\theta = 2\pi$, show that the length of the circumference is $2\pi r$.

2.37 — Question 19. A curve is defined by the parametric equations

$$x = 2 + e^{-t}(\cos t + \sin t), \quad y = 1 + e^{-t}(\cos t - \sin t)$$

for $t \geq 0$. Find the exact length of the arc of the curve between the points where $t = 0$ and $t = 3$.

2.37 — Question 20. A curve is defined by the parametric equations

$$x = 2 + 3\cos t, \quad y = 3t + 3\sin t$$

for $t \geq 0$.

a) Show that $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 36\cos^2\left(\frac{t}{2}\right)$.

b) Hence find the exact length of the curve between the points that occur when $t = 0$ and $t = \pi$.

2.37 — Question 21. A curve is defined by the parametric equations

$$x = 3\sin^3 t, \quad y = 3\cos^3 t$$

for $0 \leq t < 2\pi$. Find the exact length of the arc of the curve between the points where $t = 0$ and $t = \frac{\pi}{2}$.

2.37 — Question 22. A curve is defined by the parametric equations

$$x = 4\sin t - \sin 4t, \quad y = 4\cos t - \cos 4t$$

for $t \geq 0$.

a) Show that $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 64\sin^2\left(\frac{3t}{2}\right)$.

b) Hence find the exact length of the curve between the points that occur when $t = 0$ and $t = \frac{\pi}{3}$.

2.37 — Question 23. A designer models the shape of a curved skateboard ramp using parametric equations

$$x = k(t + \sin t), \quad y = k(1 - \cos t)$$

where x and y are measured in metres, $t \geq 0$ and k is a positive constant. The ramp section runs from the ground at $t = 0$ to the platform at $t = \pi$. Given the exact length of the ramp between the ground and the ramp is 12 metres, find the value of k. **Prerequisite Knowledge for Question 24: Calculus of Hyperbolic Functions**

2.37 — Question 24.

a) By using the substitution $u = \sinh x$, show that $\displaystyle\int \operatorname{sech} x \, dx = \arctan(\sinh x) + c$.

A curve is defined by the parametric equations

$$x = \operatorname{sech} t, \quad y = \tanh t$$

for $t \geq 0$.

b) Find the exact length of the arc of the curve between the points where $t = 0$ and $t = \ln 2$.

2.37 — Question 25.

a) Show that

$$\int \sqrt{1+x^2}\,dx = \frac{1}{2}x\sqrt{x^2+1} + \frac{1}{2}\ln\left(x+\sqrt{x^2+1}\right) + c.$$

A curve is defined by the parametric equations

$$x = at^2, \quad y = 4at$$

for $t \geq 0$ where a is a positive constant.

b) Find the exact length of the arc of the curve between the points where $t = 0$ and $t = 1$ giving your answer in terms of a.

2.38 — Question 1. In each case, the x-coordinate of two points P and Q and the curve on which they lie are given. The arc PQ is completely rotated about the x-axis. Determine the exact value of the surface area generated.

a) $x_P = 0, x_Q = 2, y = 3x + 5$

b) $x_P = 0, x_Q = 3, y = 2\sqrt{x}$

c) $x_P = 0, x_Q = 1, y = \cosh x$

d) $x_P = 1, x_Q = 4, y = \frac{1}{2\sqrt{3}}\sqrt{x}(4-x)$

2.38 — Question 2. A line is given by the equation $y = ax$ for a positive constant a, between points $x = 0$ and $x = k$ for positive constant k. Use integration to determine the surface area generated if the arc is rotated 2π radians about the x-axis.

2.38 — Question 3. A small decorative bowl is modelled by

$$y^2 = x + 16$$

for $0 \leq x \leq 6$, where x and y are measured in centimetres. The section of the curve between $x = 0$ and $x = 6$ is rotated completely about the x-axis to form the curved surface of the bowl.

a) Find the surface area of the bowl giving your answer to four significant figures.

b) The cost of one coating of paint for the bowl is £0.03 per cm^2. Find the cost of coating 20 bowls to the nearest £1

2.38 — Question 4. A curve is given by equation $y = \sqrt{x+1}$
An arc on the curve lies between the points where $x = 0$ and $x = k$ and is rotated completely about the x-axis. The surface area generated is given by $\frac{\pi}{6}(13\sqrt{13} - 5\sqrt{5})$ units^2. Determine the value of k

2.38 — Question 5. A curve is given by equation $y = \sqrt{k - x^2}$ for positive constant k between points P and Q which lie on the x-axis and y-axis respectively. By finding the surface area generated when the arc is rotated 2π radians about the x-axis, find the surface area of a sphere with radius $\sqrt{k}$ giving your answer in terms of k

2.38 — Question 6.

a) By using the substitution $x = \sinh t$, show that

$$\int \sqrt{1+x^2}\,dx = \frac{1}{2}x\sqrt{x^2+1} + \frac{1}{2}\ln(x+\sqrt{x^2+1}) + c$$

b) By using the substitution $u = 2e^{2x}$, show that the area of the surface generated when the curve $y = e^{2x}$, for $0 \leq x \leq \ln 2$, is rotated about the x-axis is 48.2 units^2, correct to three significant figures.

2.38 — Question 7. A curve is given by equation

$$y = \frac{1}{3}\cos(3x)$$

An arc on the curve lies between the points where $x = 0$ and $x = \frac{\pi}{6}$ is rotated through 2π radians about the x-axis to form surface area S

a) Show that S is given by

$$S = \frac{2\pi}{3}\int_0^{\pi/6} \cos(3x)\sqrt{1+\sin^2(3x)}\,dx$$

b) Use the substitution $u = \sin(3x)$ to find the exact value of S

2.38 — Question 8. In each case, the x-coordinate of two points P and Q and the curve on which they lie are given. The arc PQ is completely rotated about the y-axis. Determine the exact value of the surface area generated.

a) $x_P = 2, x_Q = 3, y = \frac{\sqrt{7}}{3}(x+1)$
b) $x_P = 1, x_Q = 2, y = x^2$
c) $x_P = 0, x_Q = 5, y = \frac{2}{3}(x+3)^{3/2}$
d) $x_P = 0, x_Q = \ln 2, y = \cosh x$

2.38 — Question 9. A line is given by the equation $y = ax$ for a positive constant a, between points $x = 0$ and $x = k$ for positive constant k. Use integration to determine the surface area generated if the arc is rotated 2π radians about the y-axis.

2.38 — Question 10. A curve is given by the equation $y = \operatorname{arcosh} x$
An arc on the curve, s, lies between the points where $x = 1$ and $x = 2$ and is rotated completely about the y-axis. Show that the area of the surface generated is given by $(2\pi\sqrt{3} + \pi \ln(2 + \sqrt{3}))$ units^2

Parametric Curves

2.38 — Question 11. Find the exact area of the surface generated by rotating the following arcs 2π radians about the x-axis:

a) $x = 6t, y = 2t$ for $0 \leq t \leq 3$
b) $x = t^2, y = \sqrt{4t^2 + 1}$ for $0 \leq t \leq 1$

2.38 — Question 12. Find the exact area of the surface generated by rotating the following arcs 2π radians about the y-axis:

a) $x = 4t^2, y = t^2$ for $1 \leq t \leq 2$

b) $x = \sin^2 t, y = \cos^2 t$ for $0 \leq t \leq \frac{\pi}{2}$

2.38 — Question 13. A curve is defined by the parametric equations

$$x = r\cos\theta, y = r\sin\theta$$

where r is a positive constant and $0 \leq \theta \leq \pi$

By rotating the curve about the x-axis, use integration to show that the surface area of a sphere is $4\pi r^2$

2.38 — Question 14. A curve is defined by the parametric equations

$$x = \operatorname{sech} t, y = \tanh t$$

for $t \geq 0$
The curve is rotated 2π radians about the x-axis between the points $t = 0$ and $t = \ln 5$

a) Show that $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \operatorname{sech}^2 t$

b) Find the exact area of the surface area generated.

2.38 — Question 15. The shape of a funnel may be modelled using parametric equations

$$x = 4(1+t)^{3/2}, y = 4(1-t)^{3/2}$$

where x and y are measured in centimetres for $0 \leq t \leq 0.8$ which is rotated 2π radians about the x-axis.

a) Determine the surface area of the funnel giving your answer to three significant figures.

The manufacturer wants to add a single layer of protective coating to the funnels. The cost of one layer of protective coating for the funnel is £0.02 per cm.

b) Find the cost of covering 15 funnels with one layer to the nearest £1

2.38 — Question 16. A curve is defined by the parametric equations

$$x = k\cos^3 t, y = k\sin^3 t$$

where k is a positive constant r and $\frac{\pi}{4} \leq t \leq \frac{\pi}{2}$
The curve is rotated 2π radians about the x-axis. Use integration to show that the surface area formed is $\frac{3\pi k^2}{20}(8-\sqrt{2})$ units^2.

2.38 — Question 17. A curve is defined by the parametric equations

$$x = 2t, y = \frac{1}{2}t^2 - \ln t$$

for $1 \leq t \leq 2$
The curve is rotated 2π radians about the y-axis. Determine the exact area of the surface area generated.

2.38 — Question 18.

a) Use integration by parts to show that

$$\int \frac{\ln x}{x}\, dx = \frac{1}{2}\ln^2 x + c$$

A curve is defined by the parametric equations

$$x = 2t, y = \frac{1}{2}t^2 - \ln t$$

for $1 \leq t \leq 3$. The curve is rotated 2π radians about the x-axis.

b) Find the area of the surface area generated giving your answer to three significant figures.

2.38 — Question 19. A curve is defined by the parametric equations

$$x = e^{\theta}\sin\theta, y = e^{\theta}\cos\theta$$

for $0 \leq \theta \leq \frac{\pi}{2}$. The curve is rotated 2π radians about the x-axis. Use integration to show that the surface area formed is $\frac{2\pi\sqrt{2}}{5}(e^{\pi} - 2)$ units^2.

2.38 — Question 20.

a) Show that $\int e^{-2x}(\cos x - \sin x)\,dx = -\frac{1}{5}e^{-2x}(\cos x - 3\sin x) + c$

A curve is defined by the parametric equations

$$x = 4 + e^{-t}(\cos t + \sin t), \qquad y = e^{-t}(\cos t - \sin t)$$

for $0 \leq t \leq \frac{\pi}{2}$. The curve is rotated 2π radians about the x-axis.

b) Find the area of the surface area formed giving your answer in exact form.