Pure Mathematics Year 2

Chapter 10: Numerical Methods

1. Locating roots
2. Using iteration to approximate roots of $f(x) = 0$
3. Introduction to the Newton–Raphson method

At GCSE you used iteration to obtain successively better approximations to solutions. We revisit this and introduce a more powerful method.


1. Locating roots

Understand what a root is and when a root must lie in an interval.

Show that $f(x) = x^3 - 4x^2 + 3x + 1$ has a root between $x = 1.4$ and $x = 1.5$.

2. Iteration

Rearrange $f(x) = 0$ as $x = g(x)$, then use $x_{n+1} = g(x_n)$.



3. Newton–Raphson

Follow tangents to approach a root, often faster:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

Finding a root of $f(x)$ means finding an input whose output is zero.


Solve $f(x) = 0$.

Some exact roots are complicated and difficult to calculate. For example, $x^3 + 2x^2 - 3x + 4 = 0$:

$$x = \frac{1}{3}\left(-2 - \frac{13}{\sqrt[3]{89 - 6\sqrt{159}}} - \sqrt[3]{89 - 6\sqrt{159}}\right).$$


Other equations, such as $x - \cos x = 0$, have no standard exact algebraic formula for their root.

Use a numerical approximation: the root is approximately 0.739085.



Numerical methods give progressively better approximations to solutions of $f(x) = 0$.

Show that $f(x) = e^x + 2x - 3$ has a root between $x = 0.5$ and $x = 0.6$.



If a continuous graph changes from negative to positive, its y-values must pass through zero.

Root interval


Exam tip

Calculate both endpoint values and explicitly state that there is a change of sign. Include continuity in the argument.

The Function Must Be Continuous

Luisa · 5

A student claims: "For $f(x) = 1/x$, $f(-1) = -1$ and $f(1) = 1$. There is a change of sign, so a root lies in $[-1, 1]$."




Why is this claim wrong?

The function is not continuous across this interval: it is undefined at $x = 0$, where there is a vertical asymptote. It never equals zero.

A sign change guarantees a root only when the function is continuous throughout the interval. Otherwise the graph can pass from negative to positive without crossing the axis.

No Sign Change Does Not Mean No Root

Luisa · 6



No sign change does **not** prove that an interval contains no roots.

The graph can cross the axis twice and return to the original sign. A graph can also touch the axis at a root without changing sign.

The sign-change test can therefore miss roots: it gives a sufficient condition when continuity holds, not a necessary condition.

Proving a Solution to a Given Accuracy — Luisa · 7

Edexcel C3 January 2013

$g(x) = e^{x-1} + x - 6$. Its root is α.

c. By choosing a suitable interval, show that $\alpha = 2.307$ correct to 3 decimal places.

What are the rounding boundaries?

Lower boundary

Upper boundary

A root strictly between these boundaries rounds to 2.307 to 3 decimal places.

c

Example — Logarithm and Reciprocal — Luisa · 8

a. Sketch $y = \ln x$ and $y = 1/x$ on the same axes. Explain why $f(x) = \ln x - 1/x$ has only one root.

b. Show that the root lies in $1.7 < x < 1.8$.

c. Show that $\alpha = 1.763$ to 3 decimal places.

a sketch

a

b

c

2. $f(x) = 3 + x^2 - x^3$.

a. Show that a root α lies in $[1.8, 1.9]$.

b. Use a sign change in a suitable interval to verify $\alpha = 1.864$ to 3 decimal places.

2a

2b

3. $h(x) = \sqrt[3]{x} - \cos x - 1$. Use radians.

a. Show that a root α lies in $[1.4, 1.5]$.

b. Use a sign change in a suitable interval to verify $\alpha = 1.441$ to 3 decimal places.

3a

3b

4. $f(x) = \sin x - \ln x$. Use radians. The domain is $x > 0$.

a. Show that a root α lies in $[2.2, 2.3]$.

b. Use a sign change in a suitable interval to verify $\alpha = 2.219$ to 3 decimal places.

4a

4b

Using Iteration to Approximate a Root **Luisa · 12**

Edexcel C3 January 2013

$g(x) = e^{x-1} + x - 6$.

a. Show that $g(x) = 0$ can be rearranged as $x = \ln(6 - x) + 1$.

b. Use $x_{n+1} = \ln(6 - x_n) + 1$, with $x_0 = 2$, to find x_1, x_2, x_3 to 4 decimal places.

Iteration rule

Rearrange $f(x) = 0$ into $x = g(x)$, then use $x_{n+1} = g(x_n)$. We will see why this works shortly.

a

Tip

There may be several occurrences of x to isolate. Use the target equation to guide the rearrangement.

Iteration — Calculation and Exam Technique

Luisa · 13

Edexcel C3 January 2013

$g(x) = e^{x-1} + x - 6$.

a. Show that $g(x) = 0$ can be rearranged as $x = \ln(6 - x) + 1$.

b. Use $x_{n+1} = \ln(6 - x_n) + 1$, with $x_0 = 2$, to find x_1, x_2, x_3 to 4 decimal places.

b

x_0 is the starting value. The later terms are successive approximations to a root.

Calculator tip

Enter 2 first, then enter $\ln(6 - \text{Ans}) + 1$. Press equals repeatedly for successive iterations. Retain unrounded values during the calculation.

Exam tip

Show the substitution for x_1 to earn the method mark. Then state the required values of x_2 and later terms.

Does the Starting Value Matter?

Luisa · 14

The starting value matters: iteration can approach a different root, or fail to converge. Staircase and cobweb diagrams help explain why.

$f(x) = x^3 - 3x^2 - 2x + 5$.

a. Show there is a root in $3 < x < 4$.

b. Use $x_{n+1} = \sqrt{(x_n^3 - 2x_n + 5)/3}$ to find x_1, x_2, x_3, to 4 d.p., starting from i. $x_0 = 1.5$; ii. $x_0 = 4$.

a

b(ii)

b(i)

Test Your Understanding — Iteration

Edexcel C3 June 2012, Q2

$f(x) = x^3 + 3x^2 + 4x - 12$. The equation $f(x) = 0$ has a single root between 1 and 2.

a. Show that $f(x) = 0$ can be written as $x = \sqrt{4(3-x)/(3+x)}$ for the root between 1 and 2.

b. Use $x_{n+1} = \sqrt{4(3-x_n)/(3+x_n)}$, $x_0 = 1$, to find x_1, x_2, x_3 to 2 d.p.

c. Prove that the root $\alpha = 1.272$ to 3 d.p.

a

b


c

Exam tip

Mark schemes show the minimum acceptable wording. State the endpoint values, change of sign and continuity clearly.

Why Does Iteration Work?

Solve $x^2 - x - 1 = 0$. For the positive root, rearrange as $x = \sqrt{x+1}$ and use $x_{n+1} = \sqrt{x_n + 1}$. Why does this work?



The solution of $x = \sqrt{x+1}$ is the intersection of $y = x$ and $y = \sqrt{x+1}$. Its x-coordinate, α, is a root of the original equation.


With $x_0 = 1$, find $\sqrt{x_0 + 1}$: move vertically to the curve to obtain the y-value.

This gives $x_1 = \sqrt{2} = 1.414\ldots$. Feed it back as the next input: move horizontally to $y = x$, so the y-value becomes the new x-value.

Repeat using $x_1 = 1.414\ldots$ to obtain x_2, and so

The staircase shape gives the name: a staircase diagram. The iterates converge towards α.


Another rearrangement of $x^2 - x - 1 = 0$ is $x = 1/(x-1)$. Use $x_{n+1} = 1/(x_n - 1)$. What happens?



Start with a negative value, as in this diagram. Alternate between the curve $y = 1/(x-1)$ and the line $y = x$.

The terms alternate around α and move closer to it. This is a cobweb diagram. Here iteration converges to the negative root, $\alpha = (1 - \sqrt{5})/2$.

We can also rearrange $x^2 - x - 1 = 0$ as $x = x^2 - 1$, giving $x_{n+1} = x_n^2 - 1$. What happens now?



Start to the right of the positive root. Each iteration moves farther from α, instead of closer. The approximations diverge, so this attempt fails.

Test Your Understanding — Staircase Diagram

Luisa · 19

$f(x) = x^2 - 8x + 4$.

a. Show that a nonnegative root can be written as $x = \sqrt{8x - 4}$.


b. Starting with $x_0 = 1$, use $x_{n+1} = \sqrt{8x_n - 4}$ to draw a staircase diagram. Label x_0, x_1, x_2 and the root α.

a–b

b sketch


Fixed-Point Iteration I

Luisa · 20




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

1. Mark on $y = g(x)$ the point whose x-coordinate is 4.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

2. Draw a horizontal line from this point until it reaches $y = x$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

3. Draw a vertical line from this point until it reaches the curve $y = g(x)$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

4. Draw a horizontal line from this point until it reaches $y = x$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

5. Repeat steps 3 and 4 several times. What do you notice?




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

1. Mark on $y = g(x)$ the point whose x-coordinate is 4.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

2. Draw a horizontal line from this point until it reaches $y = x$.



The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

3. Draw a vertical line from this point until it reaches the curve $y = g(x)$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

4. Draw a horizontal line from this point until it reaches $y = x$.

Fixed-Point Iteration II

Luisa · 29




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

5. Repeat steps 3 and 4 several times. What do you notice?


Fixed-Point Iteration III

Luisa · 30




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

1. Mark on $y = g(x)$ the point whose x-coordinate is 2.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

2. Draw a horizontal line from this point until it reaches $y = x$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

3. Draw a vertical line from this point until it reaches the curve $y = g(x)$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

4. Draw a horizontal line from this point until it reaches $y = x$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

5. Repeat steps 3 and 4 several times. What do you notice?




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

1. Mark on $y = g(x)$ the point whose x-coordinate is 3.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

2. Draw a horizontal line from this point until it reaches $y = x$.




The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

3. Draw a vertical line from this point until it reaches the curve $y = g(x)$.



The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

4. Draw a horizontal line from this point until it reaches $y = x$.



The graph shows $y = x$ and $y = g(x)$. The line and curve meet at a fixed point whose x-coordinate is close to 3.

5. Repeat steps 3 and 4 several times. What do you notice?

Describe the construction and its behaviour near the fixed point close to 3.

Diagram

Conclusion

Answer — Fixed-Point Iteration II

Luisa · 41

Describe the construction and its behaviour near the fixed point close to 3.

Diagram

Conclusion

Answer — Fixed-Point Iteration III

Luisa · 42

Describe the construction and its behaviour near the fixed point close to 3.

Diagram

Conclusion

Answer — Fixed-Point Iteration IV

Luisa · 43

Describe the construction and its behaviour near the fixed point close to 3.

Diagram

Conclusion

Exercise 10B — Pages 280–282

Luisa · 44

3. $f(x) = x^2 - 6x + 1$.

a. Show that $f(x) = 0$ can be written as $x = \sqrt{6x - 1}$.

b. Sketch $y = x$ and $y = \sqrt{6x - 1}$.

c. State the number of roots.

d. Explain graphically why $x_{n+1} = \sqrt{6x_n - 1}$, $x_0 = 2$, converges to a root.

e. By sketching a diagram, explain why $x_{n+1} = (x_n^2 + 1)/6$, $x_0 = 10$, moves away from the roots.

3a

3b

3c

3d

3e

3e conclusion

6. $f(x) = x^4 - 3x^3 - 6$.

a. Rearrange $f(x) = 0$ as $x = \sqrt[3]{px^4 + q}$, finding p, q.

b. With $x_0 = 0$, use $x_{n+1} = \sqrt[3]{px_n^4 + q}$ to find x_1, x_2, x_3 to 3 d.p.

c. Prove that the negative root $\alpha = -1.132$ to 3 d.p.

6a

6b

6c

Pure Mathematics Year 2 · Chapter 10

Numerical methods

Introduction to the Newton–Raphson method

Name the movie.




Solving a Cubic Numerically

Use the Newton–Raphson method to solve
$f(x) = x^3 + x^2 - 5x + 2 = 0$.


Three Roots of the Cubic

$x^3 + x^2 - 5x + 2 = 0$




$x^3 + x^2 - 5x + 2 = 0$

Zoom in on the right-hand root. Start with $x_0 = 2$.




$x^3 + x^2 - 5x + 2 = 0$

Zoom in on the right-hand root. Start with $x_0 = 2$.




$x^3 + x^2 - 5x + 2 = 0$

Zoom in on the right-hand root. Start with $x_0 = 2$.



$x^3 + x^2 - 5x + 2 = 0$

Zoom in on the right-hand root. Start with $x_0 = 2$.




Zoom in on One of the Roots

Luisa · 53

$x^3 + x^2 - 5x + 2 = 0$

Zoom in on the right-hand root. Start with $x_0 = 2$.




What Happens If We Start from the Left?

Luisa · 54


$f(x) = x^3 + x^2 - 5x + 2$

Try a starting point to the left of the right-hand root.




$f(x) = x^3 + x^2 - 5x + 2$

Try a starting point to the left of the right-hand root.




$f(x) = x^3 + x^2 - 5x + 2$

Try a starting point near the middle root.




$f(x) = x^3 + x^2 - 5x + 2$

Try a starting point near the middle root.




Example — solve $f(x) = 0$.

Set x_0 and look at the point $(x_0, f(x_0))$.



Example — solve $f(x) = 0$.


Set x_0 and look at the point $(x_0, f(x_0))$.



Example — solve $f(x) = 0$.

Set x_0 and look at the point $(x_0, f(x_0))$.

x_1 is where the tangent crosses the x-axis.




Example — solve $f(x) = 0$.

Set x_0 and look at the point $(x_0, f(x_0))$.

x_1 is where the tangent crosses the x-axis.

Gradient of tangent $= \dfrac{f(x_0)}{x_0 - x_1}$.




Example — solve $f(x) = 0$.

Set x_0 and look at the point $(x_0, f(x_0))$.

x_1 is where the tangent crosses the x-axis.

Gradient of tangent $= \dfrac{f(x_0)}{x_0 - x_1}$.

$f'(x_0) = \dfrac{f(x_0)}{x_0 - x_1}$.



Example — solve $f(x) = 0$.


Set x_0 and look at the point $(x_0, f(x_0))$.

x_1 is where the tangent crosses the x-axis.

Gradient of tangent $= \dfrac{f(x_0)}{x_0 - x_1}$.

$f'(x_0) = \dfrac{f(x_0)}{x_0 - x_1}$.

$f(x_0) = (x_0 - x_1)f'(x_0)$.



Example — solve $f(x) = 0$.

Set x_0 and look at the point $(x_0, f(x_0))$.


x_1 is where the tangent crosses the x-axis.

Gradient of tangent $= \dfrac{f(x_0)}{x_0 - x_1}$.

$f'(x_0) = \dfrac{f(x_0)}{x_0 - x_1}$.

$f(x_0) = (x_0 - x_1)f'(x_0)$.

$x_1 = x_0 - \dfrac{f(x_0)}{f'(x_0)}$.



Repeat the process to obtain the iterative formula.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

The function and its derivative must be defined, with $f'(x_n) \neq 0$.

Repeat the process to obtain the iterative formula.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

Calculator hint

Save $f(x)$ as Y1 in Graph mode. Use the Vars key to access Y1 in Run mode.

The iterative formula is

$$\text{Ans} - \frac{\text{Y1(Ans)}}{\left.\frac{d}{dx}\text{Y1}(x)\right|_{x=\text{Ans}}}.$$

Use the Newton–Raphson method to solve
$f(x) = x^3 + x^2 - 5x + 2 = 0$.

Use the Newton–Raphson method to solve
$f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

Use the Newton–Raphson method to solve
$f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

Use the Newton–Raphson method to solve
$f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Applying Newton–Raphson to the Cubic

Luisa · 71

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Applying Newton–Raphson to the Cubic

Luisa · 72

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Calculator hint

Enter the recurrence using Ans in place of x_n, then press equals repeatedly.

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Calculator hint

Enter the recurrence using Ans in place of x_n, then press equals repeatedly.

$$x_1 = \frac{18}{11} = 1.636363636\ldots$$

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Calculator hint

Enter the recurrence using Ans in place of x_n, then press equals repeatedly.

$$x_1 = \frac{18}{11} = 1.636363636\ldots$$

$$x_2 = 1.497200048\ldots$$

Applying Newton–Raphson to the Cubic — Luisa · 75

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Calculator hint

Enter the recurrence using Ans in place of x_n, then press equals repeatedly.

$$x_1 = \frac{18}{11} = 1.636363636\ldots$$

$$x_2 = 1.497200048\ldots$$

Applying Newton–Raphson to the Cubic — Luisa · 76

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Calculator hint

Enter the recurrence using Ans in place of x_n, then press equals repeatedly.

$$x_1 = \frac{18}{11} = 1.636363636\ldots$$

$$x_2 = 1.497200048\ldots$$

$$x_3 = 1.473521722\ldots$$

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Calculator hint

Enter the recurrence using Ans in place of x_n, then press equals repeatedly.

$$x_1 = \frac{18}{11} = 1.636363636\ldots$$

$$x_2 = 1.497200048\ldots$$

$$x_3 = 1.473521722\ldots$$

$$x_4 = 1.472834484\ldots, \quad x_5 = 1.472833909\ldots$$

Applying Newton–Raphson to the Cubic

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$f'(x) = 3x^2 + 2x - 5$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Using $x_0 = 2$.

Calculator hint

Enter the recurrence using Ans in place of x_n, then press equals repeatedly.

$$x_1 = \frac{18}{11} = 1.636363636\ldots$$

$$x_2 = 1.497200048\ldots$$

$$x_3 = 1.473521722\ldots$$

$$x_4 = 1.472834484\ldots, \quad x_5 = 1.472833909\ldots$$

This suggests that one root is 1.4728, correct to 4 decimal places. We still need to verify the rounding.

Verify the Root to Four Decimal Places — Luisa · 79

Show that the root is 1.4728, correct to 4 decimal places.

$f(x) = x^3 + x^2 - 5x + 2$.

Verify the Root to Four Decimal Places — Luisa · 80

Show that the root is 1.4728, correct to 4 decimal places.

$f(x) = x^3 + x^2 - 5x + 2$.

The rounding boundaries are 1.47275 and 1.47285.

Verify the Root to Four Decimal Places

Luisa · 81

Show that the root is 1.4728, correct to 4 decimal places.

$f(x) = x^3 + x^2 - 5x + 2.$

The rounding boundaries are 1.47275 and 1.47285.

$f(1.47275) = 1.47275^3 + 1.47275^2 - 5(1.47275) + 2$
$= -3.73641078\ldots \times 10^{-4} < 0.$

Verify the Root to Four Decimal Places

Luisa · 82

Show that the root is 1.4728, correct to 4 decimal places.

$f(x) = x^3 + x^2 - 5x + 2.$

The rounding boundaries are 1.47275 and 1.47285.

$f(1.47275) = 1.47275^3 + 1.47275^2 - 5(1.47275) + 2$
$= -3.73641078\ldots \times 10^{-4} < 0.$

$f(1.47285) = 1.47285^3 + 1.47285^2 - 5(1.47285) + 2$
$= 7.166\ldots \times 10^{-5} > 0.$

Verify the Root to Four Decimal Places

Luisa · 83

Show that the root is 1.4728, correct to 4 decimal places.

$f(x) = x^3 + x^2 - 5x + 2$.

The rounding boundaries are 1.47275 and 1.47285.

$f(1.47275) = 1.47275^3 + 1.47275^2 - 5(1.47275) + 2$
$= -3.73641078\ldots \times 10^{-4} < 0.$

$f(1.47285) = 1.47285^3 + 1.47285^2 - 5(1.47285) + 2$
$= 7.166\ldots \times 10^{-5} > 0.$

There is a change of sign. Since f is continuous, there is a root in $(1.47275, 1.47285)$. It therefore rounds to 1.4728 to 4 decimal places.

Find the Other Roots

Luisa · 84

Use the Newton–Raphson method to solve $f(x) = x^3 + x^2 - 5x + 2 = 0$.

$$x_{n+1} = x_n - \frac{x_n^3 + x_n^2 - 5x_n + 2}{3x_n^2 + 2x_n - 5}.$$

Choose a different starting value to find another root: use $x_0 = 0$.

Spreadsheet setup

Column A: x. Column B: $f(x)$. Column C: $f'(x)$.


Enter 0 in A2. In B2 enter:


```
=A2^3+A2^2-5*A2+2
```

Also known as Newton's method. We wish to find the root α. Start with an approximation x_0, which may be well off the mark.

Follow the direction of the curve using its tangent. If the curve were reasonably straight, the point where the tangent hits the x-axis would be close to the root.

Repeat to obtain, hopefully, increasingly accurate approximations. Can you find a formula for x_{n+1} in terms of x_n?





At $x = x_n$, the tangent has gradient $f'(x_n)$.

Using Year 1 coordinate geometry:

$$y - f(x_n) = f'(x_n)(x - x_n).$$


At the tangent's intersection with the axis, $y = 0$ and $x = x_{n+1}$.

$$-f(x_n) = f'(x_n)(x_{n+1} - x_n).$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

Newton–Raphson — Animated Demonstration

Luisa · 87



Follow each step using Next or Space. The original demonstration starts at x_1.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

$$\boldsymbol{x_{\blacksquare} = x_{\blacksquare} - \frac{f(x_{\blacksquare})}{f'(x_{\blacksquare})}.}$$

Demonstration supplied in the source slides, credited there to Wikipedia; reconstructed as native geometry.

Example — Solving x = cos x

Luisa · 88

Return to $x = \cos x$, with $x_0 = 0.5$. Use radians: the differentiation formula assumes radian measure.

Recurrence

Iterations

Calculator tip

Enter 0.5 and press equals. Then enter

$$\text{Ans} - \frac{\text{Ans} - \cos(\text{Ans})}{1 + \sin(\text{Ans})}$$

and press equals repeatedly.

Quickfire Questions — Luisa · 89

Using Newton–Raphson, state the recurrence for each function.

a. $f(x) = x^3 - 2$

a

b. $f(x) = \tan x$. Use radians; $x_n \neq \pi/2 + k\pi$, $k \in \mathbb{Z}$.

b

c. $f(x) = x^2 - x - 1$

c

Example Exam Question — Luisa · 90

Edexcel FP1 June 2013(R), Q3c

$f(x) = \frac{1}{2}x^4 - x^3 + x - 3$. The equation $f(x) = 0$ has a root β in $[-2, -1]$.

c. Starting with -1.5, apply Newton–Raphson once to obtain a second approximation to β. Give your answer to 2 decimal places. (5 marks)

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

c

Test Your Understanding — Luisa · 91

Edexcel FP1 January 2010, Q2c


$f(x) = 3x^2 - 11/x^2$.

c. Starting with 1.4, apply Newton–Raphson once to obtain a second approximation to α. Give your answer to 3 decimal places. (5 marks)


$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

c

When Does Newton–Raphson Fail? — Luisa · 92



If the starting value is a stationary point, $f'(x_0) = 0$. Newton–Raphson divides by zero. For a stationary point away from the axis, the horizontal tangent never reaches the axis.



For $f(x) = x/\sqrt{1+x^2}$, the iterates can oscillate on opposite sides of $\alpha = 0$ and move farther away. Newton–Raphson is an iteration and need not converge.

1. $f(x) = x^3 - 2x - 1$.

a. Show a root α lies in $1 < \alpha < 2$.

b. Starting with $x_0 = 1.5$, apply Newton–Raphson once. Give the second approximation to 3 d.p.

2. $f(x) = x^2 - 4/x + 6x - 10$, $x \neq 0$.

a. Find $f'(x)$. (2 marks)

A root lies in $[-0.4, -0.3]$.

b. Starting with -0.4, apply Newton–Raphson once, giving the second approximation to 3 d.p. (4 marks)

1a

2a


1b

2b

3. $f(x) = x^{3/2} - e^{-x} + 1/\sqrt{x} - 2$, $x > 0$. The diagram shows $y = f(x)$. Point A, with x-coordinate q, is stationary. A root α lies in $[1.2, 1.3]$.

a. Explain why $x_0 = q$ is unsuitable. (1 mark)

b. Starting with $x_0 = 1.2$, apply Newton–Raphson once. Give the second approximation to 3 d.p. (4 marks)



3a

3b

4. $f(x) = 1 - x - \cos(x^2)$. Use radians.

a. Show that a root α lies in $1.4 < \alpha < 1.5$. (1 mark)

b. Starting with $x_0 = 1.4$, apply Newton–Raphson once, giving the second approximation to 3 d.p. (4 marks)

c. Use a sign change in a suitable interval to show that your answer to b is correct to 3 d.p. (2 marks)

4a

4b

4c

5. $f(x) = x^2 - 3/x^2$, $x > 0$.

a. Show that a root α lies in $[1.3, 1.4]$. (1 mark)

b. Find $f'(x)$. (2 marks)

c. Starting with 1.3, apply Newton–Raphson once, giving the second approximation to 3 d.p. (3 marks)

5a

5b

5c

6. $y = f(x) = x^2 \sin x - 2x + 1$. P, Q and R are roots; A and B are stationary points with x-coordinates a, b.

a. Show a root lies in each interval: i. $[0.6, 0.7]$, ii. $[1.2, 1.3]$, iii. $[2.4, 2.5]$.

b. Explain why $x_0 = a$ is unsuitable for Newton–Raphson.

c. Starting with $x_0 = 2.4$, apply Newton–Raphson once. Give the second approximation to 3 d.p.

6a

6b

6c

7. $f(x) = \ln(3x - 4) - x^2 + 10$, $x > 4/3$.

a. Show that a root α lies in $[3.4, 3.5]$.

b. Find $f'(x)$.

c. Starting with 3.4, apply Newton–Raphson once, giving the second approximation to 3 d.p.

7a

7b

7c

Application to Modelling — Luisa · 99

The price of a car, in pounds, x years after purchase is modelled by

$f(x) = 15000(0.85)^x - 1000\sin x, \quad x > 0.$

Use radians.

a. Find its value after 10 years, to the nearest £100.

b. Show that f has a root between 19 and 20.

c. Find $f'(x)$. Starting with 19.5, apply Newton–Raphson once to estimate when the value becomes zero. Give your answer to 3 d.p.

d. Criticise the model as the car gets older.

a

b

c

d

Exercise 10D — Page 287 — Luisa · 100

1. An astronomer models a planet on an elliptical orbit. Its angle E radians is related to the angle M it would have travelled on a circular path by

   $M = E - 0.1\sin E, \quad E \geq 0.$

   She needs E when $M = \pi/6$.

a. Show that E is a root of $f(x) = x - 0.1\sin x - k$, finding k.

b. Starting with 0.6, apply Newton–Raphson once to find a second approximation to E.

c. Use a sign change in a suitable interval to show that the answer to b is correct to 3 d.p.

1a

1b

1c