Parametric Equations

Starter 2

1. Make t the subject:

a. $x = 4t - kt$

b. $y = 3t^2$

c. $y = 2 - 4\ln t$

d. $x = 1 + 2e^{-3t}$

1

2. Express using cosine:

a. $4 + 3\sin^2 x$

b. $\sin 2x$

c. $\cot x$

d. $2\cos x + \cos 2x$

2

3. Find the ranges:

a. $\ln(x+1),\ x > 0$

b. $2\sin x,\ 0 < x < \pi$

c. $x^2 + 4x - 2,\ -4 < x < 1$

d. $1/(2x+5),\ x > -2$

a

4. Find the intersections of the circle with centre $(0, 4)$, radius 5, and the line $2y - x - 10 = 0$.

4

Chapter overview

1. Convert $x = 2\cos t + 1,\ y = 3\sin t$ to Cartesian form.
2. Sketch $x = 2t,\ y = 5/t$.
3. Find where $x = t^2,\ y = 4t$ meets $x + y + 4 = 0$.
4. Model a plane using $x = (v\cos\theta)t,\ y = (v\sin\theta)t$.

A Cartesian equation relates x and y, for example $x^2 - y^2 = 5$.

Parametric equations express both coordinates using another variable, such as $x = \sin t,\ y = t^2$.

Each allowed value of t gives a point on the curve. Eliminating t does not always preserve the full domain or direction.

Converting parametric to Cartesian 5

Find a Cartesian equation and the domain and range of

$x = 2t, \quad y = t^2, \quad -3 < t < 3.$

Answer

Sketch

Further Example 6

Find a Cartesian equation and state its domain and range:

$x = \ln(t + 3), \quad y = 1/(t + 5), \quad t > -2.$

Answer

Now you try. 7

1. Eliminate t: $x = \ln(t+2),\ y = 1/(t+1),\ t > -1$.

1

2. Eliminate t: $x = \ln t,\ y = t^2 - 2,\ t > 0$.

2


Pure Year 2/AS Exercise 8A (200-202) 8

8. A curve is given by $x = 3\sqrt{t},\ y = t^3 - 2t,\ 0 \le t \le 2$.

a. Find its Cartesian equation and domain.

b. Find where $dy/dt = 0$.

c. Hence find the range of y.



a

b

c

9. $x = t^3 - t,\ y = 4 - t^2,\ t \in \mathbb{R}$.

a. Show that $x^2 = (a - y)(b - y)^2$, finding a, b.

b. Find the greatest value of y.

a

b

...when you have trig identities 10

1. Eliminate t from $x = \sin t + 2,\ y = \cos t - 3$.

1

2. Eliminate t from $x = \sin t,\ y = \sin 2t,\ -\pi/2 \leq t \leq \pi/2$.

2

Sketch

Now you try.. 11

1. $x = 2\sin t,\ y = 1 - \cos 2t,\ -\pi/2 \le t \le \pi/2$. Find a Cartesian equation and its domain.

1

2. $x = \cot t + 2,\ y = \csc^2 t - 2,\ 0 < t < \pi$. Find a Cartesian equation and sketch the curve.

2

Sketch

Exercise 8B Pure 2 pages 204-206 12

6. $x = 8\cos t,\ y = \frac{1}{4}\sec^2 t,\ -\pi/2 < t < \pi/2$.

Find a Cartesian equation, its domain and range. The endpoints $t = \pm\pi/2$ are excluded because $\sec t$ is undefined.

Answer

7. $x = 3\cot^2 2t,\ y = 3\sin^2 2t,\ 0 < t \leq \pi/4$. Find a Cartesian equation and its domain.

Answer

8. $x = \frac{1}{3}\sin t,\ y = \sin 3t,\ 0 < t < \pi/2$.

a. Show $y = ax(1 - bx^2)$, finding a, b.

b. State the domain and range.

a

b

10. $x = \tan^2 t + 5,\ y = 5\sin t,\ 0 < t < \pi/2.$

a. Find a Cartesian equation.

b. State the domain and range.

a

b

11. $x = \tan t,\ y = 3\sin(t - \pi),\ 0 < t < \pi/2$. Find a Cartesian equation and its domain and range.

Answer

Sketching a spiral

$x = \theta\cos\theta, \quad y = \theta\sin\theta, \quad 0 \le \theta \le 4\pi.$

Mark the points for $\theta = 0, \pi/2, \pi, 3\pi/2, 2\pi$. What happens during the second turn?


Sketch

Sketch

Plot points in parameter order

θ	0	$\pi/2$	π	$3\pi/2$	2π
x	0	0	$-\pi$	0	2π
y	0	$\pi/2$	0	$-3\pi/2$	0

For $x = \theta\cos\theta,\ y = \theta\sin\theta$, plot further points up to 4π. Join them smoothly and show the direction as θ increases.



Now you try.. 19

Sketch $x = 2t,\ y = t^2,\ -1 \leq t \leq 5$, showing endpoints and direction.

Sketch

Sketch

Exercise IM Level 1 Exercise 1 20

Find Cartesian equations:

i. $x = 1 - t,\ y = t^2 - 4$

ii. $x = 2t^2,\ y = 1/t,\ t \neq 0$

iii. $x = 2\cos\theta + \sin\theta,\ y = \cos\theta - 2\sin\theta$

i

ii

iii

Exercise IM Level 1 Exercise 2 — 21

For $x = t^2,\ y = t^3$, complete a table for integer t from -3 to 3, sketch the curve and find a Cartesian equation.

Table

Equation

Sketch

Exercise 8C (Pages 207-208) — 22

Find the Cartesian equation and sketch $x = 3 - t,\ y = t^2 - 2,\ -2 \leq t \leq 3$.

Answer

Sketch

Exercise 8C (Pages 207-208) 23

$x = 9\cos t - 2,\ y = 9\sin t + 1,\ -\pi/6 \le t \le \pi/2.$

a. Find a Cartesian equation.

b. Sketch the arc, giving its endpoints.

c. Find its length.

a

b

c

Sketch

Points of Intersection 24

$x = at^2 + t,\ y = a(t^3 + 8)$ passes through $(-4, 0)$. Find a and the intersections with the y-axis.

Find a

Intersections

Points of Intersection 25

1. Find where $x = t^2,\ y = 4t$ meets $x + y + 4 = 0$.

1

2. $x = \cos t + \sin t,\ y = (t - \pi/6)^2,\ -\pi/2 < t < 4\pi/3$. Find points with $y = \pi^2$, and intersections with the y-axis.


2: given y

2: axis

Now you try 26

C4 January 2013

$x = 1 - t/2,\ y = 2^t - 1$. Find the intercepts with both axes.



y-axis

x-axis

3. $x = 4at^2,\ y = a(2t - 1)$ passes through $(4, 0)$. Find a.

Answer

4. $x = b(2t - 3),\ y = b(1 - t^2)$ passes through $(0, -5)$. Find b.

Answer

5. Find the intersection of $x = 3t + 2,\ y = 1 - t$ with the line $x + y = 2$.


Answer

12. $x = 6\cos t,\ y = 4\sin 2t + 2,$ $-\pi/2 < t < \pi/2.$

a. Find the x-axis intercepts.

b. Find the parameter values for intersections with $y = 4$.

c. Give the coordinates of these points.



a

b

c

Exercise 8D (page 211-213) Q13 — 31

Show that the line $y = 2x - 5$ does not intersect $x = 2t,\ y = 4t(t-1)$.

Answer

Exercise 8D (page 211-213) Q14 — 32

$x = \sin t,\ y = \cos 2t + 1,\ 0 \leq t \leq 2\pi$.

a. For which values of k does $y = k$ meet the curve?

b. Find a Cartesian equation and the horizontal tangent.

a

b

Sketch

Modelling 33

A plane travels at constant speed v, making angle θ with the horizontal. It rises 120 m for every 600 m horizontally.

a. Find θ.

b. Give parametric equations when $v = 50\,\text{m}\,\text{s}^{-1}$.

c. Find its height after 10 seconds and a Cartesian equation. Comment on the model.


a

b

c

Further Example 34

A skater follows $x = 8\cos 20t,\ y = 12\sin(10t - \pi/3)$, where t is in minutes. Find the initial point, the self-intersection, the y-axis crossings and the time for a full circuit.



Initial point

Crossings

Circuit

Exercise 8E Page 218-220 35

1. A boat crosses a 75 m wide river, with eastward distance $x = 0.9t$ m and northward distance $y = -3.2t$ m, after t seconds.

a. Find the crossing time.

b. Find the southward drift.

c. Show the path is straight.

d. Find its speed.

a

b

c

d

Exercise 8E Page 218-220 36

2. A descending plane is modelled by $x = 80t,\ y = 3000 - 9.1t$, with distances in metres and t in minutes.

a. Find its initial height.

b. Discuss the stated domain $0 \leq t \leq 329$.

c. Find the horizontal distance at landing.

a

b


c

3. A ball is kicked from the ground at $20\,\text{m s}^{-1}$, angle $30°$. Its path is $x = 10\sqrt{3}t,\ y = 10t - 4.9t^2$.

a. Find its horizontal range.

b. Find when it is descending between heights 1.5 m and 2.5 m.

c. Find the nearest point where a player can head it in that interval.



a

b

c


4. A dolphin leaps along $x = 2t,\ y = 10t - 4.9t^2$, in metres and seconds.

a. Find the jump duration.

b. Find its horizontal distance.

c. Show the path is quadratic.

d. Find its maximum height.



a

b

c

d

5. A Ferris-wheel car has $x = 12\sin t,\ y = 12 - 12\cos t$, distances in metres and t in minutes.

a. Show its path is a circle of radius 12 m.


b. Find its maximum height.

c. Find the period and average speed.

a

b


c



6. A bowl has cross-section $x = t - 4\sin t,\ y = 1 - 2\cos t,\ -\pi/2 \le t \le \pi/2$.

a. Find the width of the opening.

b. Given that the deepest point is on the y-axis, find the depth.



a

b

Explore an implicit curve

$\sin(\sin x + \cos y) = \cos(\sin(xy) + \cos x)$.

Not every curve has a convenient explicit or parametric representation. A contour plot draws the zero set of the difference between the two sides.

