SLE₆ on Liouville quantum gravity as a growth-fragmentation process


William Da Silva


GDR Branchement


Based on joint work with Ellen Powell (Durham) and Alex Watson (UCL)

























































Bertoin, Curien, Kortchemski (2018)




Thm (BCK 18)

As $p \to \infty$, collection of perimeters scales to \mathbb{X} = growth-fragmentation process

Bertoin, Curien, Kortchemski (2018)

variant of 3/2-stable Lévy process




Thm (BCK 18)

As $p \to \infty$, collection of perimeters scales to

X =growth-fragmentation process

Bertoin, Curien, Kortchemski (2018)

variant of 3/2-stable Lévy process




Thm (BCK 18)

As $p \to \infty$, collection of perimeters scales to

Bertoin, Curien, Kortchemski (2018)

variant of 3/2-stable Lévy process




Thm (BCK 18)

As $p \to \infty$, collection of perimeters scales to

Bertoin, Curien, Kortchemski (2018)

variant of 3/2-stable Lévy process




Thm (BCK 18)

As $p \to \infty$, collection of perimeters scales to

X =growth-fragmentation process

Bertoin, Curien, Kortchemski (2018)


variant of 3/2-stable Lévy process




Thm (BCK 18)

As $p \to \infty$, collection of perimeters scales to

X =growth-fragmentation process




•
$$\gamma$$
-LQG disc: $\gamma = \sqrt{8/3}$



- γ -LQG disc: $\gamma = \sqrt{8/3}$
- space-filling curve η : SLE₆


GOAL: Build X in the continuum



- γ -LQG disc: $\gamma = \sqrt{8/3}$
- space-filling curve η : SLE₆

Branch η^z towards point $z \in \mathbb{D}$


GOAL: Build X in the continuum



•
$$\gamma$$
-LQG disc: $\gamma = \sqrt{8/3}$


• space-filling curve η : SLE₆

Branch η^z towards point $z \in \mathbb{D}$



- γ -LQG disc: $\gamma = \sqrt{8/3}$
- space-filling curve η : SLE₆
- Branching process: η^z , η^z'


GOAL: Build X in the continuum



- γ -LQG disc: $\gamma = \sqrt{8/3}$
- space-filling curve η : SLE₆

Branching process: η^z , η^z'


GOAL: Build X in the continuum



$$\gamma - LQG \ disc: \qquad \gamma = \sqrt{8/3}$$


• space-filling curve η : SLE₆






- γ -LQG disc: $\gamma = \sqrt{8/3}$
- space-filling curve η : SLE₆
- Branching process: η^z , η^z'


GOAL: Build X in the continuum



• space-filling curve η : SLE₆


Branching process: η^z , η^z





 $z \in \mathbb{D}$


- $oldsymbol{D}^z(s)$ c.c. of $\mathbb{D}\setminus \eta^z([0,s])$ containing z
- $X^{z}(s)$ (quantum) boundary length of $D^{z}(s)$



 $z \in \mathbb{D}$

- $oxed{D}^z(s)$ c.c. of $\mathbb{D}\setminus \eta^z([0,s])$ containing z
- $X^{z}(s)$ (quantum) boundary length of $D^{z}(s)$

$$\mathbb{X}(s) := \{X^z(s), z \in \mathbb{D}\}$$



 $z \in \mathbb{D}$

 $D^{z}(s)$ c.c. of $\mathbb{D} \setminus \eta^{z}([0,s])$ containing z


 $X^{z}(s)$ (quantum) boundary length of $D^{z}(s)$

 $\mathbb{X}(s) := \{X^z(s), z \in \mathbb{D}\}$

Thm (DS, Powell, Watson)

= growth-fragmentation process of BCK

variant of 3/2-stable Lévy process



Thm (DS, Powell, Watson)

★ = growth-fragmentation process of BCK

PRIOR ART

• Scaling limit from peeling Boltzmann triangulations
Bertoin, Curien, Kortchemski '18

PRIOR ART

- Scaling limit from peeling Boltzmann triangulations


 Bertoin, Curien, Kortchemski '18
- Peeling Boltzmann maps $\longrightarrow \mathbb{X}_{\alpha}$ Bertoin, Budd, Curien, Kortchemski '18


PRIOR ART


- Scaling limit from peeling Boltzmann triangulations
 Bertoin, Curien, Kortchemski '18
- Peeling Boltzmann maps $\longrightarrow \mathbb{X}_{\alpha}$ Bertoin, Budd, Curien, Kortchemski '18
- Brownian disc (metric) → time-change of X
 Le Gall, Riera '20

PRIOR ART

- Scaling limit from peeling Boltzmann triangulations
 Bertoin, Curien, Kortchemski '18
- Peeling Boltzmann maps $\longrightarrow \mathbb{X}_{\alpha}$ Bertoin, Budd, Curien, Kortchemski '18
- Brownian disc (metric) → time-change of X
 Le Gall, Riera '20
- CLE_{κ} GF on γ -LQG, $\sqrt{8/3} < \gamma < 2$ \longrightarrow \mathbb{X}_{α} , $\alpha = 4/\gamma^2$ Miller, Sheffield, Werner '22








PRIOR ART

- Scaling limit from peeling Boltzmann triangulations
 Bertoin, Curien, Kortchemski '18
- Peeling Boltzmann maps $\longrightarrow \mathbb{X}_{\alpha}$ Bertoin, Budd, Curien, Kortchemski '18
- Brownian disc (metric) → time-change of X
 Le Gall, Riera '20
- CLE_{κ} GF on γ -LQG, $\sqrt{8/3} < \gamma < 2 \longrightarrow \mathbb{X}_{\alpha}$, $\alpha = 4/\gamma^2$ Miller, Sheffield, Werner '22
- CLE₄ GF on critical LQG $\longrightarrow X_1$ Aïdékon, DS '22 Aru, Holden, Powell, Sun '23

Duplantier, Miller, Sheffield '21 Ang, Gwynne '21


unit
$$\gamma$$
-quantum disc



$$L_0 = 0, R_0 = 1$$

Duplantier, Miller, Sheffield '21 Ang, Gwynne '21


unit γ -quantum disc



$$L_0 = 0, R_0 = 1$$

Duplantier, Miller, Sheffield '21 Ang, Gwynne '21

unit γ -quantum disc




$$L_0 = 0, R_0 = 1$$

•
$$L_t = red$$
 quantum length $R_t = blue$ quantum length


Duplantier, Miller, Sheffield '21 Ang, Gwynne '21

unit γ -quantum disc




$$L_0 = 0, R_0 = 1$$

• $L_t = red$ quantum length $R_t = blue$ quantum length



Duplantier, Miller, Sheffield '21 Ang, Gwynne '21


unit γ -quantum disc




$$L_0 = 0, R_0 = 1$$


• $L_t = red$ quantum length


 R_t = blue quantum length





 $\rho(\gamma)$ -correlated Brownian excursion


























































Burdzy '85 Shimura '85 Duplantier, Miller, Sheffield '21














CONCLUSION

- **Growth-fragmentation** embedded in LQG/Brownian cone excursions
- New elementary proofs of old LQG results:

Target invariance property of SLE_6 on $\sqrt{8/3}$ -LQG

Law of area of quantum disc conditioned on perimeter

- Explicit description of BM subordinated on backward cone points (Le Gall)
- Questions about pathwise constructions of conditioned ssMPs