

Please stick the barcode label here.

Candidate Number												
------------------	--	--	--	--	--	--	--	--	--	--	--	--

MATHEMATICS Extended Part
Module 1 (Calculus and Statistics)
Question-Answer Book

8:30 am – 11:00 am (2½ hours)

This paper must be answered in English

INSTRUCTIONS

- (1) After the announcement of the start of the examination, you should first write your Candidate Number in the space provided on Page 1 and stick barcode labels in the spaces provided on Pages 1, 3, 5, 7, 9 and 11.
- (2) This paper consists of TWO sections, A and B.
- (3) Attempt ALL questions in this paper. Write your answers in the spaces provided in this Question-Answer Book. Do not write in the margins. Answers written in the margins will not be marked.
- (4) Graph paper and supplementary answer sheets will be supplied on request. Write your Candidate Number, mark the question number box and stick a barcode label on each sheet, and fasten them with string INSIDE this book.
- (5) Unless otherwise specified, all working must be clearly shown.
- (6) Unless otherwise specified, numerical answers should be either exact or given to 4 decimal places.
- (7) No extra time will be given to candidates for sticking on the barcode labels or filling in the question number boxes after the 'Time is up' announcement.



SECTION A (50 marks)

1. The table below shows the probability distribution of a discrete random variable X , where a is a constant and $p > 0$.

x	0	1	2
$P(X = x)$	$1 - 4p$	ap	p

(a) Find a .

(b) If $\text{Var}(2X + a^2) = 8\text{E}(aX - 1)$, find p .

(6 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Please stick the barcode label here.

2. Using photocopier X , the probability that there is dirt on a page of photocopy is $\frac{1}{5}$. A document of 6 pages is photocopied repeatedly by using X .

- (a) Find the probability that there is dirt on a photocopy of the document.
- (b) A photocopy of the document is called *acceptable* if there is dirt on fewer than 3 pages, otherwise the photocopy is called *unacceptable*.
- (i) Find the probability that a photocopy of the document is *unacceptable*.
- (ii) If an *unacceptable* photocopy of the document is just produced, find the expected number of *acceptable* photocopies produced between this *unacceptable* photocopy and the next *unacceptable* photocopy.

(6 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

3. Let A and B be two events. Denote the complementary event of A by A' . It is given that $P(B|A) = \frac{1}{2}$ and $P(B) = \frac{1}{3} + P(A)$. Suppose that $P(A' \cap B) = kP(A)$, where k is a constant.

- (a) By considering $P(A \cap B)$, or otherwise, prove that $k \neq \frac{1}{2}$. Also express $P(B)$ in terms of k .
- (b) Are A and B mutually exclusive? Explain your answer.
- (c) If A and B are independent, find k .

(7 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Please stick the barcode label here.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

4. A magazine publisher wants to estimate the proportion p of its current subscribers who will continue to subscribe next year. It is given that the publisher randomly selects 841 current subscribers, and 441 of them will continue to subscribe.
- (a) Find an approximate 95% confidence interval for p .
- (b) An approximate $\beta\%$ confidence interval for p is now constructed. The width of the confidence interval is 0.088. Find β correct to the nearest integer. (6 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Please stick the barcode label here.

5. Let $f(x) = (1 + k e^x)^3$, where k is a real constant.

- (a) Express, in terms of k , the constant term and the coefficient of x^2 in the expansion of $f(x)$.
- (b) If the constant term in the expansion of $f(x)$ is 27, find the coefficient of x^2 in this expansion.

(6 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

6. Define $g(x) = x + \frac{5}{x} + \ln x^4$ for all non-zero real numbers x .
- (a) Find $g'(x)$.
- (b) Someone claims that the maximum value of $g(x)$ is less than the minimum value of $g(x)$. Do you agree? Explain your answer.
- (c) Write down the equations of the two horizontal tangents to the graph of $y = g(x)$.

(6 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Please stick the barcode label here.

7. The total surface area of a solid right circular cylinder is $486\pi \text{ cm}^2$.

(a) Let $V \text{ cm}^3$ and $r \text{ cm}$ be the volume and the base radius of the circular cylinder respectively.
Find $\frac{dV}{dr}$.

(b) Can the volume of the circular cylinder exceed 5000 cm^3 ? Explain your answer.

(6 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

8. Let m be a non-zero constant.

(a) By considering $\frac{d}{dx}(xe^{mx})$, find $\int xe^{mx} dx$.

(b) If the area of the region bounded by the curve $y = xe^{mx}$, the x -axis and the straight line $x = 1$ is $\frac{1}{m}$, find m .

(7 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Please stick the barcode label here.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

SECTION B (50 marks)


9. Every morning Tom leaves home at 7:10 and walks to a certain bus stop to catch bus. He catches the earliest departing bus when he arrives at the bus stop. The time taken for Tom's walk follows a normal distribution with a mean of 15 minutes and a standard deviation of 2 minutes. There are two buses each morning, departing at 7:23 and 7:30 respectively.
- (a) Find the probability that Tom catches the bus departing at 7:23 on a certain morning. (2 marks)
- (b) Find the probability that Tom catches the bus departing at 7:30 on a certain morning. (1 mark)
- (c) Every morning Mary, Tom's student, walks to the same bus stop to catch bus. She catches the earliest departing bus when she arrives at the bus stop. It is given that the probability that Mary catches the bus departing at 7:23 on a certain morning is 0.3015 , while the probability that she catches the bus departing at 7:30 on a certain morning is 0.6328 . If Mary and Tom catch the same bus, Mary will greet Tom.
- (i) Find the probability that the 4th day in a week is the 2nd time Mary greets Tom.
- (ii) Given that Mary greets Tom on 2 certain mornings, find the probability that Mary and Tom catch the bus departing at 7:30 on these 2 mornings.
- (iii) Given that Mary greets Tom on 4 certain mornings, find the probability that Mary and Tom catch the bus departing at 7:23 on at least 1 of these 4 mornings.
- (iv) If Tom wants to have a higher chance of catching the bus departing at 7:23 than that of Mary, what is the latest time for him to leave home? Give your answer correct to the nearest minute. (10 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.



Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

10. A shopping mall launches a campaign to celebrate its fifth anniversary of the opening. A customer can throw a fair die 4 times to receive cash coupons. For each throw, a cash coupon is awarded according to the following table:

Result	1 , 2 or 3	4 or 6	5
Value of cash coupon	\$10	\$25	\$50

- (a) Find the probability that a customer receives cash coupons of a total value \$200 . (1 mark)
- (b) Find the probability that a customer receives cash coupons of a total value not less than \$150 . (2 marks)
- (c) A customer who receives cash coupons of a total value not less than \$150 can join a game. In the game, the customer presses a button 3 times. A number of cakes will show up on a screen for each press of the button, and the number of cakes shown follows a Poisson distribution with a mean of 5 . The result of each press of button is classified as follows:

Number of cakes	1 to 4	5	otherwise
Result	<i>Good</i>	<i>Excellent</i>	<i>Fair</i>


The customer receives a prize according to the following results:

Result	1 <i>Excellent</i> and 2 <i>Good</i>	2 <i>Excellent</i> and 1 <i>Good</i>	3 <i>Excellent</i>
Prize	a cup	a backpack	an oven

- (i) Find the probability that a customer joining the game receives a backpack.
- (ii) Given that a customer joining the game receives a prize, find the probability that the customer receives an oven.
- (iii) A customer who cannot join the game can still receive a cup by joining a lucky draw. In the lucky draw, the probability of receiving a cup is 0.01 . Given that a customer receives a cup, find the probability that the customer cannot join the game.

(9 marks)

Answers written in the margins will not be marked.



20 horizontal dotted lines for writing answers.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

11. The manager of a park models the rate of change of the number of adults (in thousand per month) visiting the park by

$$A(t) = 5 \ln(t^2 - 8t + 20) ,$$

where t is the number of months elapsed since the park opens. Denote the total number of adults visiting the park in the first 2 months since the park opens by α thousand. Let α_1 be the estimate of α by using the trapezoidal rule with 5 sub-intervals.

- (a) (i) Find α_1 .
- (ii) Is α_1 an over-estimate or an under-estimate? Explain your answer. (5 marks)
- (b) The manager models the rate of change of the number of children (in thousand per month) visiting the park by

$$B(t) = \frac{3^{2t+2}}{1+3^{2t}} ,$$

where t is the number of months elapsed since the park opens.

- (i) Find the total number of children visiting the park in the first 2 months since the park opens.
- (ii) The manager claims that in the first 2 months since the park opens, the difference of the total number of adults visiting the park and the total number of children visiting the park exceeds 40% of the total number of adults visiting the park. Do you agree? Explain your answer. (7 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.


Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.




A large rectangular area containing 20 horizontal dotted lines for writing answers. The lines are evenly spaced and extend across the width of the page.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.



Answers written in the margins will not be marked.

12. John studies the number of ducks P (in thousand) in a farm by

$$P = \frac{32}{a^{5+bt} + 8},$$

where a and b are constants and t ($t \geq 0$) is the number of months elapsed since the start of the study.

- (a) Express $\ln\left(\frac{32}{P} - 8\right)$ as a linear function of t . (2 marks)
- (b) John finds that the graph of the linear function obtained in (a) passes through the point $(1, \ln 2)$ and the intercept on the vertical axis of the graph is $\ln 32$.
- (i) Find a and b .
- (ii) Find $\frac{dP}{dt}$ and $\frac{d^2P}{dt^2}$.
- (iii) Estimate the number of ducks in the farm after a very long time. Hence, or otherwise, prove that the number of ducks in the farm does not exceed 4 thousand since the start of the study.
- (iv) Find the number of ducks in the farm when $\frac{dP}{dt}$ attains its greatest value.


(11 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.


Answers written in the margins will not be marked.



Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.




A large rectangular area containing 20 horizontal dotted lines for writing answers. The area is bounded by a thick black border.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.



A large rectangular area containing 20 horizontal dotted lines for writing answers. The lines are evenly spaced and extend across the width of the page.

END OF PAPER

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Standard Normal Distribution Table

<i>z</i>	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2517	.2549
0.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990
3.1	.4990	.4991	.4991	.4991	.4992	.4992	.4992	.4992	.4993	.4993
3.2	.4993	.4993	.4994	.4994	.4994	.4994	.4994	.4995	.4995	.4995
3.3	.4995	.4995	.4995	.4996	.4996	.4996	.4996	.4996	.4996	.4997
3.4	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4998
3.5	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998

Note : An entry in the table is the area under the standard normal curve between $x = 0$ and $x = z$ ($z \geq 0$). Areas for negative values of z can be obtained by symmetry.

