

A03 Bridging Neuroscience and Artificial Intelligence

Hoang Dinh

Houston City College

ITAI 4375: Artificial Intelligence in HealthCare

Professor Patricia McManus

February 4, 2026

Contents

Neural Network Comparison Analysis	2
The structure of a biological neuron, compared to familiar artificial neurons	2
How components map to artificial neural network elements.....	2
Comparison between biological and artificial neurons.....	3
Comparative Analysis of Biological and Artificial Neural Networks	3
Brain Architecture and AI Design.....	4
The hierarchical organization of the brain	4
How different regions specialize in specific tasks	5
Comparison to modern AI architectures	5
How biological emotion processing might inform AI systems.....	6
Implications for AI decision-making, risk assessment, pattern recognition, and adaptive learning	7
Integration Analysis and Future Implications	7
How understanding biological neural networks can improve AI design	7
Critical Comparison of Biological and Artificial Systems	8
Future Directions and Integration Possibilities.....	8
References	8

Neural Network Comparison Analysis

The structure of a biological neuron, compared to familiar artificial neurons

A biological neuron consists of several main parts that work together to process information.

According to WolframAlpha (n.d.), a neural cell includes a cell body, axon, and dendrites, which are responsible for receiving, integrating, and transmitting signals. Dendrites receive input from other neurons, similar to how artificial neurons receive input values in a neural network. The cell body, or soma, combines these incoming signals and determines whether the neuron will activate, which is comparable to the processing unit and activation function in artificial neurons.

The axon carries the output signal to other neurons, much like an output connection in an artificial network. Communication occurs at synapses, where the strength of connections can change over time, similar to adjustable weights in artificial models. Although biological neurons use electrical and chemical signals and artificial neurons rely on mathematical calculations, both systems are designed to process information through interconnected networks.

How components map to artificial neural network elements

The main components of a biological neuron can be directly compared to elements in an artificial neural network. WolframAlpha (n.d.) explains that a neural cell consists of dendrites, a cell body, and an axon, which work together to receive, process, and transmit information. In artificial systems, dendrites correspond to input connections because they receive signals from other neurons and pass them to the cell body. The cell body, or soma, functions as the processing unit, where incoming signals are integrated and a response is determined, similar to how an artificial neuron calculates a weighted sum and applies an activation function. The axon represents the output connection, as it carries the neuron's signal to other cells or network layers. Synapses,

which control the strength of communication between neurons, are comparable to weights in artificial neural networks because they determine how strongly one neuron influences another.

Comparison between biological and artificial neurons

Biological neurons have strongly influenced the design of artificial neurons by showing how information can be processed through interconnected cells. WolframAlpha (n.d.) describes neurons as consisting of dendrites, a cell body, and an axon that work together to receive, integrate, and transmit signals. This structure inspired artificial neural networks, where inputs are combined in a processing unit and passed forward as outputs. However, important differences exist in how these systems operate. Biological neurons communicate using electrical and chemical signals and adapt naturally through changes in synaptic strength, while artificial neurons rely on mathematical functions and learning algorithms such as backpropagation. Biological systems are highly energy-efficient, flexible, and capable of learning from small amounts of data, but they are difficult to study and control. Artificial networks are faster for large-scale computation and easier to train and modify, but they require large datasets and simplified models, limiting their ability to fully replicate human intelligence.

Comparative Analysis of Biological and Artificial Neural Networks

Biological Neural Network Characteristics	Artificial Neural Network Features	Examples from AI Architectures
Neurons consist of dendrites, cell body, and axon for signal processing (WolframAlpha, n.d.)	Artificial neurons use inputs, processing units, and outputs	Feedforward Neural Networks

Synapses adjust strength through learning and experience	Weights are updated using learning algorithms	CNNs (weight updates in convolution layers)
Signals are transmitted using electrical and chemical impulses	Information is transmitted using numerical values	RNNs (sequential numerical signals)
Learning occurs through synaptic plasticity	Learning occurs through backpropagation and optimization	Deep Neural Networks (DNNs)
Networks are highly parallel and distributed	Models run on parallel hardware (GPUs/TPUs)	CNNs and Transformer models
Neurons adapt continuously in real time	Models require retraining to adapt	RNNs and LSTM networks
Biological systems are energy efficient	Artificial systems consume high computational power	Large-scale Deep Learning models
Processing depends on timing and signal patterns	Processing depends on mathematical functions	Spiking Neural Networks (SNNs)

Brain Architecture and AI Design

The hierarchical organization of the brain

The human brain is organized in a hierarchical structure in which different regions work together at multiple levels of processing. According to WolframAlpha (n.d.), major brain structures such as the cerebellum, hypothalamus, and cerebral cortex belong to specific anatomical and

functional groups, showing how they are arranged within broader systems of the brain. For example, the primary motor cortex is part of the cerebral cortex and is responsible for voluntary movement, while the cerebellum is involved in motor coordination and learning. Lower-level structures, such as the lateral geniculate body, serve as relay centers that transmit sensory information to higher cortical areas for further processing. This layered organization allows basic sensory input and motor output to be handled by specialized regions before being integrated in higher-level cortical networks, creating an efficient and adaptable system for complex behavior.

How different regions specialize in specific tasks

Different regions of the human brain are specialized to perform specific cognitive, sensory, emotional, and motor functions. WolframAlpha (n.d.) shows that the primary motor cortex is responsible for voluntary movement, while the primary somatosensory cortex processes sensations such as touch, pain, and temperature. The cerebellum plays a key role in balance, coordination, and motor learning, helping refine physical actions. Emotional processing is mainly associated with the amygdala and hypothalamus, which regulate fear responses, mood, and hormone-related behaviors. Memory-related functions are linked to the hippocampus, where specialized neurons in the CA2 region support learning and memory formation. In addition, sensory relay centers such as the lateral geniculate body transmit visual information to the visual cortex for interpretation. This functional specialization allows the brain to efficiently divide complex tasks across interconnected regions.

Comparison to modern AI architectures

The hierarchical and specialized organization of the human brain has strongly influenced the design of modern artificial intelligence architectures. WolframAlpha (n.d.) shows that visual information in the brain is transmitted through relay centers such as the lateral geniculate body

before reaching the visual cortex for higher-level processing. This layered visual pathway is similar to how Convolutional Neural Networks (CNNs) process images through successive convolutional and pooling layers that extract increasingly complex features. Memory functions in the brain are supported by the hippocampus and its specialized neurons, including those in the CA2 region, which contribute to learning and information retention. This is comparable to Long Short-Term Memory (LSTM) networks, which are designed to store and update information over time. In addition, biological attention systems prioritize important sensory and emotional signals through interconnected regions such as the cortex and thalamus. Similarly, attention mechanisms in deep learning models focus computational resources on the most relevant parts of input data, improving learning efficiency and performance.

How biological emotion processing might inform AI systems

The amygdala plays a central role in emotional processing, especially in fear, threat detection, and emotional memory formation. According to WolframAlpha (n.d.), the amygdala is involved in emotional responses and memories of emotionally significant events. Detailed anatomical reports also show that the amygdala is part of the limbic system and is highly connected with regions responsible for decision-making and sensory processing. This close connection allows emotional information to influence perception, memory, and behavior. In artificial intelligence, this biological mechanism can inspire systems that evaluate not only factual data but also emotional or contextual importance. By modeling emotion-like processing, AI systems may become better at recognizing risk, prioritizing critical information, and responding appropriately in uncertain situations. Understanding how the amygdala filters and amplifies important signals can help researchers design AI models that balance rational analysis with adaptive emotional responses.

Implications for AI decision-making, risk assessment, pattern recognition, and adaptive learning

Biological emotion processing, especially through the amygdala and related limbic structures, has important implications for the development of intelligent artificial systems. WolframAlpha (n.d.) explains that the amygdala is responsible for evaluating emotional significance and generating appropriate behavioral responses. This ability allows humans to quickly assess risk and prioritize critical information in complex environments. In artificial intelligence, similar mechanisms could improve decision-making by enabling systems to weigh potential dangers and benefits more effectively. Emotion-inspired models may also enhance pattern recognition by highlighting unusual or threatening patterns that require immediate attention. Additionally, emotional feedback in the brain supports adaptive learning by reinforcing successful behaviors and discouraging harmful ones. By incorporating biologically inspired emotional processing, AI systems can become more flexible, context-aware, and responsive to changing environments, leading to safer and more reliable applications in areas such as healthcare, autonomous systems, and security.

Integration Analysis and Future Implications

How understanding biological neural networks can improve AI design

Understanding biological neural networks can help improve artificial intelligence by inspiring more flexible and efficient system designs. WolframAlpha (n.d.) shows that brain regions such as the hippocampus, cerebellum, and sensory relay centers work together through layered and specialized pathways. These interconnected structures suggest that future AI architectures could benefit from combining memory, optimization, and perception modules more closely. Biological systems also rely on local learning and feedback rather than centralized training, which may

inspire new biomimetic learning methods that reduce dependence on large datasets and improve robustness in uncertain environments.

Critical Comparison of Biological and Artificial Systems

Biological and artificial systems differ significantly in processing efficiency, learning ability, adaptability, and energy use. WolframAlpha (n.d.) indicates that regions such as the cerebellum and motor cortex coordinate complex actions using highly parallel and energy-efficient networks. The human brain operates using only about 20 watts of power, while large AI models require massive computational resources. Biological systems also learn continuously and generalize well from limited data, whereas artificial networks often require extensive retraining. However, artificial systems excel at speed and large-scale computation, making them suitable for data-intensive tasks despite lower adaptability.

Future Directions and Integration Possibilities

Insights from neuroscience may help address current AI limitations and guide future research. WolframAlpha (n.d.) shows that emotional and memory-related regions, such as the amygdala and hippocampus, influence decision-making and learning through interconnected feedback systems. Future AI models may incorporate emotion-inspired risk evaluation, biologically realistic memory systems, and attention-based routing mechanisms. Research may also focus on neuromorphic computing and spiking neural networks that better mimic brain signaling. Integrating biological principles with artificial systems could lead to more adaptive, efficient, and human-like intelligent technologies.

References

WolframAlpha. (n.d.). *Amygdala*. <https://www.wolframalpha.com>

WolframAlpha. (n.d.). *Cerebellum*. <https://www.wolframalpha.com>

WolframAlpha. (n.d.). *Human brain anatomy*. <https://www.wolframalpha.com>

WolframAlpha. (n.d.). *Hypothalamus*. <https://www.wolframalpha.com>

WolframAlpha. (n.d.). *Lateral geniculate body and medial geniculate body*.

<https://www.wolframalpha.com>

WolframAlpha. (n.d.). *Neural cell (neuron anatomy)*. <https://www.wolframalpha.com>

WolframAlpha. (n.d.). *Neurons in hippocampus CA2*. <https://www.wolframalpha.com>

WolframAlpha. (n.d.). *Primary motor cortex and primary somatosensory cortex*.

<https://www.wolframalpha.com>