

Does Structured Triage Context Help a Chest-Radiograph Pneumonia Model? A Time-Safe, Multi-Seed, Externally-Validated Multimodal Study with Missing-Data Cautions

Yazan Al-Dabain · Arafat Md Easin · Mufti Mahmud · Swakkhar Shatabda

Received: date / Accepted: date

Abstract Background: Emergency-department (ED) triage records structured vital signs before chest imaging, motivating multimodal models that fuse the chest radiograph (CXR) with clinical context for pneumonia detection. Whether such fusion helps—under strict time-safety and rigorous multi-seed evaluation—remains unclear, and prior studies often report single-run results without leakage or missing-data audits. **Methods:** We construct a temporally safe cohort by linking MIMIC-CXR-JPG, MIMIC-IV and MIMIC-IV-ED, anchoring every feature at the CXR acquisition time t_0 and admitting only measurements resolved at or before t_0 . On a patient-level temporal split (test $n=1,075$; prevalence 45.3%) we train image-only, concatenation- and attention-fusion, and clinical-baseline models across five random seeds, and add a time-safe early-laboratory panel as a third modality. We report AUROC, AUPRC, expected calibration error (ECE) and Brier score as mean $\pm$ SD over seeds

with paired per-seed deltas, patient-level bootstrap intervals, decision-curve and subgroup analyses, SHAP and Grad-CAM interpretability, external validation on NIH ChestX-ray14, and explicit temporal, split and preprocessing leakage audits. **Results:** A fine-tuned DenseNet-121 reaches AUROC 0.737 ± 0.003 , far above triage-only baselines (0.61). Adding triage vitals leaves discrimination unchanged (Δ AUROC $+0.004 \pm 0.009$, within a pre-specified ± 0.05 margin). A calibration advantage reported from a single checkpoint (ECE 0.040 vs 0.067) shrinks to Δ ECE -0.013 ± 0.016 across seeds—single-seed evaluation overstated it $\sim 2\times$. Adding early labs yields a small, seed-consistent AUROC gain ($+0.009 \pm 0.005$) that an ablation shows is reproduced exactly by lab *missingness indicators alone*, with no contribution from lab values, and is not leakage. Externally, discrimination transfers with minimal loss (AUROC 0.722 ± 0.005 on 112,120 NIH radiographs). **Conclusion:** The chest radiograph carries the discriminative signal; immediately available clinical context does not reliably add discrimination, and naive multimodal fusion can appear beneficial for reasons that will not generalise. We offer two transferable cautions for clinical multimodal AI: single-seed calibration inflation, and missingness masquerading as multimodal benefit.

Keywords Pneumonia detection · Chest radiograph · Multimodal fusion · Emergency department triage · Calibration · Reproducibility · Missing data · External validation

1 Introduction

Pneumonia is an acute lower respiratory tract infection and a leading cause of emergency-department (ED)

Yazan Al-Dabain
Department of Software Technology and Methodology, Faculty of Informatics, Eötvös Loránd University, Pázmány Péter Sétány 1/C, Budapest 1117, Hungary
E-mail:

Arafat Md Easin
Department of Software Technology and Methodology, Faculty of Informatics, Eötvös Loránd University, Pázmány Péter Sétány 1/C, Budapest 1117, Hungary
E-mail: arafatmd easin@inf.elte.hu

Mufti Mahmud
Department of Computer Science, Nottingham Trent University, NG11 8NS Nottingham, UK
E-mail: mufti.mahmud@ntu.ac.uk

Swakkhar Shatabda
Department of Computer Science and Engineering, BRAC University, Dhaka 1212, Bangladesh
E-mail: swakkhar.shatabda@bracu.ac.bd

presentation, hospital admission and death worldwide (Metlay et al, 2019). The frontal chest radiograph (CXR) is the recommended first-line imaging investigation, but its clinical value is bottlenecked by radiologist availability during peak ED demand. Automated CXR analysis that produces *well-calibrated* probability estimates is therefore an attractive decision-support tool, and deep convolutional networks now approach radiologist-level performance on several radiographic findings (Huang et al, 2017).

A structurally appealing but underused source of complementary information is the ED triage record. On arrival, a triage nurse records vital signs—temperature, heart rate, respiratory rate, peripheral oxygen saturation (SpO₂), blood pressure, a pain score—and an acuity level, *before* any imaging is ordered. Multimodal models that fuse the CXR with this immediately available context are widely proposed (Acosta et al, 2022; Hayat et al, 2022; Soenksen et al, 2022), on the intuition that combining the two streams should better reflect the clinical picture at the moment of decision.

Three methodological issues, however, recur across this literature and motivate the present study. First, *temporal safety*: clinical variables are frequently included without verifying they were actually available at prediction time, risking optimistic leakage from information generated after the decision point. Second, *single-seed reporting*: calibration and small discrimination effects are often reported from a single training run, although both are sensitive to training stochasticity (Guo et al, 2017; Nixon et al, 2019). Third, *missing data*: clinical inputs in the ED are heavily and non-randomly missing, so a model may exploit *whether* a measurement exists rather than its value—an effect that can masquerade as a genuine multimodal benefit.

We address a deliberately conditional question—*when, if ever, does immediately available clinical context help a CXR pneumonia model?*—under strict time-safety, with multi-seed replication and explicit missing-data and leakage audits, and report in line with prediction-model and medical-imaging guidance (Collins et al, 2024; Tejani et al, 2024). Our contribution is not a new architecture but a rigorous, reproducible, externally-validated evaluation that yields a clear answer and two transferable cautions:

- A **temporally safe, reproducible multimodal ED cohort** linking three MIMIC databases at the CXR acquisition time t_0 , with every feature verified to precede imaging (0 of 314,145 laboratory values post- t_0).
- A **multi-seed evaluation** (five seeds) of image-only, triage-fusion and laboratory-fusion models with paired bootstrap comparisons, showing

that immediately available clinical context is *discrimination-neutral*.

- **Caution 1 (single-seed calibration inflation)**: a calibration advantage that appears large from one checkpoint shrinks roughly twofold and becomes seed-fragile under replication.
- **Caution 2 (missingness as signal)**: a small “labs” gain that is reproduced entirely by laboratory *missingness indicators*, not the measured values, and is not attributable to leakage.
- **External validation** of the image model on 112,120 NIH ChestX-ray14 radiographs, showing cross-institution generalisation with an AUROC drop of only 0.016, together with subgroup, decision-curve, interpretability and modern-backbone analyses.

2 Background and Related Work

2.1 Pneumonia and ED triage

Pneumonia manifests on a frontal CXR as regions of increased opacity (consolidation) where air-filled alveoli fill with exudate. Reported CXR sensitivity for pneumonia is moderate (0.43–0.74) with high specificity, reflecting both the difficulty of early detection and reliable exclusion in unambiguous cases (Metlay et al, 2019). Triage vital signs quantify physiological perturbation and severity; they are highly correlated with pneumonia but non-specific, since tachycardia, hypoxia and high acuity accompany many respiratory presentations. Crucially, triage measurements are recorded before imaging is ordered—a workflow guarantee, not a statistical regularity—so they carry no risk of temporal leakage.

2.2 The MIMIC database family

The Medical Information Mart for Intensive Care (MIMIC) programme provides the largest publicly available de-identified clinical dataset linked to real hospital records from the Beth Israel Deaconess Medical Center (BIDMC). This study uses three interoperable components. **MIMIC-CXR-JPG** (Johnson et al, 2019) contains 227,835 chest-radiograph studies from 64,588 patients (2011–2019) with CheXpert labels for 14 findings. **MIMIC-IV** (Johnson et al, 2023b) provides hospital-admission and laboratory records, enabling temporal consistency checks and the laboratory modality. **MIMIC-IV-ED** (Johnson et al, 2023a) provides ED-specific records for all BIDMC emergency visits, including the structured triage table used as the

tabular input. All are distributed through PhysioNet under a Data Use Agreement; per-patient date shifts preserve temporal ordering and durations.

2.3 Automated labelling: CheXpert

CheXpert (Irvin et al, 2019) is an NLP pipeline that extracts binary labels for 14 thoracic conditions from free-text reports, assigning each a value of positive, negative, uncertain (-1), or not-mentioned. The uncertain label captures hedging language (“cannot exclude”, “possible pneumonia”). We adopt *u_ignore* (exclude uncertain studies) as the primary policy, which preserves label quality at the cost of a smaller, higher-prevalence evaluation set.

2.4 Deep learning and multimodal fusion

DenseNet-121 (Huang et al, 2017) propagates features through dense connectivity, $x_\ell = H_\ell([x_0, \dots, x_{\ell-1}])$, and forms a strong CXR backbone under transfer learning. Intermediate (feature-level) fusion extracts modality-specific embeddings and combines them—by concatenation, or by attention over modality tokens. Several works fuse structured clinical data with CXR for ICU or inpatient outcomes (Hayat et al, 2022; Soenksen et al, 2022); MedFuse in particular notes that healthcare modalities are collected asynchronously and are often missing, which our missing-data analysis makes central. To our knowledge, no prior work simultaneously (i) targets ED-specific pneumonia, (ii) uses ED triage vitals under a verified time-safety constraint, (iii) evaluates with a patient-level temporal split *and* multi-seed replication, and (iv) audits missingness and leakage explicitly. This study fills that gap.

3 Materials and Methods

3.1 Datasets and how they are used

The task is binary radiographic-pneumonia classification of a frontal ED CXR. Each of the three MIMIC components plays a distinct role. MIMIC-CXR-JPG supplies the *images* (JPEG radiographs) and the *labels* (CheXpert pneumonia), together with DICOM metadata (view position; StudyDate/StudyTime, from which the temporal anchor t_0 is derived). MIMIC-IV-ED supplies the *primary tabular modality*: the triage table (vitals, pain, acuity) recorded at ED intake, linked to each CXR through the ED stay. MIMIC-IV supplies the *admission records* used to verify temporal

consistency and the *laboratory events* used for the time-safe laboratory modality. The image branch consumes MIMIC-CXR-JPG; the tabular branch consumes MIMIC-IV-ED (triage) and, in the laboratory variant, MIMIC-IV (labs); the CheXpert labels supervise all models.

3.2 Cohort construction pipeline

A scripted pipeline links the sources through temporally constrained relational joins. Each CXR study is linked to a hospital admission (by `subject_id` and admission window) and to an ED stay (by `stay_id` with $t_0 \in [\text{intime}, \text{outtime}]$); one-to-one study-to-stay integrity is enforced by deduplication. The triage table is merged on `stay_id`; studies are restricted to frontal (PA/AP) projections; CheXpert pneumonia labels are merged and the *u_ignore* policy applied. The temporal anchor is $t_0 = \text{StudyDate} + \text{StudyTime}$ from DICOM metadata. The resulting cohort comprises 9,154 ED-anchored studies; the patient-level temporal split (below) assigns train 7,144 / validation 930 / test 1,080 studies. Models train and evaluate only on radiographs present on disk in our data snapshot, yielding a held-out test set of $n=1,075$ (487 positive / 588 negative; prevalence 45.3%). The entire pipeline is reproducible from a single entry point, emitting versioned manifests at each stage. Table 1 summarises the cohort: the split sizes, test-set demographics (median age 65; 55% female; predominantly White with substantial Black and Hispanic representation), the near-even PA/AP view split, and the triage vital-sign distributions with their missingness rates (3–8% per vital), which motivate the per-vital missingness indicators.

3.3 Time-safety and the patient-level temporal split

Every clinical feature is either a triage measurement (recorded at intake, before imaging by protocol) or a laboratory value with charttime in $(t_0 - 24\text{h}, t_0]$. Time-safety was verified empirically on the extracted laboratory events: *zero* of 314,145 measurements have a charttime after t_0 (median lead time 1 h). Patients are ranked by their earliest t_0 and partitioned 80/10/10 into train/validation/test, so no patient appears in more than one split (verified: 0 cross-split subjects, 0 duplicate studies). This ordinal-time construction approximates prospective deployment. The tabular preprocessor (median imputation, standardisation, one-hot encoding) is fit on the training split only.

Table 1 Cohort characteristics. Splits are patient-level and temporally ordered; demographic and vital-sign statistics are for the test split.

Characteristic	Value
<i>Splits: studies / patients / prevalence</i>	
Train	7,144 / 6,045 / 46.9%
Validation	930 / 787 / 48.8%
Test	1,080 / 890 / 45.4%
<i>Test-set demographics</i>	
Age, mean (median)	64 (65)
Female	55.0%
White / Black / Hispanic	678 / 258 / 71
Asian / Other	36 / 37
View: PA / AP	56.8% / 43.2%

Triage vital (test set)	mean±SD (% missing)
Temperature (°F)	98.5±1.4 (5.9%)
Heart rate (bpm)	87.7±19.6 (3.3%)
Respiratory rate (/min)	18.6±3.5 (4.5%)
SpO ₂ (%)	97.4±3.6 (4.8%)
Systolic BP (mmHg)	135±24 (4.0%)
Diastolic BP (mmHg)	73±16 (4.3%)
Pain (0–10)	3.8±4.3 (8.2%)
Acuity (ESI 1–5)	2.3±0.7 (1.2%)

3.4 Feature engineering

Triage features are temperature, heart rate, respiratory rate, SpO₂, systolic and diastolic blood pressure, pain score and ESI acuity, plus the radiograph view and demographics. Physiologically implausible values are clipped to clinical ranges. For each vital a binary *missingness indicator* is added and the value median-imputed; because ED missingness is not missing-at-random, the indicator itself carries information. **Laboratory features** comprise 26 concepts (complete blood count, metabolic panel, liver panel, blood gas, lactate, and the inflammatory markers CRP and procalcitonin). For each study and concept we take the last value resolved at or before t_0 within the 24 h window, and add a missingness flag recomputed after the cohort join.

3.5 Models

The **image model** is DenseNet-121, initialised from ImageNet, multilabel-pretrained on 182,637 non-ED MIMIC-CXR studies (14 CheXpert findings, masked binary cross-entropy over observed labels), then fine-tuned for binary pneumonia at 224×224 with class-balanced loss and differential learning rates (5×10^{-5} head, 10^{-5} backbone). Multilabel pretraining minimises a masked binary cross-entropy that ignores not-mentioned CheXpert labels,

$$\mathcal{L}_{\text{mBCE}} = -\frac{1}{\sum_i m_i} \sum_i m_i [y_i \log \hat{p}_i + (1 - y_i) \log(1 - \hat{p}_i)], \quad (1)$$

where $m_i \in \{0, 1\}$ masks unobserved conditions. The triage vector is encoded by a two-layer TabularMLP (batch-normalised, dropout 0.2) to a 128-d embedding e_t . The **concatenation** model concatenates the 1024-d image embedding e_v with e_t , $[e_v; e_t] \in \mathbb{R}^{1152}$, and

applies a fully connected head; the **attention** model projects each embedding to a d -dimensional token, prepends a learnable CLS token, and applies a two-layer Transformer whose scaled dot-product attention

$$\text{Attention}(Q, K, V) = \text{softmax}\left(\frac{QK^\top}{\sqrt{d_k}}\right)V \quad (2)$$

mixes the $[\text{CLS}, e_v, e_t]$ tokens; the CLS output feeds the classifier (Acosta et al, 2022). The **+labs** model appends the 26 lab values and their flags to the tabular vector. **Clinical baselines** are logistic regression and XGBoost (Chen and Guestrin, 2016) on triage features. A **modern-backbone baseline** uses a CheXpert-trained DenseNet-121 (Cohen et al, 2022) (not trained on MIMIC), zero-shot and as a frozen-feature linear probe. All deep models share the multilabel-pretrained backbone so the multi-seed analysis isolates fine-tuning stochasticity.

3.6 Training and evaluation

Each deep model is trained across five seeds $\{42, 123, 456, 789, 1000\}$ with early stopping on validation AUPRC. Primary metrics are AUROC, AUPRC, ECE and Brier score, reported as mean±SD over seeds with paired per-seed deltas against the image-only model on identical test rows; within-checkpoint uncertainty uses a 2,000-replicate patient-level paired bootstrap (patients, not studies, are resampled). A non-inferiority margin $\Delta_0 = -0.05$ AUROC was pre-specified (Piaggio et al, 2012). Calibration is assessed with the ECE

$$\text{ECE} = \sum_{b=1}^B \frac{n_b}{n} |\bar{p}_b - \bar{y}_b|, \quad (3)$$

(with $B=10$ uniform bins; $\bar{p}_b, \bar{y}_b$ the mean predicted probability and observed frequency in bin b), the maximum calibration error, Brier score and reliability diagrams; ECE bin-count/scheme sensitivity is examined.




Fig. 1 ROC curves on the held-out test set (5-seed mean predictions for the deep models; AUROC of the ensemble curve shown, slightly above the per-seed means in Table 2).

Clinical utility uses decision-curve analysis with net benefit

$$\text{NB}(p_t) = \frac{\text{TP}}{n} - \frac{\text{FP}}{n} \cdot \frac{p_t}{1 - p_t}. \quad (4)$$

Interpretability uses exact TreeSHAP (Chen and Guestrin, 2016) for the XGBoost baseline and Grad-CAM for the image branch. We additionally compute subgroup metrics (sex, age, race, acuity, view), operating points at fixed sensitivity, a laboratory *missingness-only* ablation, and external validation on all 112,120 NIH ChestX-ray14 radiographs (Wang et al, 2017) with preprocessing matched to the internal transform.

4 Results

4.1 Discrimination: imaging dominates, fusion is neutral

Table 2 and Figs. 1–3 summarise the primary results. Both CXR models exceed the triage-only baselines by more than 12 AUROC points, confirming that the radiograph carries the discriminative signal. Adding triage vitals does not change discrimination: the paired concat–image ΔAUROC is $+0.004 \pm 0.009$ (range $[-0.007, +0.014]$), with the effect’s lower bound well inside the pre-specified -0.05 non-inferiority margin. Attention fusion matches image AUROC but is severely and variably miscalibrated (ECE 0.076 ± 0.042 ; Fig. 4), illustrating that AUROC alone is insufficient for model selection. The precision-recall curves (Fig. 2) mirror the AUROC ordering at the 45.3% prevalence.




Fig. 2 Precision-recall curves at test prevalence 45.3%.

4.2 Caution 1: single-seed evaluation inflates the calibration effect



A single checkpoint suggested the concat model was markedly better calibrated (ECE 0.040 vs 0.067, a 40% reduction). Across seeds, however, the image-only ECE alone ranges 0.041–0.060, and the paired advantage shrinks to $\Delta\text{ECE} -0.013 \pm 0.016$ (4/5 seeds favour concat; the range crosses zero). The single-seed estimate thus overstated the benefit roughly twofold. The apparent gap is also binning-dependent (Fig. 6): under coarser or quantile bins it shrinks further (Nixon et al, 2019). Brier scores are essentially unchanged, so any effect is distributional alignment rather than refinement. Reliability diagrams (Fig. 5) show all deep models tracking the diagonal closely, with attention fusion the clear outlier.

4.3 Caution 2: the “labs” benefit is missing-data structure

Appending early labs gives the only seed-consistent AUROC gain (labs–image $+0.009 \pm 0.005$, 5/5 seeds, range excluding zero). Three analyses show this is neither chemistry nor leakage. *Ablation*: a model using laboratory *missingness flags only* (no values) reproduces the gain exactly (flags–image $+0.009$; Fig. 4); the lab values add nothing further to discrimination. *Subcohort*: on the 269 test studies that actually have labs, discrimination is tied (labs 0.778 vs image 0.776). *Leakage audit*: 0 of 314,145 lab measurements post- t_0 ; 0 cross-split subjects; preprocessing fit on train only; the “any lab drawn” indicator has near-null univariate AUROC (0.464). The gain therefore reflects the weak comple-

Table 2 Multi-seed test-set performance (mean $\pm$ SD over 5 seeds; $n=1,075$; prevalence 45.3%). Clinical baselines are single-fit.

Model	AUROC	AUPRC	ECE	Brier
Image-only DenseNet-121	0.737 ± 0.003	0.719 ± 0.004	0.053 ± 0.008	0.207 ± 0.001
Multimodal — concat (triage)	0.741 ± 0.006	0.715 ± 0.005	0.040 ± 0.014	0.205 ± 0.002
Multimodal — attention (triage)	0.738 ± 0.006	0.713 ± 0.012	0.076 ± 0.042	0.212 ± 0.010
Multimodal — concat (+labs)	0.747 ± 0.004	0.724 ± 0.004	0.043 ± 0.011	0.203 ± 0.002
Logistic regression (triage)	0.606	0.548	0.037	0.242
XGBoost (triage)	0.611	0.567	0.046	0.241

**Fig. 3** Per-model AUROC and ECE across five seeds (points, seeds; red bar, mean). Discrimination is stable; the image-only ECE alone varies substantially by seed.**Fig. 4** Paired fusion—image effects across seeds (mean $\pm$ SD). Every effect straddles zero; the labs and lab-flags-only deltas are identical.

mentary signal of *whether* bloodwork was ordered, is small, and lies within the patient-level bootstrap CI.

4.4 Clinical availability of laboratory results

Under time-safe matching, only 25.4% of ED-anchored CXR studies (272/1,075 test) have any resolved laboratory value at or before imaging (Fig. 7). Complete-blood-count and metabolic panels reach $\sim 25\%$, but the most pneumonia-specific markers—CRP and

procalcitonin—are present in only 0.5% and 0.6% of studies. At the moment of imaging the informative labs are essentially unavailable, which explains both the small labs effect and its missingness origin.

4.5 Operating points and clinical utility

ED triage for pneumonia prioritises sensitivity. At a fixed sensitivity of 0.90 the specificity is 0.31 (image) versus 0.32 (concat); at 0.95, 0.17 versus 0.18




Fig. 5 Reliability diagrams (10 bins). The deep models track the diagonal; attention fusion systematically under-predicts.




Fig. 6 ECE under different binning schemes; the image-concat calibration gap depends on the binning choice.

Table 3 Operating points at fixed sensitivity (mean over 5 seeds).

Model	Sens.	Spec.	PPV	NPV
Image	0.90	0.310	0.520	0.791
Concat	0.90	0.322	0.524	0.797
Image	0.95	0.173	0.488	0.808
Concat	0.95	0.179	0.490	0.814

(Table 3)—clinically equivalent operating behaviour. Decision-curve analysis (Fig. 8) shows both CXR models providing substantially higher net benefit than the triage-only baselines and the treat-all/treat-none references across the clinically relevant threshold range, with the image and multimodal curves essentially overlapping.




Fig. 7 Fraction of test studies with each laboratory value resolved at or before t_0 ; pneumonia-specific markers (red) are essentially unavailable at imaging time.




Fig. 8 Decision-curve analysis: net benefit versus threshold probability. Imaging models dominate the triage baselines and reference strategies.

4.6 Subgroup performance

Table 4 and Fig. 9 report subgroup AUROC. Performance is markedly lower for female than male patients (0.71 vs 0.77) and for Hispanic patients (0.64) relative to White (0.75); PA views outperform AP (0.75 vs 0.72). Fusion does not consistently close these gaps and, in the smallest strata, can slightly worsen calibration. These disparities echo documented underdiagnosis biases in CXR models (Seyyed-Kalantari et al, 2021; Larrazabal et al, 2020) and are reported here as a pre-requisite fairness audit rather than a solved problem; small minority strata carry wide confidence intervals.

Table 4 Subgroup AUROC (mean over 5 seeds) for image-only and concat fusion. Small strata (e.g. Hispanic, Asian) have wide CIs.

Subgroup	Image	Concat
Female	0.713	0.715
Male	0.770	0.776
White	0.751	0.756
Black	0.710	0.709
Hispanic	0.640	0.655
PA view	0.748	0.758
AP view	0.721	0.715
High acuity (1–2)	0.759	0.761
Lower acuity (3–5)	0.705	0.711




Fig. 9 Subgroup AUROC (mean±SD across seeds); fusion does not close sex, race or view gaps.

Table 5 Mean |SHAP| for the triage XGBoost model (test set).

Feature	Mean SHAP
SpO ₂	0.148
Heart rate	0.099
Temperature	0.078
Respiratory rate	0.062
PA-view indicator	0.053
Systolic BP	0.043
Diastolic BP	0.042
Acuity (ESI)	0.031
Pain score	0.012

4.7 Interpretability

For the triage-only XGBoost model, exact TreeSHAP attributes the largest importance to SpO₂ (mean |SHAP| = 0.148), followed by heart rate (0.099), temperature (0.078) and respiratory rate (0.062) (Table 5; Fig. 10)—consistent with the pathophysiology of pneumonia, in which alveolar consolidation impairs gas exchange. Grad-CAM on the image branch (Fig. 11) localises activation over regions of consolidation in confident true positives, whereas a false positive attends to overlapping opacity and a false negative shows diffuse, non-localised activation.




Fig. 10 SHAP summary (beeswarm) for the triage XGBoost model; low SpO₂ pushes predictions toward pneumonia.




Fig. 11 Grad-CAM on the image model for representative test cases: a confident true positive, a false positive, and a false negative.

4.8 Modern backbone and external validation

A CheXpert-trained DenseNet-121 (Cohen et al, 2022), used zero-shot (AUROC 0.667) or as a frozen-feature linear probe (0.687), does not outperform our task-tuned model (0.737), indicating the conclusions are not an artefact of an outdated backbone. On external validation against all 112,120 NIH ChestX-ray14 radiographs (Wang et al, 2017) (a different institution, scanner population and label pipeline; prevalence 1.28%), discrimination transfers with minimal loss: AU-




Fig. 12 Cross-institution generalisation: per-seed AUROC on internal MIMIC ($n=1,075$) versus external NIH ChestX-ray14 ($n=112,120$); mean drop -0.016 .

ROC 0.722 ± 0.005 across five seeds (ensemble 0.726), only 0.016 below the internal 0.737 (Fig. 12). As expected under the large prevalence shift ($45\% \rightarrow 1.3\%$), prevalence-dependent metrics degrade (AUPRC 0.030, ECE 0.44) because probabilities are calibrated to the internal base rate; AUROC, which is prevalence-independent, is the appropriate cross-dataset metric, and it holds despite domain shift and NLP-derived external labels.

5 Discussion

The chest radiograph carries the discriminative signal for radiographic pneumonia in this ED cohort. Immediately available structured clinical context—triage vitals, and even a broad early-laboratory panel—does not add reliable discrimination, and the image model generalises externally with minimal loss. The asymmetry is intuitive: the radiograph encodes the outcome (radiographic consolidation) almost directly, whereas vital signs and labs encode severity and physiological perturbation that are correlated with, but not specific to, pneumonia.

Two findings generalise beyond this application. First, a calibration advantage that appears substantial from a single training run (a 40% ECE reduction) is roughly halved and becomes seed-fragile under five-seed replication; single-seed calibration claims should be treated with caution, and multi-seed replication should be standard. Second, where clinical inputs are heavily and non-randomly missing, a multimodal model can exploit missingness structure and present a spurious benefit that neither reflects the measured values (the lab-present subcohort shows no value effect) nor constitutes leakage (temporal, split and preprocessing audits are clean). The missingness-only ablation makes

this mechanism explicit and is, to our knowledge, a rarely reported control in clinical multimodal studies. Both cautions are actionable and inexpensive.

Comparison to related work. Unlike prior multimodal CXR studies that report single-run gains on ICU or inpatient cohorts (Hayat et al, 2022; Soenksen et al, 2022), we evaluate ED pneumonia under a verified time-safety constraint with multi-seed replication and explicit audits, and reach a negative but well-powered conclusion on discrimination. Direct numerical comparison across cohorts and label definitions is uninformative; the methodological contribution is the rigour of the evaluation and the two cautions.

Limitations. Training and multimodal evaluation are single-institution (MIMIC) and retrospective; external validation confirms the image branch generalises to NIH but is necessarily limited to imaging, as no external dataset links ED triage to the CXR, and probabilities require recalibration to the target base rate before deployment (the *u.ignore* policy also raises internal evaluation prevalence to 45.3%). The lab-present subcohort is small ($n=269$); the modern-backbone comparison uses frozen features rather than end-to-end fine-tuning; and NIH labels are NLP-derived, adding label noise to the external estimate.

6 Conclusion

Under strict time-safety and multi-seed evaluation, fusing immediately available clinical context with the chest radiograph does not reliably improve pneumonia discrimination, and the one apparent gain is an artefact of missing-data structure. The image model itself is robust and generalises across institutions with an AUROC drop of only 0.016. Beyond the negative result, the study contributes a reproducible, time-safe multimodal ED pipeline and two reusable cautions—single-seed calibration inflation and missingness-as-signal—for the growing field of clinical multimodal AI. Future work includes end-to-end fine-tuning of modern foundation backbones, multi-site validation, and prospective evaluation with recalibration to the local base rate.

Declarations

Acknowledgements. The authors thank the MIT Laboratory for Computational Physiology and PhysioNet for maintaining and distributing the MIMIC databases, and the U.S. National Institutes of Health for the ChestX-ray14 dataset.

Authors' contributions. YAD designed and implemented the data pipeline, trained all models and performed the analyses. AME conceptualised and supervised the study and contributed to the manuscript. MM and SS contributed to methodology, interpretation and critical revision. All authors read and approved the final manuscript.

Funding. The authors received no specific funding for this work.

Conflict of interest. The authors declare no competing interests.

Ethics approval. All data are publicly available and de-identified (MIMIC under PhysioNet's credentialed Data Use Agreement; NIH ChestX-ray14 public release). No additional ethical approval was required; procedures followed the Declaration of Helsinki.

Data and code availability. MIMIC and NIH ChestX-ray14 are publicly available (MIMIC via PhysioNet credentialing). All pipeline, training and evaluation code is available at <https://github.com/wa5rx3/MultiModalPneumonia>; every reported number is regenerated by a named script tied to the fixed temporal split.

References

- Acosta JN, Falcone GJ, Rajpurkar P, Topol EJ (2022) Multimodal biomedical AI. *Nature Medicine* 28(9):1773–1784, DOI 10.1038/s41591-022-01981-2
- Chen T, Guestrin C (2016) XGBoost: A scalable tree boosting system. In: *KDD*, pp 785–794, DOI 10.1145/2939672.2939785
- Cohen JP, Viviano JD, Bertin P, Morrison P, Torabian P, Guarrera M, Lungren MP, Chaudhari A, Brooks R, Hashir M, Bertrand H (2022) TorchXRyVision: A library of chest X-ray datasets and models. In: *Proceedings of Machine Learning Research (MIDL)*, vol 172, pp 231–249
- Collins GS, Moons KGM, Dhiman P, et al (2024) TRI-POD+AI statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods. *BMJ* 385:e078378, DOI 10.1136/bmj-2023-078378
- Guo C, Pleiss G, Sun Y, Weinberger KQ (2017) On calibration of modern neural networks. In: *International Conference on Machine Learning (ICML)*, vol 70, pp 1321–1330
- Hayat N, Geras KJ, Shamout FE (2022) MedFuse: Multi-modal fusion with clinical time-series data and chest X-ray images. In: *Machine Learning for Healthcare (MLHC)*, PMLR, vol 182, pp 479–503
- Huang G, Liu Z, van der Maaten L, Weinberger KQ (2017) Densely connected convolutional networks. In: *CVPR*, pp 4700–4708, DOI 10.1109/CVPR.2017.243
- Irvin J, Rajpurkar P, Ko M, et al (2019) CheXpert: A large chest radiograph dataset with uncertainty labels and expert comparison. In: *AAAI*, vol 33, pp 590–597, DOI 10.1609/aaai.v33i01.3301590
- Johnson AEW, Pollard TJ, Berkowitz SJ, Greenbaum NR, Lungren MP, Deng Cy, Mark RG, Horng S (2019) MIMIC-CXR-JPG, a large publicly available database of labeled chest radiographs. *arXiv:1901.07042*, URL <https://arxiv.org/abs/1901.07042>
- Johnson AEW, Bulgarelli L, Pollard TJ, Horng S, Celi LA, Mark RG (2023a) MIMIC-IV-ED: Emergency department module of MIMIC-IV. URL <https://physionet.org/content/mimic-iv-ed/>
- Johnson AEW, Bulgarelli L, Shen L, et al (2023b) MIMIC-IV, a freely accessible electronic health record dataset. *Scientific Data* 10(1):1, DOI 10.1038/s41597-022-01899-x
- Larrazabal AJ, Nieto N, Peterson V, Milone DH, Ferrante E (2020) Gender imbalance in medical imaging datasets produces biased classifiers for computer-aided diagnosis. *PNAS* 117(23):12,592–12,594, DOI 10.1073/pnas.1919012117
- Metlay JP, Waterer GW, Long AC, et al (2019) Diagnosis and treatment of adults with community-acquired pneumonia (ats/idsa guideline). *American Journal of Respiratory and Critical Care Medicine* 200(7):e45–e67, DOI 10.1164/rccm.201908-1581ST
- Nixon J, Dusenberry MW, Zhang L, Jerfel G, Tran D (2019) Measuring calibration in deep learning. In: *CVPR Workshops*, URL <https://arxiv.org/abs/1904.01685>
- Piaggio G, Elbourne DR, Pocock SJ, Evans SJW, Altman DG (2012) Reporting of noninferiority and equivalence randomized trials: Extension of the CONSORT 2010 Statement. *JAMA* 308(24):2594–2604, DOI 10.1001/jama.2012.87802
- Seyyed-Kalantari L, Zhang H, McDermott MBA, Chen IY, Ghassemi M (2021) Underdiagnosis bias of artificial intelligence algorithms applied to chest radiographs in under-served patient populations. *Nature Medicine* 27:2176–2182, DOI 10.1038/s41591-021-01595-0
- Soenksen LR, Ma Y, Zeng C, Boussieux L, Villalobos Carballo K, Na L, Wiberg HM, Li ML, Fuentes I, Bertsimas D (2022) Integrated multimodal arti-

ficial intelligence framework for healthcare applications. *npj Digital Medicine* 5:149, DOI 10.1038/s41746-022-00689-4

Tejani AS, Klontzas ME, Gatti AA, Mongan JT, Moy L, Park SH, Kahn CE (2024) Checklist for Artificial Intelligence in Medical Imaging (CLAIM): 2024 update. *Radiology: Artificial Intelligence* 6(4):e240,300, DOI 10.1148/ryai.240300

Wang X, Peng Y, Lu L, Lu Z, Bagheri M, Summers RM (2017) ChestX-ray8: Hospital-scale chest X-ray database and benchmarks on weakly-supervised classification and localization of common thorax diseases. In: *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, pp 2097–2106, DOI 10.1109/CVPR.2017.369