

Visualizing the Manhattan Curve

Thesis Prospectus

William Clampitt

May 7, 2025

Topology

Definition (Topological Space)

A **topological space** X is a set together with a collection \mathcal{T} of subsets of X where \mathcal{T} contains the sets X , \emptyset , and is closed under finite intersections and arbitrary unions. The elements of \mathcal{T} are called open sets.

Topology

Definition (Topological Space)

A **topological space** X is a set together with a collection \mathcal{T} of subsets of X where \mathcal{T} contains the sets X , \emptyset , and is closed under finite intersections and arbitrary unions. The elements of \mathcal{T} are called open sets.

Definition (d -manifold)

Let $d \in \mathbb{Z}_{\geq 0}$. A **d -manifold** is topological space that is second countable, Hausdorff, and locally Euclidean of dimension d .

Topology

Definition (Topological Space)

A **topological space** X is a set together with a collection \mathcal{T} of subsets of X where \mathcal{T} contains the sets X , \emptyset , and is closed under finite intersections and arbitrary unions. The elements of \mathcal{T} are called open sets.

Definition (d -manifold)

Let $d \in \mathbb{Z}_{\geq 0}$. A **d -manifold** is topological space that is second countable, Hausdorff, and locally Euclidean of dimension d .

Definition (Locally Euclidean of Dimension d)

A topological space M is **locally Euclidean of dimension d** if every point of M is contained in an open set in M that is homeomorphic to an open subset of \mathbb{R}^d .



Surface Examples






Surface Examples





Surface Examples



Surface Examples



Surface Examples




- \mathbb{RP}^2 : the set of all lines passing through the origin in \mathbb{R}^3 .
- A basis for the topology in \mathbb{RP}^2 is the sets of lines that form a bounded angle from a fixed line ℓ in \mathbb{R}^3 .

Alternate View of \mathbb{RP}^2

Definition

A **plane** in \mathbb{R}^3 is a 2-dimensional vector subspace of \mathbb{R}^3 . An **affine plane** A is a translate of a plane P by a nonzero vector v that is not in P .




Alternate View of \mathbb{RP}^2

Lemma

Let P be a plane passing through the origin in \mathbb{R}^3 and let A be an affine plane which is a translate of P by a nonzero vector v not in P . Then,

$$\mathbb{RP}^2 \cong A \sqcup \pi(P)$$

Note: $\pi(P)$ is the subset of \mathbb{RP}^2 of lines passing through the origin and contained in P .



Fundamental Groups

Definition

Let S be a path-connected a surface. For any point $p \in S$ the **fundamental group** of S is the set of equivalence classes (under homotopy) of the loops on S based at p with the concatenation operation. This group is denoted $\pi_1(S)$.

Questions?


Hyperbolic Structures

The Upper Half Plane

Definition


The **hyperbolic plane** \mathbb{H}^2 is the metric space

$$\mathbb{H}^2 = \{(x, y) \in \mathbb{R}^2 : y > 0\} \quad d(u, v) = \inf_{u \rightarrow v} \int_0^1 \frac{\sqrt{\dot{x}(t)^2 + \dot{y}(t)^2}}{y(t)} dt$$



Hyperbolic Structure on a Surface S



- ▶ Every point $p \in S$ has a neighborhood that maps to an open subset of \mathbb{H}^2



- ▶ $\varphi_\alpha: U_\alpha \rightarrow V_\alpha$
- ▶ $\varphi_\beta: U_\beta \rightarrow V_\beta$
- ▶ $\varphi_\beta \circ \varphi_\alpha|_{\varphi_\alpha(U_\alpha) \cap \varphi_\beta(U_\beta)}$
- ▶ Needs to be isometry

- ▶ This gives a notion of distance between two points on the surface S .
- ▶ Many surfaces have lots of hyperbolic structures, but ...
- ▶ $S_{0,3}$ only has one.

The Hyperbolic Structure on $S_{0,3}$



Convex Real Projective Structures

Definition

- ▶ An open set $\Omega \subseteq \mathbb{RP}^2$ is **proper** if there exists a plane $P \subseteq \mathbb{R}^3$ passing through the origin such that $\overline{\Omega} \cap \pi(P) = \emptyset$.

Convex Real Projective Structures




Definition

- ▶ An open set $\Omega \subseteq \mathbb{RP}^2$ is **proper** if there exists a plane $P \subseteq \mathbb{R}^3$ passing through the origin such that $\overline{\Omega} \cap \pi(P) = \emptyset$.
- ▶ A proper set $\Omega \subseteq \mathbb{RP}^2$ is **convex** if, for any two points $x, y \in \Omega$, the line l_{xy} passing through x and y intersects Ω in a connected segment.

Convex Real Projective Structures

Definition

- ▶ An open set $\Omega \subseteq \mathbb{RP}^2$ is **proper** if there exists a plane $P \subseteq \mathbb{R}^3$ passing through the origin such that $\overline{\Omega} \cap \pi(P) = \emptyset$.
- ▶ A proper set $\Omega \subseteq \mathbb{RP}^2$ is **convex** if, for any two points $x, y \in \Omega$, the line l_{xy} passing through x and y intersects Ω in a connected segment.
- ▶ Ω is **strictly convex** if $\partial\Omega$ contains no straight line segments.




Convex Real Projective Structures

Given a strictly convex set Ω in \mathbb{RP}^2 .

Definition

The **Hilbert distance** between any two distinct points $a, b \in \Omega$ is given by

$$d(a, b) = \frac{1}{2} \log \text{CR}[x, a, b, y]$$




Convex Real Projective Structures on $S_{0,3}$

- ▶ Can equip $S_{0,3}$ with many convex real projective structures with their Hilbert metric

Convex Real Projective Structures on $S_{0,3}$

- ▶ Can equip $S_{0,3}$ with many convex real projective structures with their Hilbert metric
- ▶ Use reflections


$$R_{1,T} = \begin{bmatrix} -1 & 0 & 0 \\ 2T & 1 & 0 \\ \frac{2}{T} & 0 & 1 \end{bmatrix} \quad R_{2,T} = \begin{bmatrix} 1 & \frac{2}{T} & 0 \\ 0 & -1 & 0 \\ 0 & 2T & 1 \end{bmatrix}$$
$$R_{3,T} = \begin{bmatrix} 1 & 0 & 2T \\ 0 & 1 & \frac{2}{T} \\ 0 & 0 & -1 \end{bmatrix}$$



Convex Real Projective Structures on $S_{0,3}$

- ▶ Can equip $S_{0,3}$ with many convex real projective structures with their Hilbert metric
- ▶ Use reflections


$$R_{1,T} = \begin{bmatrix} -1 & 0 & 0 \\ 2T & 1 & 0 \\ \frac{2}{T} & 0 & 1 \end{bmatrix} \quad R_{2,T} = \begin{bmatrix} 1 & \frac{2}{T} & 0 \\ 0 & -1 & 0 \\ 0 & 2T & 1 \end{bmatrix}$$
$$R_{3,T} = \begin{bmatrix} 1 & 0 & 2T \\ 0 & 1 & \frac{2}{T} \\ 0 & 0 & -1 \end{bmatrix}$$



Convex Real Projective Structures on $S_{0,3}$

- ▶ Can equip $S_{0,3}$ with many convex real projective structures with their Hilbert metric
- ▶ Use reflections

$$R_{1,T} = \begin{bmatrix} -1 & 0 & 0 \\ 2T & 1 & 0 \\ \frac{2}{T} & 0 & 1 \end{bmatrix} \quad R_{2,T} = \begin{bmatrix} 1 & \frac{2}{T} & 0 \\ 0 & -1 & 0 \\ 0 & 2T & 1 \end{bmatrix}$$
$$R_{3,T} = \begin{bmatrix} 1 & 0 & 2T \\ 0 & 1 & \frac{2}{T} \\ 0 & 0 & -1 \end{bmatrix}$$



Hilbert Entropy

Definition


Let Ω be a convex real projective structure on a surface S and $p \in \Omega$ be a fixed point. The Hilbert **entropy** is given by

$$h_{\Omega} = \lim_{x \rightarrow \infty} \frac{1}{x} \log \# \{ \gamma \in \pi_1(S_{0,3}) : d_{\Omega}(p, \rho_T(\gamma)p) \leq x \}$$

► $\rho_T : \pi_1(S_{0,3}) \rightarrow \mathrm{PSL}_3(\mathbb{R})$

The Manhattan Curve

- ▶ The **Manhattan Curve** is a way use the entropy to compare two convex projective structures.



- ▶ $a + b = 1$

- ▶ $h_{T_1-T_2}(a + b)$

Properties of The Manhattan Curve

- ▶ Strictly convex

Properties of The Manhattan Curve

- ▶ Strictly convex
- ▶ Real analytic

Code to Estimate Entropy

Code to Estimate Entropy

Goals:

- ▶ Generate a lot of group elements

Code to Estimate Entropy

Goals:


- ▶ Generate a lot of group elements
- ▶ Store their singular values
 - ▶ The singular values are related to the quantity $d^H(p, \rho(\gamma))$

Code to Estimate Entropy


Goals:

- ▶ Generate a lot of group elements
- ▶ Store their singular values
 - ▶ The singular values are related to the quantity $d^H(p, \rho(\gamma))$
- ▶ Estimate entropy


Structure of Reflection Group




Structure of Reflection Group





Structure of Reflection Group



Structure of Reflection Group



Structure of Reflection Group



How the Code Works

- ▶ Takes in parameters T and a maximum depth for the layer tree.

How the Code Works

- ▶ Takes in parameters T and a maximum depth for the layer tree.
- ▶ Calculates the length of group element using the singular values.

How the Code Works

- ▶ Takes in parameters T and a maximum depth for the layer tree.
- ▶ Calculates the length of group element using the singular values.
- ▶ Counts number of elements whose lengths are in the interval $(n, n + 1)$.

How the Code Works

- ▶ Takes in parameters T and a maximum depth for the layer tree.
- ▶ Calculates the length of group element using the singular values.
- ▶ Counts number of elements whose lengths are in the interval $(n, n + 1)$.
- ▶ Uses the count to calculate the approximate entropy.

Program Execution

```
>> find_entropy  
finished row: 0  
finished row: 1  
finished row: 2  
finished row: 3  
finished row: 4  
finished row: 5  
finished row: 6  
finished row: 7  
finished row: 8  
finished row: 9  
finished row: 10  
elapsed time: 193.675ms
```

```
>> sum(count_tally(filelist)(2,:))  
ans = 4094
```

Future Directions

- ▶ Analyze symmetries of the Manhattan curve?

Future Directions

- ▶ Analyze symmetries of the Manhattan curve?
- ▶ Estimate the coordinates of the point where the slope of the tangent line of the Manhattan curve is equal to the slope of the secant line between the axes intercepts.

Future Directions

- ▶ Analyze symmetries of the Manhattan curve?
- ▶ Estimate the coordinates of the point where the slope of the tangent line of the Manhattan curve is equal to the slope of the secant line between the axes intercepts.
 - ▶ Dynamical quantity called the **correlation number**.

Future Directions

- ▶ Analyze symmetries of the Manhattan curve?
- ▶ Estimate the coordinates of the point where the slope of the tangent line of the Manhattan curve is equal to the slope of the secant line between the axes intercepts.
 - ▶ Dynamical quantity called the **correlation number**.
- ▶ What happens to the Manhattan curve $\mathcal{M}(\rho_1, \rho_T)$ when T gets large?

Future Directions

- ▶ Analyze symmetries of the Manhattan curve?
- ▶ Estimate the coordinates of the point where the slope of the tangent line of the Manhattan curve is equal to the slope of the secant line between the axes intercepts.
 - ▶ Dynamical quantity called the **correlation number**.
- ▶ What happens to the Manhattan curve $\mathcal{M}(\rho_1, \rho_T)$ when T gets large?
- ▶ Optimize code

Future Directions

- ▶ Analyze symmetries of the Manhattan curve?
- ▶ Estimate the coordinates of the point where the slope of the tangent line of the Manhattan curve is equal to the slope of the secant line between the axes intercepts.
 - ▶ Dynamical quantity called the **correlation number**.
- ▶ What happens to the Manhattan curve $\mathcal{M}(\rho_1, \rho_T)$ when T gets large?
- ▶ Optimize code
- ▶ Generate more examples for different values of T .

Thank You!