

清华大学统计学辅修课程  
Computational Statistics

1

## Lecture 4- Linear Algebra

周在莹  
清华大学统计学研究中心

<http://www.stat.tsinghua.edu.cn>



清华大学统计学研究中心



1

### Outline

2

- ▶ Orthogonality
  - Projection
  - QR Factorization
  - Gram-Schmidt Orthogonalization
  - Householder Triangularization
  - Gaussian Elimination
- ▶ Decomposition
  - Cholesky Factorization/Decomposition
  - Eigenvalue Decomposition
  - Singular Value Decomposition (SVD)
  - Least Squares Problems



清华大学统计学研究中心

2

3

# Projection



清华大学统计学研究中心

3

## Projectors

4


- A projector is an idempotent matrix, i.e. a square matrix  $P$  that satisfies

$$P^2 = P$$

- Physical picture: oblique projector, shine a light onto the subspace  $\text{range}(P)$  from just the direction  $Pv-v$ , then  $Pv$  would be the shadow projected by the vector  $v$

- $Pv-v \in \text{null}(P)$

- Whether or not all the projection lines are orthogonal to the projection plane




清华大学统计学研究中心

4

## Complementary Projectors

5

- ▶ If  $P$  is a projector,  $I - P$  is also a projector, called the complementary projector to  $P$
- ▶  $I - P$  project onto  $\text{range}(I - P)$
- ▶  $\text{range}(I - P) = \text{null}(P)$
- ▶  $\text{range}(P) = \text{null}(I - P)$
- ▶  $\text{null}(I - P) \cap \text{null}(P) = \{0\}$
- ▶  $\text{range}(P) \cap \text{null}(P) = \{0\}$
- ▶  $P$  is the projector onto  $\text{range}(P)$  along  $\text{range}(I - P)$
- ▶ Decomposition of a given vector: Using a projector we can decompose any vector  $v$  into  $v = Pv + (I - P)v$ . This decomposition is unique (Why?)



清华大学统计学研究中心

5


## Orthogonal Projectors

6

- ▶ A projector  $P$  is orthogonal if  $\text{range}(P)$  and  $\text{range}(I - P)$  are orthogonal (NOT orthogonal matrices!)
- ▶ A projector  $P$  is orthogonal if and only if  $P = P^*$

- ▶ In a statistical context,  $P$  is a symmetric matrix
- ▶ Then  $P = QQ^T$ , where the columns of  $Q_{m \times n}$  are orthonormal
- ▶ Denote  $Q = [q_1 | q_2 | \dots | q_n]$ , then


$$P = \sum_{j=1}^n q_j q_j^T, \quad q_j \in R^m$$



清华大学统计学研究中心

6

7




$v \in \mathbb{R}^m$   
 $(I-P)v$   
 $Pv$

$v = Pv + (I-P)v$   
 $P = \sum_{j=1}^n q_j q_j^T, \quad q_j \in \mathbb{R}^m$


- ▶ Rank-one orthogonal projector  $P = qq^T$  isolates the component  $Pv$  in a single direction  $q$ .  $\text{range}(qq^T) = \text{range}(q)$
- ▶ The complement  $P = I - qq^T$  is a rank  $m - 1$  orthogonal projector that eliminates the component  $Pv$  in the direction of  $q$
- ▶  $q$  is a unit vector. For arbitrary nonzero vectors  $a$ :

$$\begin{cases} P = qq^T \\ P_{\perp} = I - qq^T \end{cases}$$

the projection matrix onto  $q$



$$\begin{cases} P = \frac{aa^T}{a^T a} \\ P_{\perp} = I - \frac{aa^T}{a^T a} \end{cases}$$



清华大学统计学研究中心

7

8

## Projection with an Arbitrary Basis

- ▶ Suppose that the subspace is spanned by the linearly independent vectors  $\{a_1, \dots, a_n\}$
- ▶ Let  $A$  be the  $m \times n$  matrix whose  $j$ th column is  $a_j$
- ▶ For  $v$ , consider its orthogonal projection onto  $\text{range}(A)$ , denoted as  $y$
- ▶  $v - y$  must be orthogonal to  $\text{range}(A)$

$a_j^T(v - y) = 0$   
for every  $j$

→

$y = Ax$   
 $a_j^T(v - Ax) = 0$   
for every  $j$

→

$A^T(v - Ax) = 0$


→

$x = (A^T A)^{-1} A^T v$

→

$P = A (A^T A)^{-1} A^T$

$P = \frac{aa^T}{a^T a}$



清华大学统计学研究中心

8

9

# QR Factorization



清华大学统计学研究中心

9

## Reduced QR Factorization

10

- For many applications, we find ourselves interested in the column spaces of a matrix  $A$

$$\langle a_1 \rangle \subseteq \langle a_1, a_2 \rangle \subseteq \langle a_1, a_2, a_3 \rangle \subseteq \dots$$

- The idea of QR factorization is the construction of a sequence of orthonormal vectors  $q_1, q_2, \dots$  that span these successive spaces
- This amounts to

$$A = \hat{Q}\hat{R} \quad [a_1 | a_2 | \dots | a_n] = [q_1 | q_2 | \dots | q_n] \begin{bmatrix} r_{11} & r_{12} & \dots & r_{1n} \\ & r_{22} & \dots & r_{2n} \\ & & \ddots & \vdots \\ & & & r_{nn} \end{bmatrix}, r_{jj} \neq 0$$

when  $A_{m \times n}$  has full rank  $n$



清华大学统计学研究中心

10

## Full QR Factorization

11

- ▶  $Q_{m \times n} = [q_1 | q_2 | \dots]$ , appending additional  $m - n$  orthonormal columns to make it unitary
- ▶ So rows of zeros should be appended to  $R$  so that it becomes taller, still upper-triangular

$$A_{m \times n} = \hat{Q}_{m \times n} Q_{m \times m} \hat{R}_{m \times n}$$

The added columns span a space orthogonal to  $\text{range}(A)$ , constitute an orthonormal basis for  $\text{null}(A^T)$



清华大学统计学研究中心

11

## Gram-Schmidt Orthogonalization

12

$$\begin{bmatrix} | & | & & | \\ a_1 & a_2 & \dots & a_n \\ | & | & & | \end{bmatrix} = \begin{bmatrix} | & | & & | \\ q_1 & q_2 & \dots & q_n \\ | & | & & | \end{bmatrix} \begin{bmatrix} r_{11} & r_{12} & \dots & r_{1n} \\ & r_{22} & \dots & r_{2n} \\ & & \ddots & \vdots \\ & & & r_{nn} \end{bmatrix},$$

$r_{jj} \neq 0$

$$\begin{aligned} q_1 &= \frac{a_1}{r_{11}}, \\ q_2 &= \frac{a_2 - r_{12}q_1}{r_{22}}, \\ q_3 &= \frac{a_3 - r_{13}q_1 - r_{23}q_2}{r_{33}}, \\ &\vdots \\ q_n &= \frac{a_n - \sum_{i=1}^{n-1} r_{in}q_i}{r_{nn}}. \end{aligned}$$

▶ where

$$\begin{aligned} r_{ij} &= q_i^T a_j \quad (i \neq j), \\ r_{jj} &= \left\| a_j - \sum_{i=1}^{j-1} r_{ij}q_i \right\|_2. \end{aligned}$$



清华大学统计学研究中心

12

## Algorithm of Classical Gram-Schmidt (Unstable)

13

```

for  $j = 1$  to  $n$ 
   $v_j = a_j$ 
  for  $i = 1$  to  $j - 1$ 
     $r_{ij} = q_i^T a_j$ 
     $v_j = v_j - r_{ij} q_i$ 
   $r_{jj} = \|v_j\|_2$ 
   $q_j = v_j / r_{jj}$ 

```

- ▶ Mathematically, it offers a simple route to understanding and proving various properties of QR factorizations
- ▶ Numerically, it turns out to be unstable because of rounding errors on a computer.
- ▶ To emphasize the instability, numerical analysts refer to this as the **classical** Gram-Schmidt iteration, as opposed to the **modified** Gram-Schmidt iteration, discussed later



清华大学统计学研究中心

13

## Existence and Uniqueness

14

- ▶ All matrices have QR factorizations, and under suitable restrictions, they are unique
- ▶ Theorem 1. Every  $m \times n$  matrix  $A$  ( $m \geq n$ ) has a full QR factorization, hence also a reduced QR factorization
- ▶ Theorem 2. Each  $m \times n$  matrix  $A$  ( $m \geq n$ ) of full rank has a unique **reduced** QR factorization  $A = \hat{Q}\hat{R}$  with  $r_{jj} > 0$



清华大学统计学研究中心

14

## Solution of $Ax = b$ by QR Factorization

15

- ▶ Suppose we wish to solve  $Ax = b$  for  $x$ , where  $m \times m$  matrix  $A$  is nonsingular
- ▶ If  $A = QR$  is a QR factorization, then we can write  $QRx = b$ , or  $Rx = Q^T b$
- ▶ Method for computing the solution to  $Ax = b$ :
  1. Compute a QR factorization  $A = QR$
  2. Compute  $y = Q^T b$
  3. Solve  $Rx = y$  for  $x$
- ▶ However, Gaussian elimination is the algorithm generally used in practice, since it requires only half as many numerical operations



清华大学统计学研究中心

15

## Gram-Schmidt Orthogonalization

16



清华大学统计学研究中心

16



## Inner Products and Orthogonality

17

- ▶ The most important idea to draw from these concepts is: inner products can be used to decompose arbitrary vectors into orthogonal components

- ▶ For example, suppose that  $\{q_1, q_2, \dots, q_n\}$  is an orthonormal set, and let  $v$  be an arbitrary vector. Then

$$r = v - (q_1^* v)q_1 - (q_2^* v)q_2 - \dots - (q_n^* v)q_n = (I - \sum q_i q_i^*)v$$

is orthogonal to  $\{q_1, q_2, \dots, q_n\}$ .

Thus we see that  $v$  can be decomposed into  $n + 1$  orthogonal components:  $v = r + \sum q_i q_i^* v$

- ▶ Interpretations: View  $v$  as a sum of coefficients  $q_i^* v$  times vectors  $q_i$ ; View  $v$  as a sum of orthogonal projections of  $v$  onto the various directions  $q_i$ . The  $i$ th projection operation is achieved by the rank-one matrix  $q_i q_i^*$



清华大学统计学研究中心

17

## Gram-Schmidt Projections

18

- ▶ Use orthogonal projectors to describe the G-S iteration in another way
- ▶ Suppose that  $m \times n$  matrix  $A$  ( $m \geq n$ ) has full rank with columns  $\{a_j\}$
- ▶ Let  $Q_{j-1}$  denote the  $m \times (j-1)$  matrix containing the first  $j-1$  columns of  $Q$ ,  $Q_{j-1} = [q_1 \mid \dots \mid q_{j-1}]$

$$P_j = I - Q_{j-1} Q_{j-1}^T$$

is the  $m \times m$  matrix of rank  $m - (j-1)$  that projects  $R^m$  orthogonally onto the space orthogonal to  $\langle q_1, \dots, q_{j-1} \rangle$ ,  $j = 2, \dots, n$ ,  $P_1 = I$

- ▶  $P_j$  is the  $m \times m$  matrix of rank  $m - (j-1)$  that projects  $R^m$  orthogonally onto the space orthogonal to  $\langle q_1, \dots, q_{j-1} \rangle$ ,  $j = 2, \dots, n$ ,  $P_1 = I$

$$q_1 = \frac{P_1 a_1}{\|P_1 a_1\|}, \quad q_2 = \frac{P_2 a_2}{\|P_2 a_2\|}, \dots, \quad q_n = \frac{P_n a_n}{\|P_n a_n\|}$$

- ▶  $q_j$  is orthogonal to  $q_1, \dots, q_{j-1}$ , lies in the space  $\langle a_1, \dots, a_j \rangle$ , and has norm 1
- ▶  $v_j = P_j a_j = a_j - (q_1 q_1^T) a_j - \dots - (q_{j-1} q_{j-1}^T) a_j = a_j - (q_1^T a_j) q_1 - \dots - (q_{j-1}^T a_j) q_{j-1}$

for  $j = 1$  to  $n$   
 $v_j = a_j$   
 for  $i = 1$  to  $j-1$   
 $r_{ij} = q_i^T a_j$   
 $v_j = v_j - r_{ij} q_i$   
 $r_{jj} = \|v_j\|$   
 $q_j = v_j / r_{jj}$



清华大学统计学研究中心

18

## Modified Gram-Schmidt Algorithm

19

- Compute the same result by a sequence of  $j - 1$  projections of rank  $m - 1$

$$P_j = P_{\perp q_{j-1}} \dots P_{\perp q_2} P_{\perp q_1}$$

$$v_j^{(1)} = a_j,$$

$$v_j = P_j a_j$$

$$v_j^{(2)} = P_{\perp q_1} v_j^{(1)} = v_j^{(1)} - q_1 q_1^T v_j^{(1)},$$

$$v_j^{(3)} = P_{\perp q_2} v_j^{(2)} = v_j^{(2)} - q_2 q_2^T v_j^{(2)},$$

$$\vdots$$

$$v_j = v_j^{(j)} = P_{\perp q_{j-1}} v_j^{(j-1)} = v_j^{(j-1)} - q_{j-1} q_{j-1}^T v_j^{(j-1)}.$$

for  $i = 1$  to  $n$

$$v_i = a_i$$

for  $i = 1$  to  $n$

$$r_{ii} = \|v_i\|$$

$$q_i = v_i / r_{ii}$$

for  $j = i + 1$  to  $n$

$$r_{ij} = q_i^T v_j$$

$$v_j = v_j - r_{ij} q_i$$



清华大学统计学研究中心

19

## Comparison

20

### Classical G-S

for  $j = 1$  to  $n$

$$v_j = a_j$$

for  $i = 1$  to  $j - 1$

$$r_{ij} = q_i^T a_j$$

$$v_j = v_j - r_{ij} q_i$$

$$r_{jj} = \|v_j\|$$

$$q_j = v_j / r_{jj}$$

### Modified G-S

for  $i = 1$  to  $n$

$$v_i = a_i$$

for  $i = 1$  to  $n$

$$r_{ii} = \|v_i\|$$

$$q_i = v_i / r_{ii}$$

for  $j = i + 1$  to  $n$

$$r_{ij} = q_i^T v_j$$

$$v_j = v_j - r_{ij} q_i$$

### C/M G-S

for  $j = 1$  to  $n$

$$v_j = a_j$$

for  $i = 1$  to  $j - 1$

$$r_{ij} = q_i^T a_j \text{ (CGS)}$$

$$\left\{ \begin{array}{l} r_{ij} = q_i^T v_j \text{ (MGS)} \\ v_j = v_j - r_{ij} q_i \end{array} \right.$$

$$r_{jj} = \|v_j\|$$

$$q_j = v_j / r_{jj}$$

- Modified G-S is numerically stable (less sensitive to rounding errors)



清华大学统计学研究中心

20

## Example: Classical vs. Modified Gram-Schmidt

21

- Compare classical and modified G-S for the vectors

$$a_1 = (1, \varepsilon, 0, 0)^T, a_2 = (1, 0, \varepsilon, 0)^T, a_3 = (1, 0, 0, \varepsilon)^T$$

making the approximation  $1 + \varepsilon^2 \approx 1$

- Classical:

$$\triangleright v_1 \leftarrow (1, \varepsilon, 0, 0)^T, r_{11} = \sqrt{1 + \varepsilon^2} \approx 1, q_1 = v_1 / r_{11} = (1, \varepsilon, 0, 0)^T$$

$$\triangleright v_2 \leftarrow (1, 0, \varepsilon, 0)^T, r_{12} = q_1^T a_2 = 1, v_2 \leftarrow v_2 - r_{12} q_1 = (0, -\varepsilon, \varepsilon, 0)^T$$

$$r_{22} = \sqrt{2}\varepsilon, q_2 = v_2 / r_{22} = (0, -1, 1, 0)^T / \sqrt{2}$$

$$\triangleright v_3 \leftarrow (1, 0, 0, \varepsilon)^T, r_{13} = q_1^T a_3 = 1, v_3 \leftarrow v_3 - r_{13} q_1 = (0, -\varepsilon, 0, \varepsilon)^T$$

$$r_{23} = q_2^T a_3 = 0, v_3 \leftarrow v_3 - r_{23} q_2 = (0, -\varepsilon, 0, \varepsilon)^T, r_{33} = \sqrt{2}\varepsilon,$$

$$q_3 = v_3 / r_{33} = (0, -1, 0, 1)^T / \sqrt{2}$$



清华大学统计学研究中心

21

22

- Modified:

$$\triangleright v_1 \leftarrow (1, \varepsilon, 0, 0)^T, r_{11} = \sqrt{1 + \varepsilon^2} \approx 1, q_1 = v_1 / r_{11} = (1, \varepsilon, 0, 0)^T$$

$$\triangleright v_2 \leftarrow (1, 0, \varepsilon, 0)^T, r_{12} = q_1^T v_2 = 1, v_2 \leftarrow v_2 - r_{12} q_1 = (0, -\varepsilon, \varepsilon, 0)^T$$

$$r_{22} = \sqrt{2}\varepsilon, q_2 = v_2 / r_{22} = (0, -1, 1, 0)^T / \sqrt{2}$$

$$\triangleright v_3 \leftarrow (1, 0, 0, \varepsilon)^T, r_{13} = q_1^T v_3 = 1, v_3 \leftarrow v_3 - r_{13} q_1 = (0, -\varepsilon, 0, \varepsilon)^T$$

$$r_{23} = q_2^T v_3 = \varepsilon / \sqrt{2}, v_3 \leftarrow v_3 - r_{23} q_2 = (0, -\varepsilon/2, -\varepsilon/2, \varepsilon)^T, r_{33} = \sqrt{6}\varepsilon/2,$$

$$q_3 = v_3 / r_{33} = (0, -1, -1, 2)^T / \sqrt{6}$$

- Check orthogonality:

$$\triangleright \text{Classical: } q_2^T q_3 = (0, -1, 1, 0)(0, -1, 0, 1)^T / 2 = 1/2$$

$$\triangleright \text{Modified: } q_2^T q_3 = (0, -1, 1, 0)(0, -1, -1, 2)^T / \sqrt{12} = 0$$



清华大学统计学研究中心

22

## Operation Count

23

- ▶ Count number of floating points operations – “flops” – in an algorithm
- ▶ Each  $+$ ,  $-$ ,  $\times$ ,  $/$ , or  $\sqrt{\quad}$  counts as one flop
- ▶ No distinction between real and complex
- ▶ No consideration of memory accesses or other performance aspects



清华大学统计学研究中心

23

## Operation Count - Modified G-S

24

- ▶ Example: Count all  $+$ ,  $-$ ,  $\times$ ,  $/$  in the Modified Gram-Schmidt algorithm (not just the leading term)

- |     |                                      |  |
|-----|--------------------------------------|--|
| (1) | <b>for</b> $i = 1$ <b>to</b> $n$     |  |
| (2) | $v_i = a_i$                          |  |
| (3) | <b>for</b> $i = 1$ <b>to</b> $n$     |  |
| (4) | $r_{ii} =   v_i  $                   | $m$ multiplications, $m - 1$ additions |
| (5) | $q_i = v_i / r_{ii}$                 | $m$ divisions                          |
| (6) | <b>for</b> $j = i + 1$ <b>to</b> $n$ |  |
| (7) | $r_{ij} = q_i^T v_j$                 | $m$ multiplications, $m - 1$ additions |
| (8) | $v_j = v_j - r_{ij} q_i$             | $m$ multiplications, $m$ subtractions  |



清华大学统计学研究中心

24

- The total for each operation is

25

$$\begin{aligned}\# A &= \sum_{i=1}^n \left( m-1 + \sum_{j=i+1}^n m-1 \right) = n(m-1) + \sum_{i=1}^n (m-1)(n-i) \\ &= n(m-1) + \frac{n(n-1)(m-1)}{2} = \frac{1}{2} n(n+1)(m-1)\end{aligned}$$

$$\# S = \sum_{i=1}^n \sum_{j=i+1}^n m = \sum_{i=1}^n m(n-i) = \frac{1}{2} mn(n-1)$$

$$\begin{aligned}\# M &= \sum_{i=1}^n \left( m + \sum_{j=i+1}^n 2m \right) = mn + \sum_{i=1}^n 2m(n-i) \\ &= mn + \frac{2mn(n-1)}{2} = mn^2\end{aligned}$$

$$\# D = \sum_{i=1}^n m = mn$$



清华大学统计学研究中心

25

- And the total flop count is

26

$$\frac{1}{2} n(n+1)(m-1) + \frac{1}{2} mn(n-1) + mn^2 + mn$$

$$= 2mn^2 + mn - \frac{1}{2} n^2 - \frac{1}{2} n \sim 2mn^2$$

- The symbol  $\sim$  indicates asymptotic value as  $m, n \rightarrow \infty$  (leading term)

- Easier to find just the leading term:

– Most work done in lines (7) and (8), with  $4m$  flops per iteration

– Including the loops, the total becomes

$$\sum_{i=1}^n \sum_{j=i+1}^n 4m = 4m \sum_{i=1}^n (n-i) \sim 4m \sum_{i=1}^n i \sim 2mn^2$$



清华大学统计学研究中心

26

27

# Householder Triangularization



清华大学统计学研究中心

27

28

## Gram-Schmidt as Triangular Orthogonalization

- ▶ Gram-Schmidt multiplies with triangular matrices to make columns orthogonal, for example at the first step:

$$\begin{bmatrix} | & | & | & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & | & | \end{bmatrix} \begin{bmatrix} 1 & -r_{12} & \cdots & -r_{1n} \\ r_{11} & r_{11} & & r_{11} \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{bmatrix} = \begin{bmatrix} | & | & | & | \\ q_1 & v_2^{(2)} & \cdots & v_n^{(2)} \\ | & | & | & | \end{bmatrix}$$

- ▶ After all the steps we get a product of triangular matrices

$$\underbrace{AR_1R_2\cdots R_n}_{R^{-1}} = Q$$

- ▶ Triangular orthogonalization



清华大学统计学研究中心

28

## Householder Triangularization

29

- The Householder method multiplies by unitary matrices to make columns triangular, for example at the first step of :

$$Q_1 A = \begin{bmatrix} r_{11} & \times & \cdots & \times \\ 0 & \times & \cdots & \times \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \times & \cdots & \times \end{bmatrix}$$

- After all the steps we get a product of orthogonal matrices matrices

$$\underbrace{Q_n \cdots Q_2 Q_1}_{Q^{-1}} A = R$$

- Orthogonal triangularization

- Numerically more stable than Gram-Schmidt orthogonalization.
- Principal method for computing  $QR$  factorizations.



清华大学统计学研究中心

29

## Introducing Zeros

30

- $Q_k$  introduces zeros below the diagonal in column  $k$
- Preserves all the zeros previously introduced

$$\begin{array}{c} \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} \xrightarrow{Q_1} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \end{bmatrix} \xrightarrow{Q_2} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & 0 & \times \\ 0 & 0 & \times \end{bmatrix} \xrightarrow{Q_3} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & 0 & \times \\ 0 & 0 & 0 \end{bmatrix} \\ A \qquad \qquad Q_1 A \qquad \qquad Q_2 Q_1 A \qquad \qquad Q_3 Q_2 Q_1 A \end{array}$$



清华大学统计学研究中心

30

## Householder Reflectors

31

- Let  $Q_k$  be of the form

$$Q_k = \begin{bmatrix} I & 0 \\ 0 & F \end{bmatrix}$$

where  $I$  is  $(k-1) \times (k-1)$  and  $F$  is  $(m-k+1) \times (m-k+1)$

- Create Householder reflector  $F$  that introduces zeros:

$$x = \begin{bmatrix} \times \\ \times \\ \vdots \\ \times \end{bmatrix} \in R^{m-k+1} \xrightarrow{F} Fx = \begin{bmatrix} \|x\| \\ 0 \\ \vdots \\ 0 \end{bmatrix} = \|x\|e_1$$



清华大学统计学研究中心

31

- Idea: Reflect across hyperplane  $H$  orthogonal to  $v = \|x\|e_1 - x$ , by the unitary matrix

32

unitary/orthogonal,  
full rank( $m-k+1$ )

hermitian/symmetric


- Compare with projector

$$F = I - 2 \frac{vv^T}{v^T v}$$

so is  $Q_k$

$$P_{\perp v} = I - \frac{vv^T}{v^T v}$$

rank  $m-1$



清华大学统计学研究中心

32



## Choice of Reflector

33


- The vector  $x$  can be reflected to any multiple  $z$  of  $\|x\|e_1$  with  $|z| = 1$

- Better numerical properties with large  $\|v\|$ , for example

$$v = \text{sign}(x_1) \|x\|e_1 + x$$

- Note:

Here  $\text{sign}(0) = 1$



清华大学统计学研究中心

33

## The Householder Algorithm

34

- Compute the factor  $R$  of a QR factorization of a given  $m \times n$  matrix  $A$  ( $m \geq n$ )
- Leave result in place of  $A$ , store reflection vectors  $v_k$  for later use

### Householder QR Factorization

for  $k = 1$  to  $n$

$$x = A_{k:m,k}$$

$$v_k = \text{sign}(x_1) \|x\|e_1 + x$$

$$v_k = v_k / \|v_k\|$$

$$A_{k:m,k:n} = A_{k:m,k:n} - 2v_k(v_k^T A_{k:m,k:n})$$



清华大学统计学研究中心

34

## Applying or Forming $Q$

35

- In solving  $Ax = b$ , compute  $Q^{-1}b = Q_n \cdots Q_2 Q_1 b$  implicitly

$$\begin{array}{l} \text{Implicit Calculation of } Q^{-1}b \\ \text{for } k = 1 \text{ to } n \\ b_{k:m} = b_{k:m} - 2v_k(v_k^T b_{k:m}) \end{array}$$

- Compute  $Qx = Q_1 Q_2 \cdots Q_n x$  implicitly

$$\begin{array}{l} \text{Implicit Calculation of } Qx \\ \text{for } k = n \text{ down to } 1 \\ x_{k:m} = x_{k:m} - 2v_k(v_k^T x_{k:m}) \end{array}$$

- To create  $Q$  explicitly, apply to  $x = e_1, \dots, e_m$   
(or just  $x = e_1, \dots, e_n$  for reduced  $Q$ )



清华大学统计学研究中心

35

## Operation Count - Householder QR

36

- Most work is done by

$$A_{k:m,k:n} = A_{k:m,k:n} - 2v_k(v_k^T A_{k:m,k:n})$$

- Operations per iteration:

- $2(m-k)(n-k)$  for the dot products  $v_k^T A_{k:m,k:n}$
- $(m-k)(n-k)$  for the outer product  $2v_k(\dots)$
- $(m-k)(n-k)$  for the subtraction  $A_{k:m,k:n} - \dots$
- $4(m-k)(n-k)$  total

- Including the outer loop, the total becomes

$$\begin{aligned} \sum_{k=1}^n 4(m-k)(n-k) &= 4 \sum_{k=1}^n (mn - k(m+n) + k^2) \\ &\sim 4mn^2 - 4(m+n)n^2/2 + 4n^3/3 = 2mn^2 - 2n^3/3 \end{aligned}$$



清华大学统计学研究中心

36

## Givens Rotations

37

- Alternative to Householder reflectors

- A Givens rotation

$$R = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

rotates  $x \in \mathbb{R}^2$  by  $\theta$

- To set an element to zero, choose  $\cos \theta$  and  $\sin \theta$  so that

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x_i \\ x_j \end{bmatrix} = \begin{bmatrix} \sqrt{x_i^2 + x_j^2} \\ 0 \end{bmatrix}$$

or

$$\cos \theta = \frac{x_i}{\sqrt{x_i^2 + x_j^2}}, \quad \sin \theta = \frac{-x_j}{\sqrt{x_i^2 + x_j^2}}$$



清华大学统计学研究中心

37

## Givens QR

38

- Introduce zeros in column from bottom and up

$$\begin{array}{ccc} \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} & \xrightarrow{(3,4)} & \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ 0 & \times & \times \end{bmatrix} & \xrightarrow{(2,3)} & \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ 0 & \times & \times \\ \times & \times & \times \end{bmatrix} & \xrightarrow{(1,2)} & \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ 0 & \times & \times \\ \times & \times & \times \end{bmatrix} \\ \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} & \xrightarrow{(3,4)} & \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} & \xrightarrow{(2,3)} & \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} & \xrightarrow{(3,4)} & R \end{array}$$

- Flop count  $3mn^2 - n^3$  (50% more than Householder QR)



清华大学统计学研究中心

38

39

# Gaussian Elimination



清华大学统计学研究中心

39

## LU Factorization

40

- Gaussian elimination transforms a full linear system into an upper-triangular one by applying simple linear transformations on the left

- For  $A_{m \times m}$ ,  $L_{m-1} \dots L_2 L_1 A = U$ ,

$$A = LU$$

$$\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ \times & \times & \times & \times \\ \times & \times & \times & \times \end{bmatrix} \xrightarrow{L_1} \begin{bmatrix} \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & \times & \times & \times \end{bmatrix} \xrightarrow{L_2} \begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & \times & \times & \times \end{bmatrix} \xrightarrow{L_3} \begin{bmatrix} \times & \times & \times & \times \\ & \times & \times & \times \\ & & \times & \times \\ & & 0 & \times \end{bmatrix}$$

$A$                        $L_1 A$                        $L_2 L_1 A$                        $L_3 L_2 L_1 A$

- Triangular triangularization



清华大学统计学研究中心

40

## Two Strokes of Luck

41

► Example

$$A = \begin{bmatrix} 2 & 1 & 1 & 0 \\ 4 & 3 & 3 & 1 \\ 8 & 7 & 9 & 5 \\ 6 & 7 & 9 & 8 \end{bmatrix}$$

$$L_1 A = \begin{bmatrix} 1 & & & \\ -2 & 1 & & \\ -4 & & 1 & \\ -3 & & & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 & 0 \\ 4 & 3 & 3 & 1 \\ 8 & 7 & 9 & 5 \\ 6 & 7 & 9 & 8 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 & 0 \\ 1 & 1 & 1 & \\ 3 & 5 & 5 & \\ 4 & 6 & 8 & \end{bmatrix}$$

$$L_2 L_1 A = \begin{bmatrix} 1 & & & \\ & 1 & & \\ -3 & & 1 & \\ -4 & & & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 & 0 \\ & 1 & 1 & 1 \\ 3 & 5 & 5 & \\ 4 & 6 & 8 & \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 & 0 \\ & 1 & 1 & 1 \\ & & 2 & 2 \\ & & & 2 & 4 \end{bmatrix}$$



清华大学统计学研究中心

41

$$L_3 L_2 L_1 A = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 & 0 \\ & 1 & 1 & 1 \\ & & 2 & 2 \\ & & & 2 & 4 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 & 0 \\ & 1 & 1 & 1 \\ & & 2 & 2 \\ & & & 2 \end{bmatrix} = U$$

►  $L_k^{-1}$  is obtained by negating its subdiagonal entries

$$\begin{bmatrix} 1 & & & \\ -2 & 1 & & \\ -4 & & 1 & \\ -3 & & & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & & & \\ 2 & 1 & & \\ 4 & & 1 & \\ 3 & & & 1 \end{bmatrix}$$

►  $L = L_1^{-1} L_2^{-1} L_3^{-1}$  is just the unit lower-triangular matrix with their nonzero subdiagonal entries inserted in the appropriate places

$$\begin{bmatrix} 2 & 1 & 1 & 0 \\ 4 & 3 & 3 & 1 \\ 8 & 7 & 9 & 5 \\ 6 & 7 & 9 & 8 \end{bmatrix} = \begin{bmatrix} 1 & & & \\ 2 & 1 & & \\ 4 & 3 & 1 & \\ 3 & 4 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 & 0 \\ & 1 & 1 & 1 \\ & & 2 & 2 \\ & & & 2 \end{bmatrix}$$



清华大学统计学研究中心

A

L

U

42

## General Formulas

43

- Suppose  $x_k$  denotes the  $k$ th column of the matrix at the beginning of step  $k$

$$x_k = \begin{bmatrix} x_{1k} \\ \vdots \\ x_{kk} \\ x_{k+1,k} \\ \vdots \\ x_{mk} \end{bmatrix} \xrightarrow{L_k} L_k x_k = \begin{bmatrix} x_{1k} \\ \vdots \\ x_{kk} \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

- To do this, we subtract  $l_{jk}$  times row  $k$  from row  $j$ , where  $l_{jk}$  is the multiplier

$$l_{jk} = \frac{x_{jk}}{x_{kk}} \quad (k < j \leq m)$$

- The matrix  $L_k$  takes the form

$$L_k = \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & -l_{k+1,k} & 1 & \\ & & \vdots & & \ddots \\ & & -l_{mk} & & & 1 \end{bmatrix}$$



清华大学统计学研究中心

43

## Explain the Good Fortune

44

- Define  $l_k = [0, \dots, 0, l_{k+1,k}, \dots, l_{m,k}]^T$ , then  $L_k = I - l_k e_k^T$

- The inverse of  $L_k$  is  $I + l_k e_k^T$

$$(I - l_k e_k^T)(I + l_k e_k^T) = I - l_k e_k^T l_k e_k^T = I$$

- For

$$L = L_1^{-1} L_2^{-1} \cdots L_{m-1}^{-1} = \begin{bmatrix} 1 & & & & \\ l_{21} & 1 & & & \\ l_{31} & l_{32} & 1 & & \\ \vdots & \vdots & \ddots & \ddots & \\ l_{m1} & l_{m2} & \cdots & l_{m,m-1} & 1 \end{bmatrix}$$

$$\begin{aligned} L_k^{-1} L_{k+1}^{-1} &= (I + l_k e_k^T)(I + l_{k+1} e_{k+1}^T) \\ &= I + l_k e_k^T + l_{k+1} e_{k+1}^T \end{aligned}$$



清华大学统计学研究中心

44

## Algorithm

45

### Gaussian Elimination

```

 $U = A, L = I$ 
for  $k = 1$  to  $m - 1$ 
  for  $j = k + 1$  to  $m$ 
     $l_{jk} = u_{jk}/u_{kk}$ 
     $u_{j,k:m} = u_{j,k:m} - l_{jk}u_{k,k:m}$ 

```

#### ► Operation count

$$u_{j,k:m} = u_{j,k:m} - l_{jk}u_{k,k:m}$$

$$\sum_{k=1}^{m-1} 2(m-k+1)(m-k) \sim \frac{2}{3}m^3 \text{ flops}$$



清华大学统计学研究中心

45

## Solution of $Ax = b$ by LU Factorization

46

- $Ax = b$  is reduced to  $LUx = b$ .
- Solve two triangular systems:
  - first  $Ly = b$  for the unknown  $y$  (forward substitution)
  - then  $Ux = y$  for the unknown  $x$  (back substitution)

#### ► Operation count

$$\sum_{k=1}^m 2k \sim m^2 \text{ flops}$$

$$l_{k1}y_1 + \dots + l_{kk}y_k = b_k$$

- So if Gaussian elimination is used, the total work is  $\sim 2/3m^3$  flops, half the figure of  $\sim 4/3m^3$  flops for a solution by Householder triangularization
- Usually used rather than QR factorization to solve square systems of equations



清华大学统计学研究中心

46

## Instability of Gaussian Elimination

47

- Consider

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$$

This matrix has full rank and is well-conditioned. Nevertheless, Gaussian elimination fails at the first step

- A slight perturbation of the same matrix reveals the more general problem.

$$A = \begin{bmatrix} 10^{-20} & 1 \\ 1 & 1 \end{bmatrix} \quad \tilde{L}\tilde{U} = \begin{bmatrix} 10^{-20} & 1 \\ 1 & 0 \end{bmatrix}$$

$$L = \begin{bmatrix} 1 & 0 \\ 10^{20} & 1 \end{bmatrix} \quad U = \begin{bmatrix} 10^{-20} & 1 \\ 0 & 1 - 10^{20} \end{bmatrix} \quad \tilde{U} = \begin{bmatrix} 10^{-20} & 1 \\ 0 & -10^{20} \end{bmatrix}$$



清华大学统计学研究中心

47

## Pivoting

48

- The instability can be controlled by permuting the order of the rows of the matrix being operated on, an operation called pivoting
- Pivoting has been a standard feature of Gaussian elimination computations since the 1950s

$$\text{pivot} \begin{bmatrix} \times & \times & \times & \times & \times \\ & \boxed{x_{kk}} & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \end{bmatrix} \rightarrow \begin{bmatrix} \times & \times & \times & \times & \times \\ & x_{kk} & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \end{bmatrix}$$



清华大学统计学研究中心

48



49

$$\begin{bmatrix} \times & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ \textcolor{red}{x}_{ik} & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \end{bmatrix} \rightarrow \begin{bmatrix} \times & \times & \times & \times & \times \\ & 0 & \times & \times & \times \\ & 0 & \times & \times & \times \\ \textcolor{red}{x}_{ik} & \times & \times & \times & \times \\ & 0 & \times & \times & \times \end{bmatrix}$$

$$\begin{bmatrix} \times & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ & \times & \textcolor{red}{x}_{ij} & \times & \times \\ \times & \times & \times & \times & \times \end{bmatrix} \rightarrow \begin{bmatrix} \times & \times & \times & \times & \times \\ & \times & 0 & \times & \times \\ & \times & 0 & \times & \times \\ & \times & \textcolor{red}{x}_{ij} & \times & \times \\ & \times & 0 & \times & \times \end{bmatrix}$$

- Free to choose any nonzero entry of  $X_{k:m,k:m}$  as the pivot
- In practice, it is common to pick the largest possible
- At step  $k$ , rows/columns of the working matrix shall be permuted so as to move the pivot into the  $(k, k)$  position



清华大学统计学研究中心

49

50

## Partial Pivoting

- Only rows are interchanged

$$\begin{bmatrix} \times & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ \textcolor{red}{x}_{ik} & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \end{bmatrix} \xrightarrow{P_1} \begin{bmatrix} \times & \times & \times & \times & \times \\ \textcolor{red}{x}_{ik} & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ & \times & \times & \times & \times \end{bmatrix} \xrightarrow{L_1} \begin{bmatrix} \times & \times & \times & \times & \times \\ \textcolor{red}{x}_{ik} & \times & \times & \times & \times \\ & 0 & \times & \times & \times \\ & 0 & \times & \times & \times \\ & 0 & \times & \times & \times \end{bmatrix}$$

Pivot selection

Row interchange

Elimination

- $L_{m-1}P_{m-1}\dots L_2P_2L_1P_1A = U$



清华大学统计学研究中心

50

## $PA = LU$ Factorization and a Third Stroke of Luck

51

►  $L_{m-1}P_{m-1}\dots L_2P_2L_1P_1A = U$

$L_{m-1}P_{m-1}\dots L_2P_2L_1P_1 = (L_{m-1}'\dots L_2'L_1')(P_{m-1}\dots P_2P_1) = LP$

► For example,

$$L_3P_3L_2P_2L_1P_1 = L_3'L_2'L_1'P_3P_2P_1$$

with  $L_3' = L_3$ ,  $L_2' = P_3L_2P_3^{-1}$ ,  $L_1' = P_3P_2L_1P_2^{-1}P_3^{-1}$

$$L_k = \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & -l_{k+1,k} & 1 & \\ & & \vdots & & \ddots \\ & & -l_{mk} & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & -l_{mk} & 1 & \\ & & \vdots & & \ddots \\ & & -l_{k+1,k} & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & -l_{mk} & 1 & \\ & & \vdots & & \ddots \\ & & -l_{k+1,k} & & 1 \end{bmatrix}$$

$$L_k' = P_{m-1}\dots P_{k+1}L_kP_{k+1}^{-1}\dots P_{m-1}^{-1}$$



清华大学统计学研究中心

51

## Algorithm

52

►  $PA = LU$

►

### Gaussian Elimination with Partial Pivoting

$U = A$ ,  $L = I$ ,  $P = I$

for  $k = 1$  to  $m - 1$

    Select  $i \geq k$  to maximize  $|u_{ik}|$

$u_{k,k:m} \leftrightarrow u_{i,k:m}$  (interchange two rows)

$l_{k,1:k-1} \leftrightarrow l_{i,1:k-1}$

$P_{k,:} \leftrightarrow P_{i,:}$

    for  $j = k + 1$  to  $m$

$l_{jk} = u_{jk}/u_{kk}$

$u_{j,k:m} = u_{j,k:m} - l_{jk}u_{k,k:m}$



清华大学统计学研究中心

52

## Complete Pivoting

53

- ▶ Each elimination step with a permutation  $P_k$  of the rows applied on the left, a permutation  $Q_k$  of the columns applied on the right:

$$L_{m-1}P_{m-1}\dots L_2P_2L_1P_1 A Q_1Q_2\dots Q_{m-1} = U$$

$$(L_{m-1}'\dots L_2'L_1')(P_{m-1}\dots P_2P_1) A (Q_1Q_2\dots Q_{m-1}) = U$$

- ▶  $PAQ = LU$
- ▶ The selection of pivots takes a significant amount of time
- ▶ In practice this is rarely done, because the improvement in stability is marginal



清华大学统计学研究中心

53

54

## Fundamentals: Matrix-Vector Multiplication



清华大学统计学研究中心

54

## A Matrix Times a Vector

55

$$b = Ax = \sum_{j=1}^n x_j a_j$$

$$\begin{bmatrix} b \end{bmatrix} = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 \begin{bmatrix} a_1 \end{bmatrix} + x_2 \begin{bmatrix} a_2 \end{bmatrix} + \cdots + x_n \begin{bmatrix} a_n \end{bmatrix}$$

►  $b$  is expressed as a linear combination of the columns  $a_j$



清华大学统计学研究中心

55

## Example: Vandermonde Matrix

56

$$p(x) = c_0 + c_1 x + c_2 x^2 + \cdots + c_{n-1} x^{n-1}$$

$$A = \begin{bmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & x_m & x_m^2 & \cdots & x_m^{n-1} \end{bmatrix}$$

$$c = \begin{bmatrix} c_0 \\ c_1 \\ \vdots \\ c_{n-1} \end{bmatrix}$$

$(Ac)_i = p(x_i)$  gives  
the sampled  
polynomial values

►  $A$  is a linear map from vectors of coefficients of polynomials  $p$  of degree  $< n$  to vectors  $(p(x_1), p(x_2), \dots, p(x_m))$  of sampled polynomial values

$$A = \begin{bmatrix} 1 & x & x^2 & \cdots & x^{n-1} \end{bmatrix}$$

$Ac = p(x)$ , a single vector  
summation that at once  
gives a linear combination  
of these monomials



清华大学统计学研究中心

56

## A Matrix Times a Matrix

57

$$B = AC, \quad b_j = Ac_j = \sum_{k=1}^m c_{kj} a_k$$

$$\left[ \begin{array}{c|c|c|c} b_1 & b_2 & \dots & b_n \end{array} \right] = \left[ \begin{array}{c|c|c|c} a_1 & a_2 & \dots & a_m \end{array} \right] \left[ \begin{array}{c|c|c|c} c_1 & c_2 & \dots & c_n \end{array} \right]$$

►  $b_j$  is a linear combination of the columns  $a_k$  with coefficients  $c_{kj}$

$$c_j = \begin{bmatrix} c_{1j} \\ c_{2j} \\ \vdots \\ c_{mj} \end{bmatrix}$$



清华大学统计学研究中心

57

## Examples

58

► Outer product: product of an  $m$ -dimensional column vector  $u$  with an  $n$ -dimensional row vector  $v$

$$\begin{bmatrix} u \end{bmatrix} \begin{bmatrix} v_1 & v_2 & \dots & v_n \end{bmatrix} = \begin{bmatrix} v_1 u & v_2 u & \dots & v_n u \end{bmatrix} = \begin{bmatrix} v_1 u_1 & \dots & v_n u_1 \\ \vdots & & \vdots \\ v_1 u_m & \dots & v_n u_m \end{bmatrix}$$

►  $B = AR$

rank 1

$$\left[ \begin{array}{c|c|c} b_1 & \dots & b_n \end{array} \right] = \left[ \begin{array}{c|c|c} a_1 & \dots & a_n \end{array} \right] \begin{bmatrix} 1 & \dots & 1 \\ & \ddots & \\ & & 1 \end{bmatrix} \quad b_j = Ar_j = \sum_{k=1}^j a_k$$

The matrix  $R$  is a discrete analogue of an indefinite integral operator




清华大学统计学研究中心

58

## A Matrix Inverse Times a Vector

59

- ▶  $x = A^{-1}b$
- ▶ Think of  $x$  as the unique vector that satisfies the equation  $Ax = b$
- ▶  $A^{-1}b$  is the vector of coefficients of the expansion of  $b$  in the basis of columns of  $A$
- ▶ Multiplication by  $A^{-1}$  is a change of basis operation:



清华大学统计学研究中心

59

## Multiplication by a Unitary Matrix

60

- ▶ A square matrix  $Q$  is unitary (in the real case, orthogonal), if  $Q^* = Q^{-1}$   

conjugate transpose of  $Q$
- ▶  $Qx$  is the linear combination of the columns of  $Q$  with coefficients  $x$ , and  $Q^*b$  is the vector of coefficients of the expansion of  $b$  in the basis of columns of  $Q$
- ▶ These processes of multiplication by a unitary matrix or its adjoint preserve geometric structure in the Euclidean sense, because inner products are preserved



清华大学统计学研究中心

60

## Invariance under Unitary Multiplication

61

- ▶ Angles between vectors are preserved, and so are their lengths:  $\|Qx\| = \|x\|$
- ▶ In the real case, multiplication by an orthogonal matrix  $Q$  corresponds to a rigid rotation (if  $\det Q = 1$ ) or reflection (if  $\det Q = -1$ ) of the vector space

▶ Theorem. For any  $A \in \mathbb{C}^{m \times n}$  and unitary  $Q \in \mathbb{C}^{m \times m}$ , we have

$$\|QA\|_2 = \|A\|_2 \triangleq \sup_{\substack{x \in \mathbb{C}^n \\ x \neq 0}} \frac{\|Ax\|_2}{\|x\|_2}, \quad \|QA\|_F = \|A\|_F \triangleq \left( \sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2 \right)^{1/2}$$



清华大学统计学研究中心

61

## Cholesky Factorization/Decomposition

62



清华大学统计学研究中心

62

## The Object

63

- ▶ Hermitian positive definite matrices can be decomposed into triangular factors twice as quickly as general matrices. The standard algorithm for this, Cholesky factorization, is a variant of Gaussian elimination that operates on both the left and the right of the matrix at once, preserving and exploiting symmetry
- ▶ In a statistical context, where data is usually real, a ‘Hermitian’ matrix becomes ‘symmetric’



清华大学统计学研究中心

63

## Symmetric matrix

64

- ▶ For real symmetric matrices we have two crucial properties:
  - All eigenvalues of a real symmetric matrix are real
  - Eigenvectors corresponding to distinct eigenvalues are orthogonal
- ▶ So it's possible to choose a complete set of eigenvectors that are perpendicular (or even orthonormal)



清华大学统计学研究中心

64



## Hermitian Positive Definite Matrix

65

If  $A$  is an  $m \times m$  Hermitian positive definite matrix and  $X$  is an  $m \times n$  matrix of full rank with  $m \geq n$ , then

- ▶  $X^*AX$  is also Hermitian positive definite
- ▶ any principal submatrix of  $A$  is positive definite
- ▶ every diagonal entry of  $A$  is a positive real number
- ▶ the eigenvalues are also positive real numbers



清华大学统计学研究中心

65

## Symmetric Gaussian Elimination

66

- ▶ When a single step of Gaussian elimination is applied

$$A = \begin{bmatrix} 1 & w^* \\ w & K \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ w & I \end{bmatrix} \begin{bmatrix} 1 & w^* \\ 0 & K - ww^* \end{bmatrix}$$

- ▶ In order to maintain symmetry

$$\begin{bmatrix} 1 & w^* \\ 0 & K - ww^* \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & K - ww^* \end{bmatrix} \begin{bmatrix} 1 & w^* \\ 0 & I \end{bmatrix}$$

- ▶ Combining the operations above

$$A = \begin{bmatrix} 1 & w^* \\ w & K \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ w & I \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & K - ww^* \end{bmatrix} \begin{bmatrix} 1 & w^* \\ 0 & I \end{bmatrix}$$

- ▶ The idea of Cholesky factorization is to continue this process, zeroing one column and one row of  $A$  symmetrically until it is reduced to the identity



清华大学统计学研究中心

66

## Cholesky Factorization

67

- ▶ In general, denote  $\alpha = \sqrt{a_{11}}$
- ▶  $A = \begin{bmatrix} a_{11} & w^* \\ w & K \end{bmatrix} = \begin{bmatrix} \alpha & 0 \\ w/\alpha & I \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & K - ww^*/a_{11} \end{bmatrix} \begin{bmatrix} \alpha & w^*/\alpha \\ 0 & I \end{bmatrix} = R_1^* A_1 R_1$   
 $A_1 = R_1^{-*} A R_1^{-}$
- ▶ The upper-left entry of the submatrix  $K - ww^*/a_{11}$  should be positive
- ▶ The process is continued down to the bottom-right corner, giving us eventually a factorization

$$A = \underbrace{R_1^* R_2^* \cdots R_m^*}_{R^*} \underbrace{R_m \cdots R_2 R_1}_{R}$$



清华大学统计学研究中心

67

- ▶ Theorem. Every Hermitian positive definite matrix  $A$  has a unique Cholesky factorization

68

$$A = R^* R, \quad r_{jj} > 0,$$

where  $R$  is upper-triangular

- ▶ Algorithm:

This algorithm is always stable

### Cholesky Factorization

$R = A$  only half of the matrix being  
 for  $k = 1$  to  $m$  operated on needs to be  
 for  $j = k + 1$  to  $m$  represented explicitly.  
 $R_{j,j:m} = R_{j,j:m} - R_{k,j:m} R_{kj} / R_{kk}$   
 $R_{k,k:m} = R_{k,k:m} / \sqrt{R_{kk}}$

- ▶ Operation count:

$$GE: 2/3 m^3$$

$$\sum_{k=1}^m \sum_{j=k+1}^m 2(m-j+1) + 1 \sim \sum_{k=1}^m \sum_{j=k+1}^m 2(m-j) \sim 2 \sum_{k=1}^m \sum_{j=k+1}^m j \sim \sum_{k=1}^m k^2 \sim \frac{1}{3} m^3 \text{ flops}$$



清华大学统计学研究中心

68

## Solution of $Ax = b$

69

- ▶ If  $A$  is symmetric positive definite, the standard way to solve a system of equations  $Ax = b$  is by Cholesky factorization
- ▶ Reduces the system to  $R^*Rx = b$ , then solve two triangular systems in succession:
  - first  $R^*y = b$  for the unknown  $y$
  - then  $Rx = y$  for the unknown  $x$
- ▶ Each triangular solution requires just  $\sim m^2$  flops, so the total work is again  $\sim 1/3m^3$  flops
- ▶ This process is backward stable



清华大学统计学研究中心

69

70

## Eigenvalue Decomposition



清华大学统计学研究中心

70

## Eigenvalues and Eigenvectors

71

- ▶ Let  $A \in \mathbb{C}^{m \times m}$  be a **square** matrix. A nonzero vector  $x \in \mathbb{C}^m$  is an eigenvector of  $A$ , and  $\lambda \in \mathbb{C}$  is its corresponding eigenvalue, if

$$Ax = \lambda x$$

- ▶ Idea: the action of a matrix  $A$  on a subspace  $S$  of  $\mathbb{C}^m$  may sometimes mimic scalar multiplication
- ▶ When this happens, the special subspace  $S = E_\lambda$  is called an eigenspace, and any nonzero  $x \in S$  is an eigenvector
- ▶ The set of all the eigenvalues of a matrix  $A$  is the spectrum of  $A$ , a subset of  $\mathbb{C}$  denoted by  $\Lambda(A)$
- ▶ Properties:

$$\det(A) = \prod_{j=1}^m \lambda_j, \operatorname{tr}(A) = \sum_{j=1}^m \lambda_j$$



清华大学统计学研究中心

71

## Geometric and Algebraic Multiplicity

72

- ▶ The dimension of eigenspace  $E_\lambda$  (also the nullspace of  $A - \lambda I$ ) is known as the geometric multiplicity of  $\lambda$
- ▶ The degree  $m$  polynomial  $P_A(z) = \det(zI - A)$  is called the characteristic polynomial

$$\begin{aligned} p_A(z) &= (z - \lambda_1)(z - \lambda_2) \cdots (z - \lambda_m) \\ &= (z - \lambda_1)^{k_1} (z - \lambda_2)^{k_2} \cdots (z - \lambda_r)^{k_r}, \quad r \leq m \end{aligned}$$

- ▶ The algebraic multiplicity of an eigenvalue  $\lambda_j$  of  $A$  is its multiplicity  $k_j$  as a root of  $P_A$



清华大学统计学研究中心

72

## Eigenvalue Problems

73

- ▶ Not like the problems involving square or rectangular linear systems of equations
- ▶ Make sense only when the range and the domain spaces are the same
- ▶ This reflects the fact that in applications, eigenvalues are generally used where a matrix is to be compounded iteratively, either explicitly as a power  $A^k$  or implicitly in a functional form such as  $e^{tA}$



清华大学统计学研究中心

73

74

- ▶ Broadly speaking, eigenvalue analysis are useful:
  - Algorithmically, simplify solutions of certain problems by reducing a coupled system to a collection of scalar problems
  - Physically, give insight into the behavior of evolving systems governed by linear equations
  - Examples: the study of resonance (e.g., of musical instruments when struck or plucked or bowed) and of stability (e.g., of fluid flows subjected to small perturbations), where eigenvalues tend to be particularly useful for analyzing behavior for large times  $t$



清华大学统计学研究中心

74

## Eigenvalue Decomposition

75

►  $A = X\Lambda X^{-1}$

here  $X$  is nonsingular and  $\Lambda$  is diagonal

► Such a factorization does not always exist!

$$\left[ \begin{array}{c} A \end{array} \right] \left[ \begin{array}{c|c|c|c} x_1 & x_2 & \cdots & x_m \end{array} \right] = \left[ \begin{array}{c|c|c|c} x_1 & x_2 & \cdots & x_m \end{array} \right] \left[ \begin{array}{ccc} \lambda_1 & & \\ & \lambda_2 & \\ & & \ddots \\ & & & \lambda_m \end{array} \right]$$

► The  $j$ th column of  $X$  is an eigenvector of  $A$  and the  $j$ th entry of  $\Lambda$  is the corresponding eigenvalue



清华大学统计学研究中心

75

## $Ax = b$

76

► The eigenvalue decomposition expresses a change of basis to "eigenvector coordinates"

►  $A = X\Lambda X^{-1}$ ,  $(X^{-1}b) = \Lambda(X^{-1}x)$

► Thus, to compute  $Ax$ , we can expand  $x$  in the basis of columns of  $X$ , apply  $\Lambda$ , and interpret the result as a vector of coefficients of a linear combination of the columns of  $X$



清华大学统计学研究中心

76

## Existence

77

- ▶ An eigenvalue whose algebraic multiplicity exceeds its geometric multiplicity is a defective eigenvalue
- ▶ A matrix that has one or more defective eigenvalues is a defective matrix
- ▶ Theorem. A square matrix has an eigenvalue decomposition  $X\Lambda X^{-1}$  if and only if it is nondefective/diagonalizable
- ▶ Note: If  $X$  is highly ill-conditioned, then a great deal of information may be discarded in passing from  $A$  to  $\Lambda$



清华大学统计学研究中心

77

## Unitary Diagonalization

78

- ▶ Sometimes not only does an  $m \times m$  matrix  $A$  have  $m$  linearly independent eigenvectors, but these can be chosen to be **orthogonal**
- ▶  $A$  is unitarily diagonalizable if there exists a unitary matrix  $Q$  such that

$$A = Q\Lambda Q^*$$

- ▶ A matrix  $A$  is normal if  $A^*A = AA^*$
- ▶ Theorem. A matrix is unitarily diagonalizable if and only if it is normal



清华大学统计学研究中心

78

## Schur Factorization

79

- ▶ A Schur factorization (unitary triangularization) of a matrix  $A$  is a factorization

$$A = QTQ^*$$

where  $Q$  is unitary and  $T$  is upper-triangular

- ▶ Note that since  $A$  and  $T$  are similar, the eigenvalues of  $A$  necessarily appear on the diagonal of  $T$
- ▶ Theorem. Every square matrix has a Schur factorization



清华大学统计学研究中心

79

## Synopsis: Eigenvalue-revealing Factorizations

80

Factorizations of a matrix that reduce it to a form in which the eigenvalues are explicitly displayed

- ▶ A diagonalization  $A = X\Lambda X^{-1}$  exists if and only if  $A$  is nondefective
- ▶ A unitary diagonalization  $A = Q\Lambda Q^*$  exists if and only if  $A$  is normal
- ▶ A unitary triangularization (Schur factorization)  $A = QTQ^*$  always exists



清华大学统计学研究中心

80



81

# Singular Value Decomposition (SVD)



清华大学统计学研究中心

81


## A Geometric Observation

82

- Many problems of linear algebra can be better understood if we first ask the question:

What if we take the SVD?

- Motivation: the image of the unit sphere under any  $m \times n$  matrix is a hyperellipse (the  $m$ -dimensional generalization of an ellipse)




清华大学统计学研究中心

SVD of a 2 x 2 matrix

82

83

- Hyperellipse: the surface obtained by stretching the unit sphere in  $\mathbb{R}^m$  by some factors  $\sigma_1, \dots, \sigma_m$  (possibly zero) in some orthogonal directions  $u_1, \dots, u_m \in \mathbb{R}^m$
- Take the  $u_i$  to be unit vectors, vectors  $\{\sigma_i u_i\}$  are the principal semiaxes of the hyperellipse, with lengths  $\sigma_i$
- Exactly  $\text{rank}(A)$  of the lengths  $\sigma_i$  will be nonzero



清华大学统计学研究中心

83

## Singular Values and Vectors

84

- The lengths of the  $n$  principal semiaxes of  $AS$ ,  $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n$ , are called singular values of  $A$
- The unit vectors  $\{u_1, \dots, u_n\}$  oriented in the directions of the principal semiaxes of  $AS$ , numbered to correspond with the singular values, are called left singular vectors of  $A$
- The right singular vectors of  $A$  are the unit vectors  $\{v_1, \dots, v_n\}$  in  $S$  that are the preimages of the principal semiaxes of  $AS$ , numbered so that

$$Av_j = \sigma_j u_j, 1 \leq j \leq n$$



清华大学统计学研究中心

84

## Reduced SVD ( $m \geq n$ )

85

►  $Av_j = \sigma_j u_j, 1 \leq j \leq n$

$$\begin{bmatrix} A \end{bmatrix} \begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix} = \begin{bmatrix} u_1 & u_2 & \cdots & u_n \end{bmatrix} \begin{bmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_n \end{bmatrix}$$

full rank       $n \times n$  matrix with orthonormal columns-unitary       $m \times n$  matrix with orthonormal columns       $n \times n$  diagonal matrix with positive real entries

$$A = \hat{U} \hat{\Sigma} V^*$$



清华大学统计学研究中心

85

## Full SVD

86

► Since  $\hat{U} \in \mathbb{C}^m$ , extend it to be a unitary matrix

$$A = U \Sigma V^*$$

$$\begin{bmatrix} A \end{bmatrix} = \begin{bmatrix} U \end{bmatrix} \begin{bmatrix} \Sigma \end{bmatrix} \begin{bmatrix} V^* \end{bmatrix}$$

$m \geq n$ , full rank       $m \times m$  unitary      "silent" columns/rows       $m \times n$  diagonal with positive real entries       $n \times n$  unitary

► If  $\text{rank}(A) = r < n$ , extend both  $U$  and  $V$  to be unitary matrices




清华大学统计学研究中心

86

## Visualization

87



清华大学统计学研究中心

87

## Existence and Uniqueness

88

### Theorem

- Every matrix  $A \in \mathbb{C}^{m \times n}$  has a singular value decomposition  $A = U \Sigma V^*$
- Furthermore, the singular values  $\{\sigma_j\}$  are uniquely determined; and, if  $A$  is square and the  $\sigma_j$  are distinct, the left and right singular vectors  $\{u_j\}$  and  $\{v_j\}$  are uniquely determined up to complex signs



清华大学统计学研究中心

88

## A Change of Bases

89

$$\blacktriangleright b = Ax \Leftrightarrow U^*b = U^*U\Sigma V^*x = \Sigma V^*x$$

Denote  $b' = U^*b$ ,  $x' = V^*x$ , then

$$\Leftrightarrow b' = \Sigma x'$$

- ▶ Thus  $A$  reduces to the diagonal matrix  $\Sigma$  when the range(domain) is expressed in the basis of columns of  $U(V)$
- ▶ The SVD makes it possible for us to say that **every matrix is diagonal**-if only one uses the proper bases for the domain and range spaces



清华大学统计学研究中心

89

## SVD vs. Eigenvalue Decomposition

90

- ▶ For unitarily diagonalizable  $A = Q\Lambda Q^*$ , this is an ED, and  $A = Q|\Lambda|\text{sign}(\Lambda)Q^*$  gives a SVD
- ▶ They both have the theme of diagonalizing a matrix, while the ED  $A = X\Lambda X^{-1}$  expresses it in terms of one new basis
- ▶  $b = Ax = X\Lambda X^{-1}x \Leftrightarrow X^{-1}b = \Lambda X^{-1}x$

Denote  $b' = X^{-1}b$ ,  $x' = X^{-1}x$ , then

$$\Leftrightarrow b' = \Lambda x'$$

Here both the range and domain are expressed in the basis of eigenvectors of  $A$  (columns of  $X$ )



清华大学统计学研究中心

90

91

► Fundamental differences:

- The SVD uses two different bases (the sets of left and right singular vectors), whereas the ED uses just one (the eigenvectors)
- The SVD uses orthonormal bases, whereas the ED uses a basis that generally is not orthogonal
- Not all matrices (even square ones) have an ED, but all matrices (even rectangular ones) have a SVD
- In applications, eigenvalues tend to be relevant to problems involving the behavior of  $A^k$  or  $e^{tA}$ , whereas singular vectors tend to be relevant to problems involving the behavior of  $A$  or  $A^{-1}$



清华大学统计学研究中心

91

92

## Matrix Properties via the SVD

- 1. The rank of  $A_{m \times n}$  is  $r$ , the number of nonzero singular values
- 2.  $\text{range}(A) = \langle u_1, \dots, u_r \rangle$ ,  $\text{null}(A) = \langle v_{r+1}, \dots, v_n \rangle$
- 3.  $\|A\|_2 = \sup_{\substack{x \in \mathbb{C}^n \\ x \neq 0}} \frac{\|Ax\|_2}{\|x\|_2} = \sigma_1$ ,  $\|A\|_F = \left( \sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2 \right)^{1/2} = \sqrt{\sigma_1^2 + \dots + \sigma_r^2}$
- 4. The nonzero singular values of  $A$  are the square roots of the nonzero eigenvalues of  $A^*A$  or  $AA^*$
- 5. If  $A = A^*$ , then the singular values of  $A$  are the absolute values of the eigenvalues of  $A$
- 6. For  $A \in \mathbb{C}^{m \times m}$ ,  $|\det(A)| = \prod_{j=1}^m \sigma_j$



清华大学统计学研究中心

92

## Computational Consequences

93

- ▶ The best method for determining the rank of a matrix is to count the number of singular values greater than a judiciously chosen tolerance (Property 1)
- ▶ The most accurate method for finding an orthonormal basis of a range or a nullspace is via property 2
- ▶ QR factorization provides alternative algorithms that are faster but not always as accurate
- ▶ Property 3 represents the standard method for computing  $\|A\|_2$
- ▶ Besides these examples, the SVD is also an ingredient in robust algorithms for least squares fitting, intersection of subspaces, regularization, and numerous other problems



清华大学统计学研究中心

93

$$A = U\Sigma V^*$$

94

- ▶  $U$ : The left-singular vectors of  $A$  are a set of orthonormal eigenvectors of  $AA^*$
- ▶  $V$ : The right-singular vectors of  $A$  are a set of orthonormal eigenvectors of  $A^*A$
- ▶  $\Sigma$ : The non-zero singular values of  $A$  (found on the diagonal entries of  $\Sigma$ ) are the square roots of the non-zero eigenvalues of both  $A^*A$  and  $AA^*$
- ▶  $A$  is the sum of  $r$  rank-one matrices:

$$A = \sum_{j=1}^r \sigma_j u_j v_j^*$$

Low-rank approximations



清华大学统计学研究中心

94

95

# Least Squares Problems



清华大学统计学研究中心

95

96

## The Problem

- ▶ Least squares data-fitting has been an indispensable tool since its invention by Gauss and Legendre around 1800, with ramifications extending throughout the mathematical sciences
- ▶ The problem is the solution of an overdetermined system of equations  $Ax = b$  - rectangular, with more rows than columns
- ▶ The least squares idea is to "solve" such a system by minimizing the 2-norm of the residual  $b - Ax$



清华大学统计学研究中心

96



## Orthogonal Projection and the Normal Equations

97

- Theorem. Let  $A \in \mathbb{C}^{m \times n}$  ( $m \geq n$ ) and  $b \in \mathbb{C}^m$  be given. A vector minimizes the residual norm  $\|r\|_2 = \|b - Ax\|_2$ , thereby solving the least squares problem, if and only if  $x \in \mathbb{C}^n$ , that is,  $A^*r = 0$ , or equivalently,

$$A^*Ax = A^*b,$$

$$r \perp \text{range}(A)$$

or again equivalently,

$$Pb = Ax,$$

where  $P \in \mathbb{C}^{m \times m}$

is the orthogonal projector onto  $\text{range}(A)$

The  $n \times n$  system of equations, known as the normal equations, is nonsingular if and only if  $A$  has full rank. Consequently the solution  $x$  is unique if and only if  $A$  has full rank



清华大学统计学研究中心

97

## Pseudoinverse

98

- If  $A$  has full rank, then the solution  $x$  to the least squares problem is given by  $x = (A^*A)^{-1}A^*b$
- The pseudoinverse of  $A$  is  $A^+ = (A^*A)^{-1}A^* \in \mathbb{C}^{n \times m}$
- Then the full-rank linear least squares problem is to compute one or both of the vectors

$$x = A^+b, \quad y = Pb$$

where  $A^+$  is the pseudoinverse of  $A$  and  $P$  is the orthogonal projector onto  $\text{range}(A)$

- There are three leading algorithms for doing this



清华大学统计学研究中心

98

## Normal Equations

99

- ▶ The classical way is to solve the normal equations
- ▶ If  $A$  has full rank, this is a square, Hermitian positive definite system of equations of dimension  $n$
- ▶ The standard method of solving such a system is by Cholesky factorization,  $A^*A = R^*R$ , where  $R$  is upper-triangular, reducing the normal equations to

$$R^*Rx = A^*b$$



清华大学统计学研究中心

99

## Algorithm 1

100

### Least Squares via Normal Equations

1. Form the matrix  $A^*A$  and the vector  $A^*b$
2. Compute the Cholesky factorization  $A^*A = R^*R$
3. Solve the lower-triangular system  $R^*w = A^*b$  for  $w$
4. Solve the upper-triangular system  $Rx = w$  for  $x$

Operation count:

- ▶ To compute  $A^*A$  requires  $\sim \sum_{i=1}^n \sum_{j=1}^i 2m \sim mn^2$  flops
- ▶ Cholesky factorization takes  $\sim n^3/3$  flops
- ▶ Since these two dominate the computation, the total work requires  $\sim mn^2 + n^3/3$  flops



清华大学统计学研究中心

100

## QR Factorization

101

- ▶ The "modern classical" method, popular since the 1960s, is based upon reduced QR factorization
- ▶ Use Gram-Schmidt orthogonalization or, more usually, Householder triangularization, to get  $A = \hat{Q}\hat{R}$
- ▶ The orthogonal projector  $P = \hat{Q}\hat{Q}^*$ , so  $y = Pb = \hat{Q}\hat{Q}^*b$
- ▶  $Ax = y \leftrightarrow \hat{Q}\hat{R}x = \hat{Q}\hat{Q}^*b \leftrightarrow \hat{R}x = \hat{Q}^*b$   
 $A^*Ax = A^*b \leftrightarrow \hat{R}^*\hat{Q}^*\hat{Q}\hat{R}x = \hat{R}^*\hat{Q}^*b \leftrightarrow \hat{R}x = \hat{Q}^*b$



清华大学统计学研究中心

101

## Algorithm 2

102

### Least Squares via QR Factorization

1. Compute the reduced QR factorization  $A = \hat{Q}\hat{R}$
2. Compute the vector  $\hat{Q}^*b$
3. Solve the upper-triangular system  $\hat{R}x = \hat{Q}^*b$  for  $x$

Operation count:

- ▶ The work is dominated by the cost of the QR factorization
- ▶ If Householder reflections are used for this step, the total work requires  $\sim 2mn^2 - 2/3n^3$  flops



清华大学统计学研究中心

102

## SVD and Algorithm 3

103

- Now  $A = \hat{U}\hat{\Sigma}V^*$ , take  $P = \hat{U}\hat{U}^*$ , or  
 $A^*Ax = A^*b \leftrightarrow V\hat{\Sigma}\hat{U}^*\hat{U}\hat{\Sigma}V^*x = V\hat{\Sigma}\hat{U}^*b \leftrightarrow \hat{\Sigma}V^*x = \hat{U}^*b$

### Least Squares via SVD $A = \hat{U}\hat{\Sigma}V^*$

1. Compute the reduced SVD
2. Compute the vector  $\hat{U}^*b$
3. Solve the **diagonal** system  $\hat{\Sigma}w = \hat{U}^*b$  for  $w$
4. Set  $x = Vw$

Operation count:

- The work is dominated by the cost of the SVD, which requires  $\sim 2mn^2 + 11n^3$  flops



清华大学统计学研究中心

103

## Comparison of Algorithms

104

- Each of these methods is advantageous in certain situations
- When speed is the only consideration, Algorithm 1 may be the best
- However, solving the normal equations is not always stable in the presence of rounding errors, thus numerical analysts recommended Algorithm 2 instead
- If  $A$  is close to rank-deficient, Algorithm 2 itself has less-than-ideal stability properties, then Algorithm 3 is preferred



清华大学统计学研究中心

104

## Homework

105

- ▶ Verify that the flop count is  $\sim 3mn^2 - n^3$  for Givens QR on p. 38
- ▶ Write R functions for each algorithm, and design examples to compare their results. (CGS, MGS and Householder QR factorization, Gaussian elimination with and without partial pivoting)
- ▶ Solve sets of linear equations with your functions
- ▶ Let  $A$  be a  $10 \times 10$  random matrix with entries from the standard normal distribution, minus twice the identity. Write a program to plot  $\|e^{tA}\|_2$  against  $t$  for  $0 \leq t \leq 20$  on a log scale, comparing the result to the straight line  $e^{t\alpha(A)}$ , where  $\alpha(A) = \max_j \operatorname{Re}(\lambda_j)$  is the spectral abscissa of  $A$ . Run the program for ten random matrices  $A$  and comment on the results. What property of a matrix leads to a  $\|e^{tA}\|_2$  curve that remains oscillatory as  $t \rightarrow \infty$ ?



清华大学统计学研究中心