





# Análise Avançada de Dados Quantitativos 2019-1

Prof. Dr. Wagner de Lara Machado

# Founders of statistics




Quem sou eu?



# Formação acadêmica e atuação profissional

Nível	Tema	Ano de conclusão
Graduação em Psicologia (ULBRA)	Prevalência de ansiedade e depressão em docentes	2007
Mestrado acadêmico (UFRGS)	Adaptação da Escala de Bem-estar Psicológico	2010
Doutorado (UFRGS)	Saúde mental positiva Adaptação de medidas de saúde mental	2013
Estágio de Pós-doutorado (UFRGS)	Saúde mental positiva Centro de análise de dados (CAD)	2015
Professor da graduação e pós-graduação <i>stricto sensu</i> Psicologia	Grupo de pesquisa Avaliação Psicológica do Potencial Humano	2015 a 2018



## Grupo de Pesquisa Avaliação em Bem-Estar e Saúde Mental



-



11



42




Membro da  
Diretoria da  
Associação  
Brasileira de  
Psicologia  
Positiva



# Links

- <https://scholar.google.com.br/citations?user=fH6qCDoAAAAJ&hl=en>
- <https://github.com/wagnerLM>






# Análise Avançada de Dados Quantitativos

- <https://github.com/wagnerLM/quantia>








Livro texto  
base da  
disciplina

- Ver link no github
- <https://raw.githubusercontent.com/wagnerLM/quantia/master/linklivro>

# Livros auxiliares sobre R e estatística

- Acesse o link:

<https://www.dropbox.com/sh/4kc8cjy51nkhuvr/AACLK0vJonv66OJSG3ZHId6Ra?dl=0>



## QUALITATIVAS

## QUANTITATIVAS

NOMINAIS OU  
CATEGÓRICAS

ORDINAIS

DISCRETAS

CONTÍNUAS

Sexo  
Cor do olho  
Profissão

NSE  
Nível educacional  
Escala Likert

NÚMEROS NATURAIS (nº DE  
OBJETOS)

Altura  
Idade  
Tempo de reação

# Tipos de variáveis

Nominal	Ordinal	Intervalar	Razão
<p>Podemos atribuir números como rótulo, porém sem manter as propriedades dos números.</p> <p>Ex. Sexo</p> <p>1 – Masculino</p> <p>2 – Feminino</p>	<p>É possível estabelecer uma ordem de grandeza ou magnitude, porém a distância não é conhecida.</p> <p>Ex. Dureza dos materiais</p> <p>Nível educacional</p> <p>1 – Fundamental</p> <p>2 – Médio</p> <p>3 – Superior</p> <p>4 – Pós-graduação</p>	<p>A distância entre as unidades de medida é conhecida, podem ser realizadas a maioria das operações matemáticas.</p> <p>Ex. Escores modelados de instrumentos psicométricos</p>	<p>Escala intervalar com um zero conhecido e não arbitrário.</p> <p>Ex. Quando zero significa ausência do atributo</p>

## Níveis de mensuração

# Questões sobre método

- Tipos (papéis) de variáveis
  - Confundidora
  - Controle
  - Dependente
  - Independente
  - Binária ou dicotômica
  - Covariável
  - Critério
  - “Dummy”
  - Endógena
  - Exógena
  - Latente
  - Observada
  - Mediadora
  - Moderadora
  - Preditora
  - Desfecho
  - Instrumental

Questões  
sobre  
método  
quantitativo

FREQUENTISTA	BAYESIANA
$P(D   H)$	$P(H   D)$

- Inferência estatística, teste de hipótese

Sendo D = dados e H = hipótese

- Interpretação do “p” ou Bayes Factor 0,4




# Software R

---

- R Project
- John Chambers
- Software livre, colaborativo
- Possui um repositório CRAN
- Bibliotecas e pacotes (códigos e algoritmos)
- 100% FREE!!!








# O quê é a linguagem r?

- R é um “dialeto” da linguagem S
- 1993 R é lançado ao público
- 1995 General Public License
- 1997 o R Core Group é formado –  
Controla a fonte de códigos do R
- 2000 R version 1.0.0 é lançado
- 2015 version 3.2.1




# Porquê usar a linguagem r?

- Capacidades gráficas muito sofisticadas e melhores que muitos softwares
- Linguagem de programação que possibilita o desenvolvimento de novas ferramentas
- Comunidade de usuários muito ativa e participativa



# Porquê usar a linguagem r?

# Porquê usar a linguagem r?





## Passos iniciais

---

- É o sistema básico – feito de forma colaborativa – Possui funções básicas que podem ser combinadas em pacotes mais avançados
- R Studio – Uma interface mais amigável para o uso do R. Permite análise, escrita (tem gente que escreve a tese por aqui), e publicações.





# R desing

- O sistema R é dividido em duas partes:
  - A “base” do sistema R que pode-se fazer o download no CRAN
  - Todo o resto – Funcionalidades chamadas de packages
  - Atualmente existem mais de 6000 packages



```
gibbs.cpp x
Source on Save
Source

1 #include <Rcpp.h>
2 using namespace Rcpp;
3
4 // [[Rcpp::export]]
5 NumericMatrix gibbs(int N, int thin) {
6
7     NumericMatrix mat(N, 2);
8     double x = 0, y = 0;
9
10    for(int i = 0; i < N; i++) {
11        for(int j = 0; j < thin; j++) {
12            x = R::rgamma(3.0, 1.0 / (y * y + 4));
13            y = R::rnorm(1.0 / (x + 1), 1.0 / sqrt(2 * x + 2));
14        }
15        mat(i, 0) = x;
16        mat(i, 1) = y;
17    }
18
19    return(mat);
20 }

19:15 gibbs(int N, int thin): NumericMatrix C/C++
```

<https://www.r-project.org/>



[\[Home\]](#)

**Download**

[CRAN](#)

**R Project**

[About R](#)

[Logo](#)

[Contributors](#)

[What's New?](#)

[Reporting Bugs](#)

[Development Site](#)

[Conferences](#)

[Search](#)

**R Foundation**

[Foundation](#)

[Board](#)

[Members](#)

[Donors](#)

[Donate](#)

# The R Project for Statistical Computing

## Getting Started

R is a free software environment for statistical computing and graphics. It compiles and runs on a wide variety of UNIX platforms, Windows and MacOS. To **download R**, please choose your preferred [CRAN mirror](#).

If you have questions about R like how to download and install the software, or what the license terms are, please read our [answers to frequently asked questions](#) before you send an email.

## News

- **R version 3.5.1 (Feather Spray)** has been released on 2018-07-02.
- The R Foundation has been awarded the Personality/Organization of the year 2018 award by the professional association of German market and social researchers.
- **R version 3.5.0 (Joy in Playing)** has been released on 2018-04-23.

## News via Twitter




The R Foundation Retweeted



**Peter Dalgaard**

[@pdalgd](#)

[#rstats](#) 3.5.1 "Feather Spray" is released (source version)



# **A (very) short introduction to R**

Paul Torfs & Claudia Brauer

Hydrology and Quantitative Water Management Group

Wageningen University, The Netherlands

16 April 2012

<https://www.dropbox.com/s/vppgdkx1u909r3n/Torfs%20Brauer%202012%20-%20introR.pdf?dl=0>

[Home](#)[Learn](#)[Teach](#)[Contribute](#)[Blog](#)[FAQ](#)[Help](#)

Learn R, in R.

swirl teaches you R programming and data science  
interactively, at your own pace, and right in the R  
console!

<https://swirlstats.com/>



R version 3.1.3 (2017-11-30) Rite Editing RICE  
Copyright (C) 2017 The R Foundation for Statistical Computing  
Platform: x86\_64-w64-mingw32/x64 (64-bit)

R é um software livre e vem sem GARANTIA ALGUMA.  
Você pode redistribuí-lo sob certas circunstâncias.  
Digite 'license()' ou 'licence()' para detalhes de distribuição.

R é um projeto colaborativo com muitos contribuidores.  
Digite 'contributors()' para obter mais informações e  
'citation()' para saber como citar o R ou pacotes do R em publicações.

Digite 'demo()' para demonstrações, 'help()' para o sistema on-line de ajuda,  
ou 'help.start()' para abrir o sistema de ajuda em HTML no seu navegador.  
Digite 'q()' para sair do R.

> |

# RStudio

Open source and enterprise-ready  
professional software for R

[Download](#)[Download](#)[Discover RStudio](#)[Discover](#)




Figure 1 The editor, workspace, console and plots windows in RStudio.


# RStudio

1 – script ou comandos; visualização de bancos de dados ou texto

2 – repositório de objetos e funções

3 – log de atividade

4 – gráficos e documentação



## Using R and psych for personality and psychological research

### User manual and help files

1. The [psych user manual](#) (pdf)
2. The [individual help files for the psych package](#) in html

### Vignettes

1. [introduction](#) (pdf) to the psych package

[Overview of psychometric functions](#) (pdf) by David C. Howell

[R](#) is a very powerful open source system for statistical computation and graphics. It consists of a language plus a run-time environment with graphics, a debugger, access to certain system functions, and the ability to run programs stored in script files. Base R is a foundation upon which more than 11,000 "packages" have been built. It is the use of these packages that makes R such a powerful tool for research.

The psych package has been developed at the [Personality, Motivation and Cognition](#) laboratory, University of Colorado Boulder.

## Conteúdos

- Análise fatorial exploratória
- <http://personality-project.org/r/psych/>

# lavaan

latent variable analysis

[About lavaan](#) [Tutorial](#) [Resources](#) [Version History](#)

---

## About lavaan

[Welcome](#)

[Getting started](#)

[Features](#)

[Development](#)

[Support](#)


[About](#)

## News:

- (22 Sept 2018): lavaan version 0.6-3 has been released on [CRAN](#). See [Version History](#) for more information.
- (10 Jun 2018): the blavaan paper (on Bayesian SEM with a lavaan syntax) is [published](#) in the Journal of Statistical Software.
- (18 Dec 2017): a tutorial on 'The Pairwise Likelihood Method for Structural Equation Modelling with ordinal variables and data with missing values using the R package lavaan' prepared by Myrsini Katsikatsou has been added to the (new) [tutorial](#) page of the resources section.
- (16 July 2017): a recording of my keynote presentation 'Structural Equation Modeling: models, software and stories' given at the [useR!2017 Conference](#) is available [here](#).

- Análise fatorial confirmatória
- Análise de trajetórias e modelagem estrutural de equações
- <http://lavaan.ugent.be/>





Psychological Bulletin  
1991, Vol. 110, No. 2, 305-314

Copyright 1991 by the American Psychological Assoc  
0033-29

## Conventional Wisdom on Measurement: A Structural Equation Perspective

Kenneth Bollen  
Sociology Department  
University of North Carolina at Chapel Hill

Richard Lennox  
Institute for Research in Social Science  
University of North Carolina at Chapel Hill

# Análise fatorial confirmatória



---

# *Journal of Statistical Software*

May 2012, Volume 48, Issue 6.

<http://www.jstatsoft.org/>

---

## **mirt: A Multidimensional Item Response Theory Package for the R Environment**


R. Philip Chalmers  
York University

- Teoria de Resposta ao Item
- <https://www.rdocumentation.org/packages/mirt/versions/1.30>
- <https://www.jstatsoft.org/article/view/v048i06>

# A History of Social Science Measurement

Benjamin D. Wright

*University of Chicago, MESA Psychometric Laboratory*



# Análise de Rede

- Capítulo de livro
- **MACHADO, W. L.**; VISSOCI, J. R. N. ; EPSKAMP, S. (2015). Análise de rede aplicada à psicometria e à avaliação psicológica. In C. S. Hutz; D. R. Bandeira & C. M. Trentini (Eds.). *Psicometria* (p. 125-146). Porto Alegre: Artmed.

(vii) `qgraph(E3,directed=FALSE)`  
`qgraph(E3,layout="circle")`  
`qgraph(E3)`  
`qgraph(E3,directed=FALSE,layout="spring")`




Figura 5. Redes ponderadas, direcionais e não-direcionais, sem (acima) e com (abaixo) o emprego do algoritmo de posicionamento.




Figura 13. Modelo de traço latente (esquerda) e de rede (direita) da comorbidade.

- Sacha Epskamp
- Assistant Professor in Psychological Methods and Psychometrics at the University of Amsterdam
- João Vissoci
- Pesquisador na divisão de Emergency Medicine do departamento de Cirurgia, e na divisão Duke Global Neurosurgery and Neuroscience (DGNN) do departamento de Neurocirurgia, na Duke University



### **Análise de rede aplicada à psicometria e a avaliação psicológica**

Wagner de Lara Machado

Pontifícia Universidade Católica de Campinas

João Ricardo Nickenig Vissoci

Faculdade Ingá e Duke University

Sacha Epskamp

Universiteit van Amsterdam

Positive Mental Health Scale: Validation of the *Mental Health Continuum – Short Form*

Wagner de Lara Machado – Pontifícia Universidade Católica de Campinas, Campinas, São Paulo, Brasil  
Denise Ruschel Bandeira – Universidade Federal do Rio Grande do Sul, Porto Alegre, Brasil




Figure 1. Map of items and persons. The vertical axis indicates the scale in *logits*. The distribution was obtained by fixing the contrary measure so the mean would be equal to zero.

Table 2  
Item Factor Loadings and Reliability Measures of the MHC-SF in the Bifactor Model

Item (summarized content)	Factor loading			
	General factor	EWB	SWB	PWB
1 – Happy	.70	.65		
2 – Interested	.78	.32		
3 – Satisfied	.78	.36		
4 – Contribute to society	.69		.07	
5 – Belong to community	.67		.23	
6 – Society is becoming a better place	.62		.63	
7 – People are good	.62		.41	
8 – Way society works makes sense	.57		.56	
9 – Likes own personality	.79			.21
10 – Manages responsibility well	.70			.19
11 – Relationships with others	.72			.24
12 – Grow and become a better person	.62			.39
13 – Confident to express own ideas	.68			.52
14 – Life has direction or meaning	.83			.12
Average extracted variance	.49	.22	.19	.10
Composite reliability	.93	.43	.47	.34

Note. EWB = emotional well-being, SWB = social well-being, PWB = psychological well-being.




Figure 2. Network of positive mental health indicators. The covariance structure indicates that the MHC-SF items are strongly associated with the others, resulting in a dense component. The items in the emotional (EWB), social (SWB) and psychological (PWB) well-being subscales exhibit moderate to strong crossed associations. Purpose in life and self-acceptance are the central nodes of the system, meaning that they are more strongly associated with the remainder of the items. The stronger line represents the correlation between “happy” and “satisfied” ( $r_{1,3} = .78$ ); the fainter line represents the correlation between “relations” and “coherence” ( $r_{6,11} = .35$ ).





# The Experience of Sexual Stigma and the Increased Risk of Attempted Suicide in Young Brazilian People from Low Socioeconomic Group

Angelo Brandelli Costa<sup>1\*</sup>, Andrew Pasley<sup>2</sup>, Wagner de Lara Machado<sup>3</sup>, Ernesto Alvarado<sup>4</sup>, Luciana Dutra-Thomé<sup>4</sup> and Silvia Helena Koller<sup>4</sup>




FIGURE 1 | Overall network with correlation (left) and regularized regression (gLASSO) functions (right). Variables names match those in Table 1.


This study was intended to analyze the intersection of experience of sexual stigma low-socioeconomic status, and suicide attempt amongst young Brazilians (11–24 years old). In each of the data collection periods (2004–2006:  $n = 7185$ ; 2010–2012:  $n = 2734$ ), participants completed a questionnaire-based instrument. Network analysis provided support for a Minority Stress Model, oriented around whether participants had experienced sexual stigma. Although suicide attempts decreased by 20% for participants

# Psychometric Perspectives on Diagnostic Systems



Denny Borsboom (2008)

*University of Amsterdam*



*Figure 1.* The left panel shows the relation between panic disorder and its symptoms from a latent variable modeling point of view. The right panel shows a representation of these symptoms as a causal system.

# Comorbidity: A network perspective

**Angélique O. J. Cramer**  
Department of Psychology, University of Amsterdam, 1018 WB Amsterdam,  
The Netherlands  
A.O.J.Cramer@uva.nl  
www.aojcramer.com

**Lourens J. Waldorp**  
Department of Psychology, University of Amsterdam, 1018 WB Amsterdam,  
The Netherlands  
L.J.Waldorp@uva.nl  
http://users.fmg.uva.nl/lwaldorp

**Han L. J. van der Maas**  
Department of Psychology, University of Amsterdam, 1018 WB Amsterdam,  
The Netherlands  
H.L.J.vanderMaas@uva.nl  
http://users.fmg.uva.nl/hvandermaas/

**Denny Borsboom**  
Department of Psychology, University of Amsterdam, 1018 WB Amsterdam,  
The Netherlands  
D.Borsboom@uva.nl  
http://sites.google.com/site/borsboomdenny/dennyborsboom




Figure 1. A model of comorbidity between disorders A and B, under the standard assumptions of latent variable modeling. The circles represent the disorders (i.e., latent variables) and the rectangles represent the observable core symptoms of those disorders (i.e.,  $X_1 - X_5$  for disorder A, and  $Y_1 - Y_5$  for disorder B). In this model, comorbidity is viewed as a correlation between the latent variables, visualized by the thick bidirectional edge between disorders A and B.





Figure 2. Comorbidity under a network approach. Disorder A consists of bidirectionally related symptoms  $X_1 - X_5$ , and disorder B consists of symptoms  $Y_1 - Y_5$ . Symptoms  $B_1$  and  $B_2$  are *bridge symptoms* that overlap between disorders A and B. In this model, comorbidity arises as a result of direct relations between the bridge symptoms of two disorders.

# Network Analysis: An Integrative Approach to the Structure of Psychopathology

Denny Borsboom and Angélique O.J. Cramer

Department of Psychology, University of Amsterdam, Amsterdam 1018 XA, The Netherlands;  
email: D.Borsboom@uva.nl



- Redes Bayesianas
- <http://www.bnlearn.com/>

