

### Elliptic curves: homework 8

Mastermath / DIAMANT, Spring 2019

Martin Bright and Marco Streng

1. Show that being isogenous is an equivalence relation on the set of complex tori, and that there are uncountably many isogeny classes of complex tori.
2. The *multiplier ring* of a lattice  $\Lambda$  is defined as

$$\mathcal{O}(\Lambda) = \{\alpha \in \mathbb{C} \mid \alpha\Lambda \subset \Lambda\}.$$

- (a) Show that  $\mathcal{O}(\Lambda)$  is a subring of  $\mathbb{C}$  isomorphic to the endomorphism ring  $\text{End}(\mathbb{C}/\Lambda)$  of the torus  $\mathbb{C}/\Lambda$ , that is, the ring of isogenies from  $\mathbb{C}/\Lambda$  to itself.
- (b) Show that we have  $\mathcal{O}(\Lambda) = \mathbb{Z}$  unless  $\Lambda$  is homothetic to a lattice of the form  $\mathbb{Z} + \mathbb{Z}\lambda$ , with  $\lambda \in \mathbb{C} \setminus \mathbb{R}$  a zero of an irreducible quadratic polynomial  $aX^2 + bX + c \in \mathbb{Z}[X]$ , and that in this exceptional case we have  $\mathcal{O}(\Lambda) = \mathbb{Z}[\frac{D+\sqrt{D}}{2}]$  with  $D = b^2 - 4ac < 0$ .

[In the exceptional case, we say that  $\mathbb{C}/\Lambda$  has *complex multiplication* by  $\mathcal{O}(\Lambda)$ .]

3. (a) For  $\Lambda = \mathbb{Z} + i\mathbb{Z}$ , show

$$\wp_{\Lambda}(iz) = -\wp_{\Lambda}(z), \quad \wp'_{\Lambda}(iz) = i\wp'_{\Lambda}(z), \quad G_k(\Lambda) = 0$$

for  $k \geq 3$  not divisible by 4.

- (b) Give the analogous results for  $\Lambda = \mathbb{Z} + \rho\mathbb{Z}$  where  $\rho = \exp(2\pi i/3) = \frac{i\sqrt{3}-1}{2}$ .
4. Show that the subrings of  $\mathbb{C}$  that are lattices correspond bijectively to the set of negative integers  $D$  satisfying  $D \equiv 0, 1 \pmod{4}$  under the association  $D \mapsto \mathcal{O}(D) = \mathbb{Z}[\frac{D+\sqrt{D}}{2}]$ . Show that there exists a ring homomorphism  $\mathcal{O}(D_1) \rightarrow \mathcal{O}(D_2)$  if and only if  $D_1/D_2$  is a square in  $\mathbb{Z}$ .
  5. Compute the structure of the group  $\text{Hom}(\mathbb{C}/\Lambda_1, \mathbb{C}/\Lambda_2)$  for each of the following choices of  $\Lambda_1$  and  $\Lambda_2$ :
    - (a)  $\Lambda_1 = \Lambda_2 = \mathbb{Z} + \mathbb{Z}i$ ;
    - (b)  $\Lambda_1 = \mathbb{Z} + \mathbb{Z}i$  and  $\Lambda_2 = \mathbb{Z} + \mathbb{Z}2i$ ;
    - (c)  $\Lambda_1 = \mathbb{Z} + \mathbb{Z}i$  and  $\Lambda_2 = \mathbb{Z} + \mathbb{Z}\sqrt{-2}$ .
  6. See also Exercise 6.6 of [Silverman] Define the  $j$ -invariant of a lattice  $\Lambda$  by

$$j(\Lambda) = 1728 \frac{g_2(\Lambda)^3}{g_2(\Lambda)^3 - 27g_3(\Lambda)^2}.$$

Prove that, if  $\Lambda$  and  $\Lambda'$  are homothetic lattices, then  $j(\Lambda) = j(\Lambda')$ .

7. For  $\tau \in \mathbb{C}$  with  $\Im(\tau) > 0$ , define  $j(\tau) := j(\mathbb{Z} + \mathbb{Z}\tau)$ .
- (a) Prove that  $j$  is a holomorphic function of  $\tau$ .
  - (b) Show  $j(\frac{a\tau+b}{c\tau+d}) = j(\tau)$  for all  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ .
  - (c) Compute  $j(i)$  and  $j(\rho)$  for  $\rho$  as in Problem 3.