

# Algebraic Topology II - Assignment 7

Matteo Durante, s2303760, Leiden University

15th May 2019

## Exercise 2

*Proof.* (a) It is sufficient to notice that, for any element  $[f] \in \pi_n(S^n) \cong \mathbb{Z}$ , we have by definition that  $h_{S^n}([f]) = f_*([\alpha]) = \deg(f) \cdot [\alpha]$ . Since  $[\text{Id}_{S^n}] \in \pi_n(S^n)$  is s.t.  $\text{Id}_{S^n}$  has degree 1 because it induces the identity isomorphism on  $H_n(S^n) \cong \mathbb{Z}$ , we have then the surjectivity.  $\square$

*Proof.* (c) The two maps trivially agree up to sign, for they are isomorphisms from  $\pi_n(S^n) \cong \mathbb{Z}$  to  $H_n(S^n) \cong \mathbb{Z}$ .  $\square$

## Exercise 3

*Proof.* By the usual argument about cellular maps,  $\pi_m(X) = 0 = \pi_m(X) \otimes \mathbb{Q}$  for  $m < n$ .

By [1, thm. 12.1], all of the homotopy groups of  $X$  are abelian and finitely generated, hence they can be described as  $\pi_k(X) = \mathbb{Z}^r \oplus \pi_k(X)^{\text{tors}}$  for some  $r \in \mathbb{N}$ . Also,  $\pi_k(X) \otimes \mathbb{Q} = \mathbb{Q}^r$ . We will then work with the Hurewicz theorem mod  $\mathcal{C}$ , where  $\mathcal{C}$  is the class of torsion abelian groups.

First of all, we shall compute  $H_n(X) \otimes \mathbb{Q}$  for all  $n$  and  $k$ .

Using the description of  $X$  as a finite CW-complex, we see that its homology corresponds to the homology of the cellular chain complex  $(C_\bullet, \partial)$ , where  $C_0 = \mathbb{Z}$ ,  $C_n = \mathbb{Z}$ ,  $C_{n+1}$  and  $C_{n+1} \xrightarrow{\partial_n} C_n$  is given by  $m \mapsto km$ . It follows that  $H_n(X) = \mathbb{Z}/k\mathbb{Z} \in \mathcal{C}$ ,  $H_0(X) = \mathbb{Z}$ ,  $H_m(X) = 0$  for  $m \neq 0, n$  and  $H_0(X) \otimes \mathbb{Q} = \mathbb{Q}$ ,  $H_t(X) \otimes \mathbb{Q} = 0$  for all other  $t$ .

By Hurewicz,  $\pi_n(X) = H_n(X) = \mathbb{Z}/k\mathbb{Z}$  and  $\pi_n(X) \otimes \mathbb{Q} = 0$ .

We also have that  $P_n X$  is a  $K(\mathbb{Z}/k\mathbb{Z}, n)$ . We may then consider the fibration sequence  $X\langle n \rangle \rightarrow X \rightarrow K(\mathbb{Z}/k\mathbb{Z}, n)$ , which then gives us the following one:  $\Omega K(\mathbb{Z}/k\mathbb{Z}, n) = K(\mathbb{Z}/k\mathbb{Z}, n-1) \rightarrow X\langle n \rangle \rightarrow X$ .

By [1, lemma 13.16],  $H_n(K(\mathbb{Z}/k\mathbb{Z}, m)) \in \mathcal{C}$  for all  $m \in \mathbb{N}_{>0}$  and by [1, lemma 13.15] the same goes for  $\pi_{n+1}(X\langle n \rangle) = \pi_{n+1}(X)$ . It follows that  $\pi_{n+1}(X) \otimes \mathbb{Q} = 0$ .  $\square$

## References

- [1] Heuts Gijs and Meier Lennart. *Algebraic Topology II*. 2019.