

# Self-Sustained Penrose Excitation of Accretion Disks: A Spin-Regulated Mechanism for Super-Eddington Quasar Luminosities

JUN WAKABAYASHI<sup>1</sup>

<sup>1</sup>*Independent Researcher, Japan*

## ABSTRACT

We present a high-spin, near-threshold equatorial radiative channel that (i) sustains super-Eddington luminosities and (ii) self-consistently enforces a sub-unity spin ceiling, explaining why spins do not exceed unity in practice. Compactness-limited transparency ( $\ell$ ) gates the observable power, and a simple torque balance links the luminosity to the same control that sets the ceiling. The framework is immediately testable via three co-occurring hooks at high spin and high accretion rate: (1) a 20–120 keV high-energy shoulder favored over a featureless continuum by  $\Delta\text{AICc}$  or  $\Delta\text{BIC} \geq 6$ ; (2) lag hardening from X to UV/optical; and (3) equator-aligned X-ray polarization with a rising degree. The path to falsification is explicit: in vetted high-spin, high- $\lambda$  quasars, failure to recover at least two hooks—after excluding heavy absorption, extreme inclination, and high-compactness cases—would refute the scenario. We also note that rare geometries may allow partial direct leakage of the quasi-beam, potentially producing fast-variability events (speculative; see Sec. 6.4).

*Keywords:* Quasars — Accretion, accretion disks — Black hole physics — Relativistic processes

## EXECUTIVE SUMMARY

1. Near-extremal spins enable a compactness-gated *equatorial* radiative channel that sustains  $L/L_{\text{Edd}} > 1$  while self-limiting the spin.
2. The lever is repeated Penrose-like coupling at the ISCO-ergoregion interface, depositing power back into the disk and extracting angular momentum.
3. The framework predicts three joint hooks: a hard X/MeV shoulder (20–120 keV, obs), lag hardening (X→UV/optical), and equator-aligned X-ray polarization.
4. Screens remove confounders: Compton-thick absorption, extreme inclination, and high compactness ( $\ell < 30$  required).
5. **Decisive falsification criterion:** in vetted high-spin/high- $\lambda$  quasars, failure to recover  $\geq 2$  hooks (after screens) refutes the framework.

## 1. INTRODUCTION

This paper develops a concrete, spin-regulated mechanism — Self-Sustained Penrose Excitation (SSPE) — that can both power super-Eddington quasar luminosities and enforce a near-extremal spin ceiling, while remaining explicitly falsifiable by X-ray and polarimetric data.

Accreting supermassive black holes (SMBHs) display a remarkable range of radiative efficiencies and Eddington ratios. Standard thin-disk theory struggles to account for quasars radiating persistently at factors of a few above the Eddington limit without invoking either extremely fine-tuned parameters or poorly constrained “slim” accretion flows. At the same time, near-extremal Kerr spins appear to be common among luminous active galactic nuclei (AGN), yet the dimensionless spin parameter  $a_*$  is expected to remain bounded below unity by basic causal and thermodynamic arguments. Any viable model for the most luminous quasars must therefore accomplish two tasks at once: (i) provide an additional, repeatable energy source that can sustain  $L \sim 2\text{--}3 L_{\text{Edd}}$  (with occasional excursions to  $5\text{--}10 L_{\text{Edd}}$ ), and (ii) explain why the spin distribution shows an apparent “ceiling” just short of  $a_* = 1$ .

A second, independent tension has emerged from observations of jet-dominated, Blandford–Znajek (BZ) favorable systems. In several objects where powerful relativistic jets clearly dominate the large-scale energy budget, the thermal accretion disk appears more extended and hotter at large radii than expected from canonical BZ + thin-disk models. In other words, the disk seems to be “puffed up and overheated” on scales where standard radiative and viscous heating would predict a cooler, geometrically thinner structure. This combination — BZ-dominated power extraction *and* a conspic-

uously hot, inflated outer disk — suggests that a non-negligible fraction of the black-hole spin energy might be being redirected back into the equatorial flow.

*Tensions across probes: local fixes vs. global synthesis.*— Taken in isolation, each of several long-running tensions can be accommodated by source-by-source adjustments within existing frameworks:

- (i) hard X/MeV shoulders can be fit with multi-component Comptonization or corona-geometry variants;
- (ii) wavelength-dependent lag spectra can be absorbed into more complex reprocessing geometries;
- (iii) microlensing and reverberation sizes can be softened by modest disk thickening and radiative-transfer complications; and
- (iv) in jet-favorable systems, outer-disk heating can be attributed to jet feedback or environment-dependent illumination.

However, the recurring appearance of these tensions *in the same high-spin, high- $\lambda$  population* suggests that treating them as unrelated, probe-specific anomalies is an increasingly high-dimensional patchwork.

This paper frames SSPE as a single *equatorial inner-disk heating channel* that can be confronted with data across these disparate probes under a shared screening logic (compactness, absorption, inclination). The intent is not to declare uniqueness, but to provide a unifying, explicitly falsifiable target that invites coordinated scrutiny by the spectral-timing, microlensing/reverberation, and GRMHD/plasma communities.

This paper introduces and develops *Self-Sustained Penrose Excitation* (SSPE): a quasi-equatorial, Penrose-like energy-transfer channel that couples the ergoregion to the innermost accretion flow. When the Kerr spin exceeds a near-threshold value  $a_* \gtrsim a_{\text{th}} \simeq 0.97$ , a modest fractional coupling of the rotational-energy reservoir, parameterized by  $\epsilon_{\text{coup}}(a_*) \sim 10^{-2}$ – $10^{-1}$ , can continuously re-inject power into the inner disk while providing a counter-torque that self-limits the spin.

Our goal is to establish SSPE as a quantitatively testable hypothesis. First, we formalize the geometry and kinematics of the ISCO–ergoregion interface and introduce a minimal parameterization of the coupling efficiency and threshold (Section 2). Second, we present a semi-analytic parameter study exploring how  $L/L_{\text{Edd}}$ , duty cycle, and spin evolution respond to changes in  $\epsilon_{\text{coup}}$ , coupling scale, and effective screening of high- $\ell$  influx (Section 3). Third, we translate each observational hook into concrete signal-to-noise, model-selection, and polarimetric requirements, and state an explicit falsification criterion (Section 4). Fourth, we apply the

framework to BZ-favorable, jet-dominated quasars with anomalously hot/extended disks, arguing that equatorial feedback can inflate and overheat the outer disk even in such systems (Section 6.3).

*Scope and modeling stance.*—We present a phenomenological framework: the microphysics of beam formation, transport, and equatorial deposition is intentionally kept agnostic, while the *observable consequences* of such coupling are made explicit and falsifiable. This stance allows clean confrontation with data now and provides a clear interface to future GRMHD/plasma studies that could instantiate the coupling agent. Throughout we deliberately restrict attention to a single Kerr SMBH and its immediate accretion environment; issues of cosmological SMBH growth, seeding, and host-galaxy evolution are deferred to separate work.

*Boundary conditions are not fixed near  $a_* \rightarrow 1$ .*—Much of the literature treats the inner disk, magnetosphere, and ergoregion with effectively fixed cross-component boundary conditions. We instead emphasize that near-extremal spin the boundaries themselves evolve: frame dragging enlarges and reshapes the ergoregion, the ISCO approaches it, and the equatorial return path becomes thin and resistive. In this coupled regime an equatorial current sheet is generically required by the global field topology; shear and flux loading drive plasmoid-dominated reconnection; and the resulting split redistributes  $(E, L)$  so that a negative-energy branch is absorbed by the hole while a gain branch vents as a narrow equatorial quasi-beam, part of which returns to heat the inner disk.

## 2. THEORETICAL FRAMEWORK

**In brief:** Near-threshold Kerr coupling at the ISCO–ergoregion interface supplies disk heating while extracting spin, under a compactness gate  $\ell < 30$ .

---

All elements invoked below act outside the event horizon; the coupling operates in the ergoregion and deposits energy in the equatorial flow, avoiding assumptions about interior or singularity-scale physics. **Abbreviations and references.** We use *FFE* for the force-free, magnetically dominated limit ( $\rho_e \mathbf{E} + \mathbf{J} \times \mathbf{B} \approx 0$ ,  $\mathbf{J} \cdot \mathbf{E} = 0$ ) and *GRMHD* for ideal magnetohydrodynamics evolved on a Kerr background (S. S. Komissarov 2004; S. E. Gralla & T. Jacobson 2014; W. E. East & H. Yang 2018; Z. Pan 2018). We use *PIC* for first-principles kinetic simulations that resolve reconnection and plasmoid formation in the ergospheric current sheet (K. Parfrey et al. 2019; A. Bransgrove et al. 2021). Throughout we group references as (FFE/GRMHD) versus (PIC).

### 2.1. Kerr Energy Reservoir

**In brief:** A near-extremal Kerr hole stores  $\sim 0.29Mc^2$  of rotational energy that can power luminous episodes.

For a Kerr black hole of mass  $M$  and spin  $a_*$ , the extractable rotational energy is

$$E_{\text{rot}}(a_*) = \left[ 1 - \sqrt{\frac{1}{2} \left( 1 + \sqrt{1 - a_*^2} \right)} \right] Mc^2, \quad (1)$$

reaching  $\sim 0.29 Mc^2$  as  $a_* \rightarrow 1$ .


### 2.2. Spin-Triggered Coupling

**In brief:** A steep activation above  $a_{\text{th}} \simeq 0.97$  sets both the luminosity lever and a self-imposed spin ceiling.

We posit a threshold spin  $a_{\text{th}}$  above which an ergoregion-coupled agent activates. The effective coupling follows

$$\epsilon_{\text{coup}}(a_*) = \begin{cases} 0, & a_* \leq a_{\text{th}}, \\ \epsilon_{\text{max}} \left( \frac{a_* - a_{\text{th}}}{1 - a_{\text{th}}} \right)^n, & a_* > a_{\text{th}}, \end{cases} \quad (2)$$

where  $\epsilon_{\text{max}}$  is the saturation cap and  $n$  controls activation sharpness. **Activation threshold.** The equatorial coupling turns on steeply once the spin exceeds a threshold  $a_{\text{th}}$ . In our fiducial calibration, the rise is rapid near  $a_* \simeq 0.97$ , setting the lever for both luminosity boost and spin regulation (cf. Eqs. (2)–(11)).



**Figure 1. Activation turns on steeply above the threshold.** Peak  $L_{\text{tot}}/L_{\text{Edd}}$  rises rapidly once  $a_* > a_{\text{th}}$ , setting the lever for luminosity and spin regulation (cf. Eqs. (2)–(11)).

Unless noted, we adopt the following fiducials for figures and estimates:  $M = 10^9 M_\odot$ ,  $\eta_{\text{acc}} = 0.1$ ,  $\epsilon_{\text{max}} = 0.1$ ,  $n = 2$ ,  $R_0 = 10^2$ , and  $\alpha = 50$ ; the illustrative threshold is  $a_{\text{th}} \simeq 0.97$ . *Micro-bridge.* While we keep microphysics agnostic, the fiducial range  $\epsilon_{\text{coup}} \sim 10^{-2}$ – $10^{-1}$  is broadly consistent with energy-release fractions seen in near-horizon, plasmoid-dominated reconnection in recent GRMHD studies. For the present framework we only assume that  $\epsilon_{\text{coup}}$  rises sharply above  $a_{\text{th}}$  and saturates below  $\epsilon_{\text{max}}$ ; detailed calibration is left for future simulations.

*Phenomenological parameters and physical ranges.*—The cap  $\epsilon_{\text{max}}$  limits equatorial deposition efficiency by energy-budget and pair-compactness constraints;  $n$  controls activation sharpness above  $a_{\text{th}}$  as an effective criticality index of the coupling geometry.  $(R_0, \alpha)$  regulate the dissipation footprint to maintain transparency, trading compactness against reprocessing. We restrict these to physically plausible ranges and view them as interfaces for future GRMHD/plasma calibration, not curve-fitting knobs.

*Sensitivity and saturation.*—The activation  $\epsilon_{\text{coup}}(a_*)$  is intentionally steep:  $(a_{\text{th}}, n)$  set the narrowness of the “spin ceiling”. In practice  $\epsilon_{\text{coup}}$  and the leakage fraction depend nonlinearly on flux loading and  $\dot{M}$ , with possible saturation arbitrarily close to the extremal limit. We therefore treat  $(a_{\text{th}}, n, \epsilon_{\text{max}}, \alpha)$  as calibration parameters to be fixed by future GRMHD-in-ergoregion testbeds.

*Power partition.*—We allow the extracted rotational power to partition between a polar BZ jet and an equatorial feedback channel:

$$P_{\text{ext}} = P_{\text{jet}} + P_{\text{eq}}, \quad f_{\text{BZ}} + f_{\text{eq}} = 1.$$

Near  $a_* \rightarrow 1$ ,  $f_{\text{eq}}$  can become substantial (radio-quiet, radiation-dominated states), while  $f_{\text{BZ}}$  dominates in

radio-loud systems; hybrid states are possible in transient MAD-like regimes.

### 2.3. Mechanism sketch: a leaky equatorial return path

**In brief:** Equatorial reconnection ejecta split in the ergoregion; a negative-energy branch is absorbed while a quasi-beam returns to heat the inner disk.

---

*Projectile sufficiency.*—Reconnection ejecta (plasmoids) in the ergoregion carry specific energy  $E$  and angular momentum  $L$ . A Penrose split requires a branch with  $E - \Omega_H L < 0$  (horizon condition).<sup>2</sup> Tension-driven redistribution during ejection changes  $L$  by  $\Delta L \sim \mathcal{O}(r_g v_\phi)$  and reconnection outflows can reach  $v \sim 0.1\text{--}0.5c$  in high- $S$  sheets; hence tens-of-percent shifts in  $L$  suffice to place a fraction on the negative-energy branch, with the complement forming the *positive-energy branch* (equatorial quasi-beam). Near-extremal spin brings the ISCO into close contact with the ergoregion and enforces an equatorial return current sheet at their interface. Shear and flux loading thin the sheet until it becomes tearing-unstable; reconnection then ejects plasmoids as a chain of narrow, equator-following pulses. These ejecta provide the “projectile” required by a Penrose-like energy split: within the ergoregion, a fraction of the flow is placed on negative-energy-at-horizon trajectories while the counterpart gains energy and escapes. The extraction condition is simply

$$E - \Omega_H L < 0. \quad (3)$$

cf. [R. Penrose 1969](#); [R. Penrose 2002](#). so the black hole’s rotational energy pays for the escaping branch. The centrifugal barrier and toroidal tension form an equatorial nozzle, so a growing fraction of the circuit power vents as a collimated equatorial quasi-beam while the DC return still closes globally. A modest back-flow coupling ( $\epsilon_{\text{coup}} \sim 10^{-2}\text{--}10^{-1}$ ) suffices to heat the inner disk and regenerate magnetic flux, closing a self-sustained loop. The active zone and leakage fraction increase monotonically with spin (and also depend nonlinearly on flux loading and accretion rate), with possible saturation arbitrarily close to the extremal limit.

<sup>2</sup> Notation follows [Fig. 2](#).



**Figure 2. Equatorial Penrose trigger: negative vs. positive branches.**

Plasma from the inner disk/plunging region enters the ergosphere and splits kinematically: a *negative-energy branch* with  $E - \Omega_H L < 0$  falls through the *event horizon* ( $r_H$ ), while a *positive-energy branch* emerges as an equatorial quasi-beam that reheats the inner rim near  $r_{\text{ISCO}}$  (R. Penrose 1969; R. Penrose 2002). Landmarks—ergosurface,  $r_{\text{ISCO}}$ , and the equatorial return current sheet (blue ribbon)—are indicated; branches are shown as *red solid* (positive) and *blue dashed* (negative) to match Eq. (3).

(FFE/GRMHD: S. S. Komissarov 2004; W. E. East & H. Yang 2018; Z. Pan 2018; PIC: K. Parfrey et al. 2019; A. Bransgrove et al. 2021).

*Why an equatorial, quasi-collimated branch?*—Near  $a_* \rightarrow 1$  the combination of (i) the equatorial centrifugal barrier, (ii) strong toroidal fields generated by shear (hoop stress), and (iii) pressure deficits carved by intermittent reconnection outflows forms an “equatorial nozzle”. The *positive-energy branch* therefore propagates as a narrow, radiation-supported quasi-beam rather than escaping along the poles. Because the transport remains largely collisionless/Poynting-dominated until it reprocesses in the inner disk, and because the dissipation footprint is extended ( $R_{\text{eff}} \sim 10^2\text{--}10^3 r_g$ ), the pair compactness along the beam stays low, consistent with the transparency requirement of Eq. (6).

#### 2.4. Self-Sustained Penrose Excitation

**In brief:** A modest feedback fraction  $\epsilon_{\text{coup}} \sim 10^{-2}\text{--}10^{-1}$  closes the loop, sustaining bright states without pair-choking.

---

The mean extraction power is  $\langle P_{\text{ext}} \rangle = E_{\text{rot}}/\tau$ , and with duty cycle  $d$  the instantaneous power during active phases is  $P_{\text{ext}} \sim \langle P_{\text{ext}} \rangle/d$ . Here  $\tau$  denotes the effective extraction e-folding timescale of the rotational-energy reservoir, i.e.,  $\langle P_{\text{ext}} \rangle = E_{\text{rot}}/\tau$  using Eq. (1).

#### 2.5. Disk Dissipation and Scale

**In brief:** An extended footprint  $R_{\text{eff}} \sim 10^2\text{--}10^3 r_g$  keeps  $\ell$  low so the hard component remains transparent.

---

A fraction  $\epsilon_{\text{coup}}(a_*)$  of  $P_{\text{ext}}$  is deposited into the disk:

$$L_{\text{self}} = \epsilon_{\text{coup}} P_{\text{ext}}, \quad (4)$$

and the dissipation spreads over an effective radius

*Self-transparency (physical note).*—The scaling  $R_{\text{eff}} \approx R_0[1 + \alpha(\epsilon_{\text{coup}}/\epsilon_{\text{max}})]r_g$  phenomenologically captures geometric spreading, multi-zone deposition, and increased scattering mean free paths as the equatorial quasi-beam heats and rarefies the inner disk corona. Our results require  $R_{\text{eff}} \sim 10^2\text{--}10^3 r_g$  during bright episodes to keep the pair compactness low (cf. Sec. 2.6).

$$R_{\text{eff}} \approx R_0 \left[ 1 + \alpha \left( \frac{\epsilon_{\text{coup}}}{\epsilon_{\text{max}}} \right) \right] r_g, \quad (5)$$

with  $r_g = GM/c^2$ , fiducial  $R_0 \sim 10^2$  and  $\alpha \sim 50$ .

#### 2.6. Transparency (Compactness Constraint)

**In brief:** Transparency requires  $\ell \lesssim 30$ , linking power deposition to geometry and spectrum.

Transparency requires pair compactness

$$\ell = \frac{L_{\text{self}} \sigma_T}{4\pi R_{\text{eff}} m_e c^3} \lesssim 30, \quad (6)$$

which couples Eqs. (4) and (5) and motivates  $R_{\text{eff}} \sim 10^2\text{--}10^3 r_g$  during bright episodes. We adopt  $\ell \lesssim 30$  as a conservative transparency threshold following classic compactness arguments; the precise value depends on geometry and spectrum and can be re-tuned in data applications. See, e.g., (R. Svensson 1984; A. P. Lightman & A. A. Zdziarski 1987).

#### 2.7. Spin and Mass Evolution

**In brief:** Torque balance with  $\Omega_H$  enforces  $a$ -drift toward an equilibrium  $a_{\text{eq}}$  below unity.

---

The horizon angular frequency is

$$\Omega_H = \frac{a_* c^3}{2GM(1 + \sqrt{1 - a_*^2})}, \quad (7)$$

where  $r_H = r_g(1 + \sqrt{1 - a_*^2})$  and  $r_g = GM/c^2$ . Evolution obeys

$$\frac{dM}{dt} = \frac{dM_{\text{acc}}}{dt} - \frac{P_{\text{ext}}}{c^2}, \quad (8)$$

$$\frac{dJ}{dt} = \frac{dJ_{\text{acc}}}{dt} - \frac{P_{\text{ext}}}{\Omega_H}, \quad (9)$$

$$\frac{da_*}{dt} = \frac{c}{GM^2} \frac{dJ}{dt} - 2a_* \frac{1}{M} \frac{dM}{dt}. \quad (10)$$

#### 2.8. Net Luminosity

**In brief:** Total luminosity combines accretion and equatorial feedback, generically yielding  $L/L_{\text{Edd}} \sim 2\text{--}3$ .


---

The total luminosity is

$$L_{\text{tot}} = L_{\text{acc}} + L_{\text{self}}, \quad L_{\text{acc}} \approx \eta_{\text{acc}} \left( \frac{dM_{\text{acc}}}{dt} \right) c^2, \quad (11)$$

which links the dynamical solution (Eqs. (8)–(10)) to observables.

*Energetic sanity check (one-line).*—With  $M_{\text{BH}} = 10^9 M_\odot$  [ $Mc^2 \simeq 1.8 \times 10^{63}$  erg] and  $E_{\text{rot}} \sim 0.1 Mc^2$ , a reservoir e-fold  $\tau = 10^7$  yr,  $\epsilon_{\text{coup}} = 0.05$ , and duty  $d = 0.2$  yield  $L_{\text{self}} \sim (E_{\text{rot}}/\tau)(\epsilon_{\text{coup}}/d) \approx 1.4 \times 10^{47}$  erg s $^{-1} \sim 1.1 L_{\text{Edd}}$ , and  $2\text{--}3 L_{\text{Edd}}$  when combined with concurrent accretion, while  $da_*/dt < 0$  prevents overspin.



**Figure 3. Spin evolution and equilibrium vs. coupling.** *Top:* Spin drift rate  $da_*/dt$  from Eqs. (8)–(10) for representative accretion states; negative values drive  $a_*$  below unity. *Bottom:* Equilibrium spin  $a_{eq}$  obtained from  $da_*/dt = 0$  as a function of  $\epsilon_{coup}$ ; ... the shaded band  $\epsilon_{coup} \sim 10^{-2}$ – $10^{-1}$  is the fiducial range used elsewhere. See also Eq. (7).



**Figure 4. Spin-down on Myr scales under equatorial extraction.** Time evolution of the spin parameter  $a_*(t)$  obtained by integrating Eqs. (8)–(9)–(10) with horizon frequency (7) for a fiducial coupling  $\epsilon_{coup}$  (gold). The monotonic drift toward  $a_{eq} < 1$  illustrates the self-limiting ceiling; see Sec. 6.



**Figure 5. Equilibrium timescale and luminosity mapping.** *Top:* Time to equilibrium  $\tau_{eq}$  versus coupling  $\epsilon_{coup}$ , computed from the spin-evolution system Eqs. (8)–(10) with horizon frequency (7); the shaded band  $\epsilon_{coup} \sim 10^{-2}$ – $10^{-1}$  is the fiducial range used elsewhere. *Bottom:* Total luminosity at equilibrium,  $L_{tot}/L_{Edd}$  from Eq. (11) evaluated at  $a_* = a_{eq}$ , showing a typical 2–3 band (blue) with labeled  $\epsilon_{coup}$  values along the trend.

### 3. SEMI-ANALYTIC PARAMETER STUDY

**In brief:** Model trends map directly to hooks; figures follow their defining equations to prevent misalignment.

The semi-analytic exploration above yields three robust trends that do not depend on the microphysical details of how the equatorial branch is realized. First, once the coupling activates above the threshold spin  $a_* \gtrsim a_{th}$ , even a modest efficiency  $\epsilon_{coup} \sim 10^{-2}$ – $10^{-1}$  can supply an inner-disk feedback luminosity  $L_{self} = \epsilon_{coup} P_{ext}$  that is comparable to  $L_{Edd}$  for plausible reservoir and duty-cycle choices, so that the combined output  $L_{tot} = L_{acc} + L_{self}$  generically reaches the observed  $L/L_{Edd} \sim 2$ – $3$  band. Second, the same extraction term necessarily carries angular momentum and therefore introduces a counter-torque; integrating the minimal evolution system generically drives  $a_*$  toward an equilib-



rium  $a_{\text{eq}} < 1$  on Myr-scale equilibration times, with stronger coupling yielding a lower  $a_{\text{eq}}$ . Third, compactness links the observability of the hard component to geometry: the requirement  $\ell \lesssim 30$  couples  $(\epsilon_{\text{coup}}, d)$  to the dissipation footprint  $R_{\text{eff}}$ , carving a transparency-limited window in which bright states remain pair-safe. These trends map directly onto the observational program in Sec. 4. In the compactness-safe window (i.e., after screening out high- $\ell$  states), SSPE predicts that the hard component associated with equatorial feedback should be simultaneously (i) spectrally visible as a high-energy shoulder, (ii) temporally imprinted via lag-energy hardening consistent with inner-disk reheating and reprocessing, and (iii) geometrically encoded through disk-parallel hard X-ray polarization. Because activation is steep near  $a_{\text{th}}$ , the hooks are expected to concentrate in the high-spin, high- $\lambda$  subset rather than appearing as unrelated, source-by-source anomalies. Conversely, failure to recover at least two hooks in vetted high-spin/high- $\lambda$  quasars (after excluding heavy absorption, extreme inclination, and compactness-choked cases) constitutes a decisive falsification criterion. Finally, the same bookkeeping clarifies why SSPE should be treated as an equatorial complement to polar BZ extraction: across states, a changing partition between  $P_{\text{BZ}}$  and  $P_{\text{eq}}$  implies an anti-correlation between jet dominance and equatorial reheating diagnostics, with rare geometries potentially allowing partial direct leakage discussed separately in Sec. 6.4.



#### 4. OBSERVATIONAL HOOKS AND FALSIFIABILITY

In this paper we treat SSPE as a concrete, falsifiable working hypothesis rather than a completed theory. The microphysics of the coupling between the ergoregion, quasi-equatorial beams, and the inner disk is intentionally left agnostic and absorbed into a small set of phenomenological parameters. What is fixed — and offered to observers and simulators — are three testable hooks and an explicit falsification criterion: (1). A hard X/MeV “shoulder” in the  $\sim 20$ – $120$  keV band (Hook 1), (2). Lag hardening in Fourier-resolved timing, in which harder bands lead softer bands over a well-defined frequency range (Hook 2), and (3). X-ray polarization preferentially aligned with the disk plane in the hard component (Hook 3). These are complemented by a population-level falsification criterion: high-spin, high-Eddington-ratio, radio-quiet quasars must not *systematically* lack Hooks 1–3. The goal of this paper is not to close the question, but to make it straightforward for the community to confirm, refine, or decisively rule out SSPE using existing and near-future data.

##### 4.1. Observational Hooks

**In brief:** After screens, three hooks: MeV shoulder, lag hardening, and equator-aligned polarization—should co-occur in the bright, high-spin subset.

---

We emphasize that the observational hooks and thresholds were specified *a priori* as a falsification-oriented protocol, and only subsequently confronted with existing reports and archival indicators to assess feasibility. This ordering is intended to minimize post hoc tuning and to keep the framework portable across instruments and subfields.

**Why this matters.** What to look for and how to falsify, in plain terms.

**Recall.** Activation  $\epsilon_{\text{coup}}(a_*)$  and power  $P_{\text{eq}} = \epsilon_{\text{coup}} \dot{M} c^2 g(\ell)$  (cf. Eqs. (2), (11)), with transparency set by compactness  $\ell$  (Eq. (6)). These establish the trends in the plots below.

**Screens (apply first).** Exclude heavy absorption (Compton-thick etc.), extreme inclination, and high compactness ( $\ell \gtrsim 30$ ).

**Hooks (apply jointly under the screens).**

- (1) **Hard X/MeV shoulder (20–120 keV, obs):** model comparison favors a shoulder over a featureless baseline by  $\Delta\text{AICc}$  or  $\Delta\text{BIC} \geq 6$  (rest  $\sim 0.3$ – $1$  MeV at  $z \gtrsim 6$ ).

- (2) **Lag hardening (X→UV/optical):** energy-resolved lags increase *monotonically* with photon energy within an epoch.
- (3) **Equator-aligned X-ray polarization (2–10 keV):** EVPA  $\parallel$  equator; polarization degree increases toward high spin/high  $\lambda$ .

**Falsification.** In vetted high-spin, high- $\lambda$  quasars, *failure to recover at least two hooks* after the screens above *refutes* the framework.

##### Observer Quickstart (Tier-1 archival screening; non-committal).

1. **Hook 1 (spectral):** fit a baseline (e.g., cutoff power-law) and a shoulder-enabled alternative over 20–120 keV, and report preference using  $\Delta\text{BIC}$  or  $\Delta\text{AICc}$  (threshold:  $\geq 6$ ).
2. **Hook 2 (timing):** compute the lag–energy relation in the same epoch and test only the sign/monotonicity, requiring  $d(\text{lag})/d \log E > 0$  over the adopted band.
3. **Decision:** sources satisfying both Tier-1 hooks motivate Tier-2 follow-up; persistent null results in vetted high-spin/high- $\lambda$ , radio-quiet samples contribute directly to falsification under Sec. 4.1.

**At a glance.** How power repartitions at high  $\lambda$  across mechanisms.


**Table 2.** Operational hook specification

Item	Observable	Band / Product	Pass criterion	Primary confounders
Hook 1	Hard X/MeV shoulder	20–120 keV	$\Delta\text{BIC} \geq 6$	model degeneracy
Hook 2	Lag hardening	lag vs. $E$ slope	positive/monotone	dilution
Hook 3	Equatorial polarization	2–10 keV	EVPA $\parallel$ disk	jet contamination

NOTE—Screens applied first; failure to recover  $\geq 2$  hooks falsifies SSPE.

**Table 1. Bottom line:** the equatorial channel sustains  $L/L_{\text{Edd}} > 1$  and imposes a spin ceiling  $a_{\text{eq}} \approx a_{\text{th}}$ , complementary to slim-disk (no ceiling) and BZ/MAD (polar power,  $a_{\text{eq}} \sim 0.5\text{--}0.7$ ).

	$L/L_{\text{Edd}} > 1$	spin ceiling	channel
Slim disk	yes (advective)	no	radiation (disk)
BZ/MAD	indirect	yes ( $a_{\text{eq}} \sim 0.5\text{--}0.7$ )	polar Poynting (jet)
<b>This work</b>	<b>yes (equatorial)</b>	<b>yes (<math>a_{\text{eq}} \approx a_{\text{th}}</math>)</b>	<b>equatorial Penrose-like</b>



**Figure 6. Hook observables vs. equilibrium spin.** *Left:* Fe K $\alpha$  reverberation lag  $\tau_{\text{FeK}}$  (in  $r_g/c$ ) as a function of  $a_{\text{eq}}$ ; lags increase toward lower  $a_{\text{eq}}$  (stronger coupling). *Right:* Observable fraction  $f_{\text{hook}}$  (hard X/MeV shoulder or 2–10 keV polarization) vs.  $a_{\text{eq}}$  from the toy population mapping; the fraction rises steeply toward low  $a_{\text{eq}}$ . See Sec. 4.1 and App. A for definitions and pass criteria.

## 5. POSITION IN CONTEXT

**In brief:** SSPE is not a replacement for existing spin-extraction channels, but an equatorial complement. Thorne-limit arguments remain valid in their original thin-disk regime; BZ jets remain the natural outlet for MAD, strongly magnetized, radio-loud systems. SSPE instead targets the high- $\lambda$ , high-spin, radio-quiet sector where neither tool fully constrains the phenomenology.

Classical spin-evolution work shows that radiation capture in a razor-thin Novikov–Thorne disk drives an asymptotic limit  $a_* \simeq 0.998$  (K. S. Thorne 1974). We regard this as a clean limiting case, not a universal ceiling: real quasars host geometrically thick, magnetized, outflow-bearing inner flows that violate the assumptions of the original derivation. Likewise, the Blandford–Znajek (BZ) mechanism provides a robust path for launching powerful polar jets in magnetically arrested disks (e.g., R. D. Blandford & R. L. Znajek 1977; A. Tchekhovskoy et al. 2011), and can efficiently bleed spin energy in radio-loud, MAD-like systems. However, BZ by itself does not explain the combination of high inferred spins, high Eddington ratios, and weak radio emission seen in many luminous quasars.

In this landscape, SSPE is best viewed as an *equatorial exhaust channel*. When the ISCO approaches the ergoregion and the inner flow becomes thick and strongly coupled, a quasi-equatorial beam can transfer rotational energy back into the inner disk and cocoon. This plausibly supports mildly to strongly super-Eddington luminosities while enforcing an effective “spin ceiling”  $a_* \lesssim 0.97$ – $0.99$  for systems that lack a powerful jet. BZ and SSPE then form a see-saw: polar extraction dominates in MAD, radio-loud states, while equatorial extraction dominates in high- $\lambda$ , radio-quiet states. The observational hooks and falsification criterion in Sec. 4.1 are designed precisely to test this equatorial branch.

In what follows, we use the shorthand “see-saw” to denote an anti-correlated power partition between the polar and equatorial channels, i.e.,  $P_{\text{BZ}}$  increasing as  $P_{\text{eq}}$  decreases, at fixed spin-energy budget.

## 6. DISCUSSION

### 6.1. *Synthesis across probes*

Taken in isolation, each observational tension motivating SSPE admits a plausible local explanation. The core claim here is *not* that such explanations are impossible, but that their repeated co-occurrence in the same high-spin, high- $\lambda$ , often radio-quiet subset makes a single shared inner-disk heating channel a lower-complexity hypothesis. SSPE is offered as such a channel in a form that can be rejected decisively: after applying the screens (absorption, inclination, compactness), the high-spin/high- $\lambda$  subset should recover at least two of the three hooks (Sec. 4.1); systematic failure would support  $f_{\text{eq}} \rightarrow 0$  and render SSPE unnecessary.

**In brief:** The equatorial branch complements polar BZ; an anti-correlated partition is expected across states.

---


These bounds align with Fig. 7–8 and are sufficient to reach  $L/L_{\text{Edd}} \sim 2\text{--}3$  with rarer  $5\text{--}10\times$  excursions. Slim disks allow modest super-Eddington flows but no spin ceiling. BZ/MAD explain jet power yet not the radiative dominance of radio-quiet quasars. This framework ties hyperluminous output and the spin ceiling via near-extremal, equatorial coupling. Consistent with GRFFE/GRMHD studies, an *equatorial return-current sheet forms within the ergoregion* and becomes plasmoid-unstable; detached plasmoids unavoidably split into branches with  $E - \Omega_H L \lesssim 0$  (captured vs. escaping), enabling rotational-energy extraction via a Penrose-like radiative channel (R. Penrose 1969; R. Penrose 2002; S. S. Komissarov 2004; W. E. East & H. Yang 2018; Z. Pan 2018; K. Parfrey et al. 2019; A. Bransgrove et al. 2021).

**Relation to Blandford–Znajek jets.** BZ is polar/Poynting-dominated; our mechanism is equatorial/radiative. Hybrid states and an anti-correlated partition are expected; counterexamples (simultaneously strong jets and high radiative output) can occur in transitional MAD-like regimes. Importantly, SSPE is not posed as a competitor to BZ, but as an equatorial channel that can coexist with the global magnetospheric topology required for BZ. In this sense, SSPE may utilize the equatorial return current and inner-disk coupling present in BZ-favorable configurations in principle. The relevant emission / deposition radius  $R_{\text{eff}}$  for the SSPE-driven equatorial beam is not known a priori: it depends on microphysics that current GRMHD and kinetic simulations have not yet fully resolved. In this paper we therefore treat  $R_{\text{eff}}$  as a sensitivity parameter rather than a fitted quantity.


### 6.2. *Parameter space for quasi-beam transparency*

Figure 7–8 illustrates how the compactness screen  $\ell = \ell_{\text{crit}}$  carves the  $(\epsilon_{\text{coup}}, d)$  plane for two representative choices of  $R_{\text{eff}}$ ; the figure is meant as a diagnostic guide, not as a prediction for a specific object. Crucially, the map is an *upper-limit transparency bound*: it states how large the dissipation footprint may be *if* an  $\ell < 30$  quasi-beam component is to escape pair-choking, without assuming where or how the beam is launched.

Interpreted conservatively, increasing  $R_{\text{eff}}$  enlarges the *allowed* transparent window, not the likelihood of direct escape.



**Figure 7. Transparency selects the viable high-luminosity regime.** Peak Eddington ratio  $L_{\text{tot}}/L_{\text{Edd}}$  (color) on the  $(\epsilon_{\text{coup}}, d)$  plane; the black contour marks  $\ell=30$  from Eq. (6) with components defined in Eqs. (4,5). For  $R_{\text{eff}} = 100 r_g$ , the allowed region lies *outside* the  $\ell=30$  contour, where  $L_{\text{tot}}/L_{\text{Edd}} \sim 2\text{--}3$  is typical with rarer  $5\text{--}10\times$  excursions; see App. A.



**Figure 8. A larger dissipation footprint widens the transparent window.** Peak  $L_{\text{tot}}/L_{\text{Edd}}$  (color) on the  $(\epsilon_{\text{coup}}, d)$  plane for a larger footprint  $R_{\text{eff}} = 1000 r_g$ . The  $\ell=30$  boundary (black; from Eq. (6) with components in Eqs. (4,5)) shifts so that higher peaks in  $L_{\text{tot}}/L_{\text{Edd}}$  remain compactness-safe, enlarging the viable region compared to Fig. 7.

### 6.3. Power partition and spin budget

We model the spin-extraction power as a partition between a polar Blandford–Znajek (BZ) channel and an equatorial feedback channel,

$$P_{\text{tot}} = P_{\text{BZ}} + P_{\text{eq}}, \quad (12)$$

$$P_{\text{BZ}} = f_{\text{BZ}} P_{\text{ext}}, \quad P_{\text{eq}} = f_{\text{eq}} P_{\text{ext}}, \quad (13)$$

$$f_{\text{BZ}} + f_{\text{eq}} = 1. \quad (14)$$

A practical implication is that SSPE can affect systems that appear, by their large-scale phenomenology, to be BZ-favorable. Even when the polar channel dominates the long-term power budget (i.e.,  $P_{\text{BZ}} \gg P_{\text{eq}}$  on average), a nonzero equatorial branch provides an internal heating pathway that acts directly on the disk photosphere rather than through external irradiation. In such states, the observable tension is not “a jet plus a thin disk,” but a jet coexisting with a disk that looks *too extended and too hot* for a strictly passive, radiatively efficient thin-disk interpretation—as suggested by microlensing/reverberation size inferences and wavelength-dependent lag amplitudes that exceed simple reprocessing expectations. Within SSPE, the same bookkeeping that permits strong jet power also permits intermittent equatorial reheating episodes whose signatures should correlate with the spectral-timing hooks (Sec. 4) while remaining compatible with radio-loudness. This is not an attempt to fit any single source; rather, it motivates a concrete null test: in jet-dominated quasars exhibiting anomalously large continuum sizes or overheated outer disks, the absence of *both* hard-component hooks (Hook 1/2) *and* disk-parallel hard-X polarization (Hook 3), after controlling for absorption and compactness screening, would disfavor equatorial feedback as the common cause.

This partition predicts an anti-correlation between radio-jet dominance and equatorial reheating diagnostics within the high- $a_*$ , high- $\lambda$  subset, with hybrid states possible in transient MAD phases. Our claim is modest: in the high-spin, high-Eddington subset,  $f_{\text{eq}}$  is statistically non-zero and sometimes dominant; in others,  $f_{\text{BZ}}$  may prevail. Outside this subset we make no claim.

### 6.4. Rare direct quasi-beam events as ergoregion probes

We stress that the following subsection is explicitly speculative and *not* required for the core SSPE framework: the primary observational program remains the hook-based spectral-timing-polarimetric tests in Sec. 4.1. **In brief:** In most SSPE-active systems the quasi-equatorial beam is reprocessed into Hook 1/2. In rare geometries a fraction may escape directly, producing ultra-fast, ultra-

hard flares that could offer an unusually direct probe of the ergoregion.

---

**Normal outcome: reprocessing into Hooks 1/2.** In the working picture adopted here, SSPE acts primarily as an *equatorial power channel*: a narrow quasi-beam extracts rotational energy near the ISCO–ergoregion interface and re-injects it into the inner disk and surrounding cocoon. Most of the time this energy is promptly reprocessed into the hard X/MeV shoulder (Hook 1) and lag-hardening signatures (Hook 2), with only modest direct leakage beyond the disk and torus. These reprocessed components constitute the primary hooks discussed in Sec. 4.1.

**Rare outcome: partial beam leakage.** However, there is no guarantee that the inner flow is always geometrically and optically thick in the relevant directions. In special configurations—for example, low  $H/R$  at the inner rim, local warps or gaps, or mild misalignment between the spin axis and the large-scale disk—a fraction of the quasi-equatorial beam may find a comparatively transparent channel and emerge without full reprocessing. In that case SSPE could produce *direct* signatures on timescales of a few  $r_g/c$ , much shorter than the thermal and viscous response of the disk.

**Qualitative signature set.** We emphasize that we do *not* attempt a full classification here, but the following features are natural in the present framework:

- (1) **Ultra-fast, ultra-hard flares:** isolated spikes or bursts with variability timescales  $\sim$  few  $r_g/c$ , spectral peaks extending into the hard X-ray / soft  $\gamma$ -ray band, and hardness ratios exceeding those of the quasi-steady shoulder.
- (2) **Hard-to-soft evolution:** individual events in which the hardest photons lead a softer tail, consistent with a narrow beam depositing energy into a thicker reprocessing layer.
- (3) **Transient polarization swings:** in principle, a direct beam component should exhibit strong polarization with EVPA tied to the local equatorial geometry; rapid changes in the effective beam direction could induce short-lived swings in hard X-ray polarization angle.

**Candidate hunting grounds.** At the level of working hypotheses, two regimes appear especially promising:

- (i) **Lamppost fits with extremely compact source heights** ( $h \sim$  a few  $r_g$ ). In some bright quasars and Seyferts, lamppost spectral fits drive the inferred source height to values  $h \sim$  few  $r_g$  or

lower, where the physical interpretation becomes ambiguous. Within SSPE such cases could instead correspond to quasi-equatorial beams emerging just above the horizon, rather than a compact source suspended on the spin axis.

- (ii) **High- $\lambda$  narrow-line Seyfert 1s (NLS1s).** These systems host relatively low-mass SMBHs accreting at high Eddington ratios and often display strong, rapid X-ray variability. A subset with very hard, short flares and strong high-energy excess would be natural laboratories for rare beam-leakage episodes.

We stress that these are *suggestive* classes, not firm identifications: detailed modeling and joint timing–spectral–polarimetric analyses are required before any specific source can be claimed as a direct quasi-beam detection candidate.

**Relationship to the main hooks.** From the perspective of this Paper, rare direct quasi-beam events are not an additional mandatory hook but a high-risk, high-reward extension. The primary falsifiable content of the SSPE framework remains the *co-occurrence* of Hooks 1–3 under the screening criteria of Sec. 4.1. A confirmed direct quasi-beam event would instead provide an exceptionally clean window into the ergoregion dynamics, placing much sharper constraints on the microphysics of the coupling than our phenomenological treatment assumes.

## 6.5. Validation Roadmap

**In brief:** Polarization–spin trends, intra-epoch lag slopes, and compactness screens lead; GRMHD+RT+population studies anchor the pipeline; Table 3 summarizes the minimal products and falsification criterion triggers.

---

### Observational fronts.

- (i) **Polarization vs. spin.** Test that EVPA is equator-aligned and that the degree rises with  $a_*/\lambda$ ; control for jet and scattering-cone geometries; report rank-correlation  $p$ -values and effect sizes.
- (ii) **Lag hardening.** Within single epochs, verify that  $X \rightarrow UV$ /optical lags increase monotonically with photon energy; mask absorption events; provide cadence/noise tests.
- (iii) **Spectral compactness screen.** Use cutoff-energy/pair proxies to flag high- $\ell$  cases (non-transparent) and separate them from the transparent sample used for hooks.

**Table 3. Validation roadmap matrix.** What can be tested now, and what would falsify SSPE under the stated screens.

Hook	Minimal data products	Executable now?	Population-level falsification criterion trigger
Hook 1	Hard X-ray spectrum (background-controlled) + model comparison (AICc/BIC)	<b>Yes</b>	After screens, vetted high-spin/high- $\lambda$ sample systematically prefers featureless continua (no shoulder)
	Simultaneous (or tightly interleaved) X-ray + UV/optical light curves + Fourier lags		After screens, lag-energy slopes fail to be positive/monotone within epochs across the bright subset
Hook 2	Hard-component X-ray polarimetry + EVPA reference to disk plane	<b>Partly</b>	After screens, EVPA shows no preference for equatorial alignment and no trend
Hook 3	+ spin/ $\lambda$ proxies		with inferred spin/ $\lambda$

NOTE—The primary falsification criterion used in this work is the joint-hook rule: absence of  $\geq 2$  hooks in vetted high-spin/high- $\lambda$  quasars after screens.



- (iv) **BZ–equatorial power partition (see-saw).** Using Sec. 6.3 as the bookkeeping ( $P_{\text{tot}} = P_{\text{BZ}} + P_{\text{eq}}$ ;  $f_{\text{BZ}} + f_{\text{eq}} = 1$ ), test whether states (or sources) with larger jet-dominance proxies (e.g., radio loudness, core dominance, or independent jet-power estimates) systematically show weaker equatorial reheating diagnostics (Hooks 1–3), at fixed spin/ $\lambda$  and after screens. Report partial correlations (controlling for inclination and absorption) and state-by-state transitions where available. This see-saw test is a cross-check on the power partition, not an additional hook in the primary falsification protocol.

### Simulation fronts.

- (i) GRMHD near the ergoregion. Resolve the equatorial return current sheet; quantify energy partition into positive/negative branches ( $E - \Omega_H L \gtrless 0$ ); compare with torque-balance trends.
- (ii) Radiative transfer with pairs. Propagate hard X/MeV emission through compactness-limited media; map when the shoulder survives vs. is quenched (link to  $\ell$  cut).
- (iii) Population modeling. Combine spin distributions with transparency and coupling priors to predict the fraction of sources passing  $\geq 2$  hooks.

*Additional falsifiers (secondary).*—The primary falsification protocol is the hook-based falsification protocol in Sec. 4.1. The following are *supporting* population-level patterns that, if established robustly, would further disfavor a non-zero equatorial fraction  $f_{\text{eq}}$ :

1. Systematic absence of the Hard-X/MeV shoulder (20–120 keV, obs) *and* systematically weak high-ionization lines (He II, NV, C IV) relative to the parent population.
2. Reverberation lags from X→UV showing no energy dependence (no inward reheating signature).
3.  $L_{\text{bol}}$  not anti-correlated (even weakly) with  $a_*$  within the bright subset (no self-regulated spin-down imprint).
4. Optical/X-ray polarization failing to favor equatorial angles or degrees when near-threshold spins are inferred.

If *two or more* items above hold simultaneously for the same high-spin, high- $\lambda$  subset, we would consider  $f_{\text{eq}} \rightarrow 0$  supported and this mechanism unnecessary.

## 7. CONCLUSION

**In brief:** Recovering  $\geq 2$  hooks under screens supports the framework; systematic absence in high-spin/high- $\lambda$  would falsify it.

---

Self-sustained Penrose excitation near extremal Kerr SMBHs can inject rotational energy into the accretion disk, yielding sustained 2–3× Eddington with rarer 5–10× episodes while enforcing a spin ceiling.

## APPENDIX

## A. OPERATIONAL PROTOCOL (SELF-CONTAINED)

**Screens.** Exclude Compton-thick, extreme inclination; require  $\ell < 30$ .

**Hook (i):** Shoulder passes if  $\Delta\text{AICc}$  *or*  $\Delta\text{BIC} \geq 6$  (20–120 keV obs).

**Hook (ii):** Lag hardening passes if  $d\tau/d\log E > 0$  and monotone within an epoch.

**Hook (iii):** X-ray polarization: EVPA  $\parallel$  equator; degree increases with spin/ $\lambda$ .

**Verdict.** After screens, absence of  $\geq 2$  hooks  $\Rightarrow$  falsified; else passes.

**Table 4. Candidate Gallery** — Representative objects; values are indicative.

Name	$z$	$M_{\text{BH}}/M_{\odot}$	$L/L_{\text{Edd}}$	Notes
TON 618	2.219	$\sim 6.6 \times 10^{10}$	$\sim 3$	Extremely massive; radio-loud; literature refs. <sup><i>a</i></sup>
J2157-3602	4.75	$\sim 3.4 \times 10^9$	$\gtrsim 10$	Hyper-luminous; super-Eddington episode indications. <sup><i>b</i></sup>
J0100+2802	6.30	$\sim 1.2 \times 10^{10}$	$\sim 2$	$z > 6$ luminous quasar. <sup><i>c</i></sup>
J0439+1634	6.51	$\sim 7 \times 10^9$	$\sim 2-3$	Possible lensing history. <sup><i>d</i></sup>

NOTE—Values are illustrative; per-object sourcing is out of scope.

<sup>*a*</sup>Virial-factor dominated systematics; radio-loud bias possible.

<sup>*b*</sup>Bolometric/line-width systematics  $\gtrsim 0.3$  dex.

<sup>*c*</sup>High- $z$  mass methods differ (reflection vs. continuum).

<sup>*d*</sup>Historical lensing debate; values assume de-lensing consensus.

## CAUTIONARY NOTE.

Claims here are modular and falsifiable. Alternative mechanisms may dominate in other classes.

## AUTHOR CONTRIBUTIONS

Conceptualization, modeling, analysis, visualization, and writing: J. Wakabayashi.

## COMPETING INTERESTS

The author declares no competing interests.

## DATA AND CODE AVAILABILITY

All figures can be regenerated from scripts in the accompanying repository; data sources and acquisition steps are documented in a README.

An archived OSF snapshot (Version 1.1) is available: <https://doi.org/10.17605/osf.io/62gzv>.

## COMMUNICATION AND MEDIA

Media note: The paper and its reproducibility package are the sole authoritative sources.

## ACKNOWLEDGMENTS

This work stands on decades of insight into black-hole accretion, spin, and energy extraction. I am indebted to the community that built the modern framework of quasar physics—from classical Penrose energy extraction and disk theory to spin-jet coupling and polarimetry—and to teams who made public data and tools available.

**Use of large language models.** Large language model assistants (Google Gemini and OpenAI ChatGPT) were used for drafting support (editing for clarity, formatting suggestions, and figure placement/LaTeX troubleshooting). No novel data, equations, or results were produced by these tools. No confidential or unpublished data were provided to them. All analysis, derivations, and conclusions are by the author, who takes full responsibility for the content; the models are not authors.

I also thank colleagues and readers who provided critical comments on early drafts. Any remaining errors are mine.

*Software:* `latexmk`, AASTeX701

## REFERENCES

- Abramowicz, M. A., Czerny, B., Lasota, J.-P., & Szuszkiewicz, E. 1988, *The Astrophysical Journal*, 332, 646, doi: [10.1086/166683](https://doi.org/10.1086/166683)
- Blandford, R. D., & Znajek, R. L. 1977, *Monthly Notices of the Royal Astronomical Society*, 179, 433, doi: [10.1093/mnras/179.3.433](https://doi.org/10.1093/mnras/179.3.433)
- Bransgrove, A., Beloborodov, A. M., & Parfrey, K. 2021, *Physical Review Letters*, 127, 055101, doi: [10.1103/PhysRevLett.127.055101](https://doi.org/10.1103/PhysRevLett.127.055101)
- Cackett, E. M., Bentz, M. C., & Kara, E. 2021, *iScience*, 24, 102557, doi: [10.1016/j.isci.2021.102557](https://doi.org/10.1016/j.isci.2021.102557)
- Connor, T., Stern, D., Bañados, E., & Mazzucchelli, C. 2021, *ApJL*, 922, L24, doi: [10.3847/2041-8213/ac37b5](https://doi.org/10.3847/2041-8213/ac37b5)
- East, W. E., & Yang, H. 2018, *Physical Review D*, 98, 023008, doi: [10.1103/PhysRevD.98.023008](https://doi.org/10.1103/PhysRevD.98.023008)
- Gralla, S. E., & Jacobson, T. 2014, *Monthly Notices of the Royal Astronomical Society*, 445, 2500, doi: [10.1093/mnras/stu1690](https://doi.org/10.1093/mnras/stu1690)
- Komissarov, S. S. 2004, *Monthly Notices of the Royal Astronomical Society*, 350, 427, doi: [10.1111/j.1365-2966.2004.07672.x](https://doi.org/10.1111/j.1365-2966.2004.07672.x)
- Lai, S., Wolf, C., Onken, C. A., & Bian, F. 2023, *MNRAS*, 521, 3682, doi: [10.1093/mnras/stad651](https://doi.org/10.1093/mnras/stad651)
- Lightman, A. P., & Zdziarski, A. A. 1987, *ApJ*, 319, 643
- Pan, Z. 2018, *Physical Review D*, 98, 043023, doi: [10.1103/PhysRevD.98.043023](https://doi.org/10.1103/PhysRevD.98.043023)
- Parfrey, K., Philippov, A. A., & Cerutti, B. 2019, *Physical Review Letters*, 122, 035101, doi: [10.1103/PhysRevLett.122.035101](https://doi.org/10.1103/PhysRevLett.122.035101)
- Penrose, R. 1969, *Nuovo Cimento Rivista Serie*, 1, 252
- Penrose, R. 2002, *General Relativity and Gravitation*, 34, 1141, doi: [10.1023/A:1016578408204](https://doi.org/10.1023/A:1016578408204)
- Reynolds, C. S. 2014, *SSRv*, 183, 277, doi: [10.1007/s11214-013-0006-6](https://doi.org/10.1007/s11214-013-0006-6)
- Sun, M., Xue, Y., Brandt, W. N., et al. 2020, *Astrophysical Journal*, 902, 7, doi: [10.3847/1538-4357/ab789e](https://doi.org/10.3847/1538-4357/ab789e)
- Svensson, R. 1984, *MNRAS*, 209, 175, doi: [10.1093/mnras/209.2.175](https://doi.org/10.1093/mnras/209.2.175)
- Tchekhovskoy, A., Narayan, R., & McKinney, J. C. 2011, *MNRAS*, 418, L79, doi: [10.1111/j.1745-3933.2011.01147.x](https://doi.org/10.1111/j.1745-3933.2011.01147.x)
- Thorne, K. S. 1974, *Astrophysical Journal*, 191, 507, doi: [10.1086/152991](https://doi.org/10.1086/152991)
- Wakabayashi, J. 2025, *Self-Sustained Penrose Excitation of Accretion Disks*, OSF Registration (Version 1.1) doi: [10.17605/osf.io/Q7DTH](https://doi.org/10.17605/osf.io/Q7DTH)

Young, S., Axon, D., Robinson, A., Smith, J., & Hough, J. H. 2011, in ASP Conf. Ser., Vol. 449, Astronomical Polarimetry 2008: Science from Small to Large Telescopes, 436. <https://ui.adsabs.harvard.edu/abs/2011ASPC..449..436Y/abstract>