

Self-Sustained Penrose Excitation

A Spin-Regulated Mechanism for Super-Eddington Quasar Luminosities

1. The Puzzle: Super-Eddington Quasars & Spin Ceiling

Quasars often shine brighter than the Eddington limit (L_{Edd}), sometimes by factors of 2-10x. Standard models struggle to explain this. Furthermore, observations suggest SMBH spins (a_*) rarely exceed unity, hinting at a self-regulating mechanism.

2. Core Mechanism: Equatorial Penrose Excitation

Near extremal spin ($a_* \rightarrow 1$), an equatorial return current sheet forms at the ISCO-ergoregion interface. Magnetic reconnection ejects plasmoids. Within the ergoregion, these plasmoids undergo a **Penrose split**:


- A **negative-energy branch** ($E - \Omega_H L < 0$) falls into the black hole.
- A **positive-energy branch** escapes as a narrow equatorial quasi-beam.

A fraction (ϵ_{coup}) of this beam re-enters the inner disk, boosting luminosity and extracting angular momentum, which enforces the spin ceiling.

3. Simulator: Spin & Luminosity

The plot shows the total luminosity ($L_{\text{tot}} = L_{\text{acc}} + L_{\text{self}}$) vs. spin (a_*), assuming a base accretion of $L_{\text{acc}} = 1 L_{\text{Edd}}$ and a fiducial coupling $\epsilon_{\text{coup}} = 0.01$.

Note the sharp, non-linear rise as a_* surpasses the threshold $a_{\text{th}} \approx 0.97$.



4. Key Observational Signatures (The 'Hooks')

Target: high spin ($a_* > a_{\text{th}}$), high λ , transparency ($\ell \lesssim 30$)

- **Hook (Support):** energy-dependent Fe K α /UV lags; spin-dependent polarization; EUV/MeV shoulder