

The Spin-Regulated Quasar

A New Mechanism for Super-Eddington Luminosities

The Quasar Conundrum

Astrophysics faces two major questions regarding supermassive black holes: How do quasars get so bright, often exceeding the Eddington limit ($L > L_{\text{Edd}}$)? And why do observed black hole spins appear to "max out" just below the theoretical maximum ($a_* \approx 1$), even with continuous accretion?


The Luminosity Problem

$$L > L_{\text{Edd}}$$

Many observed quasars are brighter than the Eddington limit, which theoretically should blow away accreting gas.

The Spin Ceiling Problem

The Spin Ceiling Problem



Observed Max Spin ($a_* < 0.9985$)

Standard accretion theory predicts a spin limit of $a_* \approx 0.9985$ but fails to explain why many are lower, suggesting a self-regulating mechanism to enforce a sub-unity ceiling.

A Proposed Solution: Self-Sustained Penrose Excitation

A new model suggests a "Self-Sustained Penrose Excitation" process. Active at high spin rates and high accretion rates, this mechanism operates at the boundary of the ergoregion (where spacetime is dragged) and the inner edge of the accretion disk (ISCO).

How It Works: A Spin-Regulated Feedback Loop
$P_{\text{ext}} = P_{\text{BZ}} + P_{\text{eq}}$
Total extracted power is the sum of equatorial (Eq) and polar (BZ) components.
$f_{\text{BZ}} + f_{\text{eq}} + f_{\text{res}} = 1$
The fractions must sum to unity.

This mechanism acts as a "lever" that taps into the black hole's rotational energy to power the disk's luminosity, creating a self-sustaining and self-limiting cycle.

1. High Accretion Gas flows in at super-Eddington rates.	2. Penrose Excitation At the ISCO/Ergoregion boundary, disk energy taps BH spin.	3. Sustained $L > L_{\text{Edd}}$ The extracted rotational energy powers the disk, exceeding the Eddington limit.
4. Spin Regulation (Feedback) The process extracts angular momentum, enforcing a spin ceiling, which in turn regulates the excitation process.		

Key Observational Signatures (The "Hooks")

This model is not just theoretical; it makes three specific, testable predictions ("hooks") that should co-occur in high-spin, high-accretion quasars.

