

# FPT Algorithms – Incremental Dominating Set Solutions

Kai Wang

Master by Research

School of Engineering and IT Faculty of EHSE

Charles Darwin University
Darwin

2014

## ACKNOWLEDGEMENTS

## **CONTENTS**

1	Intr	oductio	on	1
	1.1	Motiv	vation	1
	1.2	Contr	ibution Overview	1
	1.3	Thesis	s Overview	1
2	Prel	iminar	ies and Notation	3
	2.1	Set Th	neory	3
	2.2	Graph	Theory	3
		2.2.1	Dominating Set	3
	2.3	Comp	olexity Theory	4
		2.3.1	Decision Problems	4
		2.3.2	NP,NP-Hard, NP-complete	4
		2.3.3	Growth Rate of Function	4
		2.3.4	Fixed Parameter Tractability	4
		2.3.5	W-Hierarchy	4
		2.3.6	Kernelization	4

Contents

		2.3.7	DOMINATING SET	4
		2.3.8	Hamming Distance	5
		2.3.9	INCREEMENTAL DOMINATING SET	6
	2.4	Redu	ction Rules	7
3	Con	ıclusior	ı	9

# **ABSTRACT**

8 Contents

# List of Figures

10 List of Figures

# LIST OF TABLES

## Introduction

- 1.1 Motivation
- 1.2 Contribution Overview
- 1.3 Thesis Overview

2 1. Introduction

## PRELIMINARIES AND NOTATION

## 2.1 Set Theory

//TODO: add later  $\mathbb N$ 

- 2.2 Graph Theory
- 2.2.1 Dominating Set

//TODO:add later

- 2.3 Complexity Theory
- 2.3.1 Decision Problems
- 2.3.2 NP,NP-Hard, NP-complete

//TODO:add later

- 2.3.3 Growth Rate of Function
- 2.3.4 Fixed Parameter Tractability
- 2.3.5 W-Hierarchy

//TODO:add later

- 2.3.6 Kernelization
- 2.3.7 DOMINATING SET

Dominating set is one of natural properties of graphs, while DOMINATING SET problem is one of complex problems studied in complexity theory. DOMINATING SET problem is categorized in NP-complete class [GJ79].

## DOMINATING SET

Instance A graph G=(V,E) and  $k \in \mathbb{N}$ .

Question Is there a dominating set  $D \subseteq V$  for G such that  $|D| \leqslant k$ ?

[G]79]

**DOMINATING SET** has been proved to be a W[2] – complete problem by Downey and Fellows in 1995 [DF95]. In another words, this problem is not a FPT problem and does not have kernel. Nevertheless, the incremental edition of this problem, INCREEMENTAL DOMINATING SET problem, can be classified as a FPT problem [RGD14].

#### 2.3.8 Hamming Distance

Before talking about the INCREEMENTAL DOMINATING SET problem, we need clarify some concepts to help us understand the increment problem.

Assuming there are two vectors with the same length, we can check if the symbols in the corresponding positions of the two vectors are same or different. We call the number of positions where the symbols are different as  $Hamming\ Distance\ [Ham50]$ . Obviously, this technique can be applied in graphs. Firstly, given two graphs G=(V,E) and G'=(V,E'), both have the same set of vertices but different set of edges. We say that set E' is obtained by a series of  $edge\ edit\ operations$  from E, which refers to edge deletion and edge addition. Secondly, we can establish two 0/1 vectors to indicate E and E'. Thirdly, we can

find the Hamming distance between E and E', which is denoted by  $d_e(G,G')$ . We call  $d_e(G,G')$  as  $edge \ edit \ distance$ . In the fourth step, if there exists a solution of vertex set  $S \subset V$  for graph G and there may or may not exist another solution  $S' \subset V$  for G' with respect to a certain graph problem, we can also establish two 0/1 vectors to indicate S and S'. Finally, we can define the Hamming distance  $d_H(S,S')$  as the  $vertex \ solution \ set \ distance \ [RGD14].$ 

#### 2.3.9 INCREEMENTAL DOMINATING SET

With the assistance of the Hamming distance of  $d_e(G, G')$  and  $d_H(S, S')$ , we can define INCREEMENTAL PROBLEM.

#### INCREEMENTAL PROBLEM (INC-PROBLEM)

Instance A graph G = (V, E) and a set  $S \subseteq V$  where S has a certain property P for G,

A graph G' = (V, E') with  $d_e(G, G') \leq k$ ,

 $k, r \in \mathbb{N}$ 

Parameter(k, r)

Question Is there a set  $S' \subseteq V$  such that S' has property P for G' and

 $d_H(S,S') \leqslant r$ 

[RGD14]

With regards to the property of dominating set of graph, the definition of INCREEMENTAL DOMINATING SET problem can be presented in the following form:

2.4 Reduction Rules

7

#### INCREEMENTAL DOMINATING SET (INC-DS)

Instance A graph G=(V,E) and a dominating set  $S\subseteq V$  for G, A graph G'=(V,E') with  $d_e(G,G')\leqslant k$ ,  $k,r\in\mathbb{N}$ Parameter (k,r)Question Is there a dominating set  $S'\subseteq V$  such that  $d_H(S,S')\leqslant r$ [RGD14]

### 2.4 Reduction Rules

An improved FPT Algorithm for Vertex Cover

**Crown Reduction Rule** 

## Conclusion

10 3. Conclusion

## APPENDIX A

**Source Code** 

12 3. Conclusion

## **BIBLIOGRAPHY**

- [DF95] R. G. Downey and M. R. Fellows, *Parameterized computational feasibility*. Springer, Boston, 1995. 5
- [GJ79] M. Garey and D. Johnson, Computers and Intractability: A
  Guide to the Theory of NP-Completeness. W. H. Freeman, New
  York, 1979. 4, 5
- [Ham50] R. W. Hamming, "Error detecting and error correcting codes,"

  Bell System technical journal, vol. 29, no. 2, pp. 147–160, 1950. 5
- [RGD14] M. R. F. F. A. R. P. S. Rodney G. Downey, Judith Egan, "Incremental dominating set," processing, 2014. 5, 6, 7