

Simulation summary

Xuelong Wang

7/12/2018

Contents

1	Background of the environmental study	1
2	Estimation of the Cumulative (main) effect	1
2.1	GCTA method	1
2.2	Proposed Method (uncorrelating)	2
3	Estimation of the interactive effect	3
3.1	Model with interaction terms	3
3.2	Issues of estimating the interaction effect	3
3.3	Normality's affect on the performance of signal effect estimation	3
3.4	Questions	5
4	Further work	5

1 Background of the environmental study

The overall goal of this study is to find the relation between chemical exposures and health outcome. Due to the complexity of the problem, we have to tackle the problem step by step. More specifically, we want to find a model to estimate the **cumulative(main)**, **interactive(interaction)**, and **separate** effects.

2 Estimation of the Cumulative (main) effect

Since the magnitudes of the covariates from the environmental study (e.g. PCB data) is small, the signal of the environmental factors will probably be weak. This situation is very similar with what we got in the **GWAS** study. Therefore, it's natural to pick up the approaches used by GWAS studies. For example, the GCTA method.

2.1 GCTA method

2.1.1 Model

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_m x_{im} + \epsilon,$$

The matrix format is

$$Y = X\beta + \epsilon,$$

If x 's are standardized and independent from ϵ ,

$$\text{var}(y) = \text{var}(\sum \beta_i x_i) + \text{var}(\epsilon) = \sum \beta_i^2 + \sigma_\epsilon^2 = \sigma_\beta^2 + \sigma_\epsilon^2$$




Figure 1: Main effect estimation by proposed method box plot




Figure 2: Main effect estimation by proposed method hist

2.2 Proposed Method (uncorrelating)

To adopt GCTA approach, we need to transform the original data to uncorrelated data by using a linear transformation.

It seems that the correlation problem can be solved well based on the simulation's result. Although there are high correlation among the environmental data, the proposed method method can estimate the main effect unbiasedly.

2.2.1 Model

$$y_i = \alpha_0 + \alpha_1 z_{i1} + \cdots + \alpha_m z_{im} + \epsilon,$$

The matrix format is

$$Y = Z\alpha + \epsilon,$$

where $Z = XA^{-1}$ and $\alpha = A\alpha$,

Then we have

$$\text{var}(Z\alpha) = \text{var}(X\beta)$$

2.2.2 Simulation result

The full details can be found here

3 Estimation of the interactive effect

In this case, we're not only interested in the main effect but also in the interaction effect. It is possible that interaction effect also has a contribution to the response. Besides, under the environmental study, the total number of the covariates is much smaller than the GWAS study so that considering the the interaction is also feasible.

3.1 Model with interaction terms

$$y_i = \sum_{j=1}^m x_j * \beta_j^{(main)} + \sum_{j=1}^{m(m-1)/2} x_j^{(inter)} \gamma_j^{(inter)} + \epsilon_i$$

The variance of y could be decomposed as following:

$$var(y_i) = \sigma_\beta^2 + \sigma_\gamma^2 + \sigma_\epsilon^2$$

3.2 Issues of estimating the interaction effect

If consider to use the GCTA approach, there is a large bias on the estimation of the interaction effect. Based on the simulation results from the proposal, the marginal distribution of the covariates may have affected the proposed method's performance. It seems that it works well under the normal distribution even give the correlation structure between covariates.

3.3 Normality's affect on the performance of signal effect estimation

To check what influence of the normality will have on the performance, I conduct several simulations studying which includes tranforming and subsetting the covariates.

3.3.1 Transformation

Since the data is right skewed, I consider log and square root transformation. Besides, I also consider the rank and normal quantile transformation. Categorized transformation is also used to improve the normality

3.3.1.1 simulation result

Basically, all the transformation methods improve the performance. However, the bias issue is still not solved, especially for the proposed method.

Following is the result of rank and normal quantile transformation. For the graphs you can tell that the proposed method still has bias after the tranformation

3.3.2 Subset

To further improve the normality, I just remove several covariates which have a very un-symmetric empirical pdf (even after normal transformation). Therefore, after this step, most of the covariates should have a nice symmteric bell-shape distribution.




Figure 3: Rank Transformation result box plot




Figure 4: Rank Transformation result hist




Figure 5: Normal Quantile Transformation result box plot





Figure 6: Normal Quantile Transformation result hist

3.3.2.1 simulation result



3.4 Questions

1. Since we consider the interaction terms, the covariates cannot be independent any more
2. Interaction terms are also not standardized, which means that the $E(x^{inter}) \neq 0$
3. What's will be the sparsity of the interaction terms?

4 Further work