

CILE Clinical Consultation Report

Patient Vitals & Demographics

Age: 8 | Gender: M
Weight: 25.0 kg | SrCr: SrCr was not given mg/dL
Estimated CrCl: CrCl could not be calculated mL/min

Predictive Clinical Alerts & Profiles

Drug: Codeine

[Drug Classification]: NARCOTIC, PEDIATRIC
[RoA]: IV, Oral, SubQ
[PGx Markers Found] Targets: RECEPTOR, METABOLIZER, CYP2D6

- > [CLINICAL INTERCEPT] Ultra-rapid metabolizers (CYP2D6) risk fatal respiratory depression. Contraindicated in children <12 and post-tonsillectomy.
- > [FDA ALERT] WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF CODEINE SULFATE TABLETS Addiction, Abuse, and Misuse Because the use of Codeine Sulfate Tablets exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions
- > [FDA ALERT] Life-Threatening Respiratory Depression Serious, life-threatening, or fatal respiratory depression may occur with use of Codeine Sulfate Tablets, especially during initiation or following a dosage increase
- > [FDA ALERT] Accidental Ingestion Accidental ingestion of even one dose of Codeine Sulfate Tablets, especially by children, can result in a fatal overdose of codeine .