

CILE Clinical Consultation Report

Patient Vitals & Demographics

Age: 68 | Gender: M
Weight: 80.0 kg | SrCr: 1.2 mg/dL
Estimated CrCl: 66.67 mL/min

Predictive Clinical Alerts & Profiles

Drug: Amlodipine

[Drug Classification]: GERIATRIC
[RoA]: IV, Oral
[PGx Markers Found] Targets: CYP3A5, RECEPTOR

- > [CLINICAL INTERCEPT] Calcium channel blocker. High risk of peripheral edema and orthostatic hypotension in older adults.
- > [FDA ALERT] In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Drug: Metformin

[Drug Classification]: RENAL
[RoA]: IV, Oral, SubQ
[PGx Markers Found] Targets: CYP2C9, CYP3A4, CYP2C8

- > [CLINICAL INTERCEPT] Monitor eGFR; contraindicated if eGFR <30 mL/min due to risk of lactic acidosis; do not initiate if eGFR 30-45.
- > [FDA ALERT] 8.5 Geriatric Use ZITUVIMET In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy and the higher risk of lactic acidosis
- > [FDA ALERT] Renal function should be assessed more frequently in elderly patients.