Session 7: Survival Analysis 2

Levi Waldron

CUNY SPH Biostatistics 2

Learning objectives and outline

Learning objectives

1. Define proportional hazards
2. Perform and interpret Cox proportional hazards regression
3. Define time-dependent covariates and their use
4. Identify the differences between parametric and semi-parametric survival models
5. Identify situations when a parametric survival model might be useful

Outline

1. Review of survival and hazard functions
2. The Cox proportional hazards model
   - interpretation and inference
   - what are proportional hazards
   - when hazards aren't proportional
3. Parametric vs semi-parametric survival models

- Vittinghoff sections 6.1-6.2, 6.4

Review of survival and hazard functions

Recall leukemia Example

- Study of 6-mercaptopurine (6-MP) maintenance therapy for children in remission from acute lymphoblastic leukemia (ALL)
- 42 patients achieved remission from induction therapy and were then randomized in equal numbers to 6-MP or placebo.
- Survival time studied was from randomization until relapse.

Leukemia follow-up table


Table 3.13 Follow-up table for placebo patients in the leukemia study

Week of follow-up	No. followed	No. relapsed	No. censored	Conditional prob. of remission	Survival function
1	21	2	0	$19/21 = 0.91$	0.91
2	19	2	0	$17/19 = 0.90$	$0.90 \times 0.91 = 0.81$
3	17	1	0	$16/17 = 0.94$	$0.94 \times 0.81 = 0.76$
4	16	2	0	$14/16 = 0.88$	$0.88 \times 0.76 = 0.67$
5	14	2	0	$12/14 = 0.86$	$0.86 \times 0.67 = 0.57$
6	12	0	0	$12/12 = 1.00$	$1.00 \times 0.57 = 0.57$
7	12	0	0	$12/12 = 1.00$	$1.00 \times 0.57 = 0.57$
8	12	4	0	$8/12 = 0.67$	$0.67 \times 0.57 = 0.38$
9	8	0	0	$8/8 = 1.00$	$1.00 \times 0.38 = 0.38$
10	8	0	0	$8/8 = 1.00$	$1.00 \times 0.38 = 0.38$

Figure 1: leukemia Follow-up Table

This is the **Kaplan-Meier Estimate** $\hat{S}(t)$ of the Survival function $S(t)$.

Leukemia Kaplan-Meier plot




The hazard function h(t)

- *Definition*: The *survival function* at time t, denoted $S(t)$, is the probability of being event-free at t. Equivalently, it is the probability that the survival time is greater than t.
- *Definition*: The *cumulative event function* at time t, denoted $F(t)$, is the probability that the event has occurred by time t, or equivalently, the probability that the survival time is less than or equal to t. $F(t) = 1 - S(t)$.
- *Definition*: The *hazard function* $h(t)$ is the short-term event rate for subjects who have not yet experienced an event.
  - $h(t)$ is the probability of an event in the time interval $[t, t + s]$ (s is small), given that the individual has survived up to time t

$$h(t) = \lim_{s \to 0} \frac{Pr(t \leq T < t + s | T \geq t)}{s}$$


Leukemia dataset $S(t)$

Survival function $S(t)$



Leukemia dataset $h(t)$

Hazard function $h(t)$



SAS and R Source code

The Hazard Ratio (HR)

- If we are comparing the hazards of a control and a treatment group, it could in general be a function of time:
  - $HR(t) = h_T(t)/h_C(t)$
- Interpretation: the risk of event for the treatment group compared to the control group, as a function of time

The Proportional Hazards Assumption

- *Definition*: Under the *proportional hazards assumption*, the hazard ratio does not vary with time. That is, $HR(t) \equiv HR$.
- In other words, HR does not vary with time
  - $HR(t)$ is a constant, HR, at *all times* t
  - this assumption is about the population, of course there will be sampling variation

A nice proportional hazards dataset






Fig. 6.3 Hazard functions for 6-, 11-, and 21-year-old transplant recipients


Figure 2: Vittinghoff Figure 6.3, p. 210

The hazard function h(t) (leukemia dataset)





Log-minus-log plot



Recall previous regression models

$$E[y_i|x_i] = \beta_1 x_{1i} + \beta_2 x_{2i} + ... + \beta_p x_{pi}$$

- x_p are the predictors or independent variables
- y is the outcome, response, or dependent variable
- $E[y|x]$ is the expected value of y given x
- β_p are the regression coefficients

For logistic regression:

$$Logit(P(x_i)) = log\left(\frac{P(x_i)}{1 - P(x_i)}\right) = \beta_1 x_{1i} + \beta_2 x_{2i} + ... + \beta_p x_{pi}$$

For log-linear regression:

$$log(E[y_i|x_i]) = \beta_1 x_{1i} + \beta_2 x_{2i} + ... + \beta_p x_{pi}$$

Cox proportional hazards model

Cox proportional hazards model

- Cox proportional hazards regression assesses relationship between a right-censored, time-to-event outcome and predictors:
  - categorical variables (e.g., treatment groups)
  - continuous variables

$$log(HR(x_i)) = log\frac{h(t|x_i)}{h_0(t)} = \beta_1 x_{1i} + \beta_2 x_{2i} + ... + \beta_p x_{pi}$$

- $HR(x_i)$ is the hazard of patient i relative to baseline
- $h(t|x_i)$ is the time-dependent hazard function $h(t)$ for patient i
- $h_0(t)$ is the *baseline hazard function*

Multiplicative or additive model?

Interpretation of coefficients

- Coefficients β for a categorical / binary predictor:
  - β is the *log* of the ratio of hazards for the comparison group relative to reference group ($log(HR)$)
- Coefficients β for a continuous predictor:
  - β is the *log* of the ratio of hazards for someone having a one unit higher value of x (1 year, 1mm Hg, etc)
- If the hazard ratio ($exp(\beta)$) is close to 1 then the predictor does not affect survival
- If the hazard ratio is less than 1 then the predictor is protective (associated with improved survival)
- If the hazard ratio is greater than 1 then the predictor is associated with increased risk (= decreased survival)

Hypothesis testing and CIs

- Wald Test or Likelihood Ratio Test for coefficients
  - $H_0 : \beta = 0, H_a : \beta \neq 0$
  - equivalent to $H_0 : HR = 1, H_a : HR \neq 1$
- CIs typically obtained from Wald Test, reported for HR

Cox PH regression for Leukemia dataset

```
## Call:
## coxph(formula = Surv(time, cens) ~ group, data = leuk)
##
##   n= 42, number of events= 30
##
##              coef exp(coef) se(coef)      z Pr(>|z|)
## group6 MP -1.5721    0.2076   0.4124 -3.812 0.000138 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
##           exp(coef) exp(-coef) lower .95 upper .95
## group6 MP    0.2076      4.817   0.09251    0.4659
##
## Concordance= 0.69  (se = 0.041 )
## Likelihood ratio test= 16.35  on 1 df,   p=5e-05
## Wald test            = 14.53  on 1 df,   p=1e-04
## Score (logrank) test = 17.25  on 1 df,   p=3e-05
```

Cox PH is a semi-parametric model

- Cox proportional hazards model is *semi-parametric*:
  - assumes proportional hazards (PH), but no assumption on $h_0(t)$
  - robust if PH assumption is not violated
  - time-dependent covariates may resolve apparent violations of the PH assumption.

Summary: proportional hazards assumption

- Constant hazard *ratio* between groups over time (proportional hazards)
- A linear association between the natural log of the relative hazard and the predictors (log-linearity)
    - A multiplicative relationship between the predictors and the hazard
- Uninformative censoring

What to do when proportional hazards doesn't hold?

- **Time-dependent covariates**
- **Definition**: A time-dependent covariate is a predictor whose values may vary with time.
- **Basic rule**: You cannot look into the future in your analysis (even though it took place in the past) E.g.:
  - breast cancer chemotherapy patients divided into groups based on how much of the planned dose they received
  - patients divided into groups based on early response to treatment (shrinkage of tumor, lowering of cholesterol, etc)
  - interpolation of the values of a laboratory test linearly between observation times
  - removing subjects who do not finish the treatment plan
  - imputing the date of an adverse event as midway between observation times

Source: Using Time Dependent Covariates and Time Dependent Coefficients in the Cox Model

Immortal time bias example

- Immortal time bias is an example of looking into the future.
- E.g. Yee *et al.* reported that new statin users reported a 26% reduction in the risk of diabetes progression with one year or more of treatment relative to never-users (adjusted HR 0.74, 95% CI: 0.56 to 0.97).
  - New users excludes those who had received a lipid lowering drug from three years before to six months after cohort entry
- HR>1 was expected:people whose diabetes progresses are more likely to develop cardiovascular disease, an indication for statins.
- This is a result of an analysis error. Why?

- Yee *et al.* Statin use in type 2 diabetes mellitus is associated with a delay in starting insulin (http://onlinelibrary.wiley.com/doi/10.1111/j.1464-5491.2004.01263.x/full)

Immortal time bias example (cont'd)

- What was the analysis error?
  - "new statin user" group was defined based on future initiation: knowledge unknown at the time of entry into the study
  - guaranteed no events for statin users from cohort entry to start of statin use
  - thus all persons in the *treated* group are "immortal" from time 0 until the initiation of statin treatment
  - this period of immortality made treatment look more effective

Parametric survival models

What are "parametric" survival models?

- "Parametric" models estimate additional *parameters* for the baseline hazard, e.g.:
  - **Weibull**: hazard function is a polynomial
  - **exponential**: hazard function is constant over time, survival function is exponential (special case of Weibull):
    e.g. healthy population with randomly occurring events
  - many other options for assumption of distributions
- In most common implementation a log-transform of the time variable is used
  - then can be interpreted as *Accelerated Failure Time* (AFT) models.

Coefficients in parametric models

- The interpretation of β coefficients is different:
  - Cox model: $log(HR)$
  - AFT models: $log(survivaltimeratio)$
  - The sign is *opposite* (*i.e.* if one is positive the other is negative)

Why use a parametric survival model?

- Can be more powerful if assumption is correct
  - may help with small numbers of events
- Extra capabilities:
  - smooth estimation of baseline hazard
  - extrapolation
  - complicated censoring
- Easy to interpret: coefficients are $log(survivaltimeratio)$
- Easy to fit: replace `survival::coxph` with `survival::survreg`

Why not to use a parametric survival model?

- Depend on correct specification of baseline hazard model
- Even if correctly specified, may not provide much improvement in efficiency
- Still make a proportionality assumption, on survival functions