

Session 4: loglinear regression part 1

Levi Waldron

CUNY SPH Biostatistics 2

Learning objectives and outline

Learning objectives

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

- 1 Define log-linear models in GLM framework
- 2 Identify situations that motivate use of log-linear models
- 3 Assess model fit of log-linear models
- 4 Define multi-collinearity

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

- 1 brief review of GLMs
- 2 Motivating example for log-linear models
 - Poisson regression
- 3 Checking model assumptions and fit: Residual Analysis
- 4 Note on collinearity

Reading: Vittinghoff textbook chapter 8.1-8.3

**Session 4:
loglinear
regression
part 1**

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

Review

Components of GLM

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

- **Random component** specifies the conditional distribution for the response variable - it doesn't have to be normal but can be any distribution that belongs to the "exponential" family of distributions
- **Systematic component** specifies linear function of predictors (linear predictor)
- **Link** [denoted by $g(\cdot)$] specifies the relationship between the expected value of the random component and the systematic component, can be linear or nonlinear

Linear Regression as GLM

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

- **The model:**

$$y_i = E[y|x] + \epsilon_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi} + \epsilon_i$$

- **Random component** of y_i is normally distributed:

$$\epsilon_i \stackrel{iid}{\sim} N(0, \sigma_\epsilon^2)$$

- **Systematic component** (linear predictor):

$$\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi}$$

- **Link function** here is the *identity link*:

$g(E(y|x)) = E(y|x)$. We are modeling the mean directly, no transformation.

Logistic Regression as GLM

- **The model:**

$$\text{Logit}(P(x)) = \log \left(\frac{P(x)}{1 - P(x)} \right) = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi}$$

- **Random component:** y_i follows a Binomial distribution (outcome is a binary variable)
- **Systematic component:** linear predictor

$$\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi}$$

- **Link function:** *logit* (Converts Prob \rightarrow log-odds)

$$g(P(x)) = \text{logit}(P(x)) = \log \left(\frac{P(x)}{1 - P(x)} \right)$$

$$P(x) = g^{-1}(\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi})$$

Additive vs. Multiplicative models

- Linear regression is an *additive* model
 - e.g. for two binary variables $\beta_1 = 1.5$, $\beta_2 = 1.5$.
 - If $x_1 = 1$ and $x_2 = 1$, this adds 3.0 to $E(y|x)$
- Logistic regression is a *multiplicative* model
 - If $x_1 = 1$ and $x_2 = 1$, this adds 3.0 to $\log(\frac{P}{1-P})$
 - Odds-ratio $\frac{P}{1-P}$ increases 20-fold: $\exp(1.5 + 1.5)$ or $\exp(1.5) * \exp(1.5)$

Motivating example for log-linear models

Effectiveness of a depression case-management program

- Research question: can a new treatment reduce the number of needed visits to the emergency room, compared to standard care?
- *outcome*: # of emergency room visits for each patient in the year following initial treatment
- *predictors*:
 - *race* (white or nonwhite)
 - *treatment* (treated or control)
 - *amount of alcohol consumption* (numerical measure)
 - *drug use* (numerical measure)

Statistical issues

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

- 1 about $1/3$ of observations are exactly 0 (did not return to the emergency room within the year)
- 2 highly nonnormal and cannot be transformed to be approximately normal
- 3 even $\log(y_i + 1)$ transformation will have a “lump” at zero + over $1/2$ the transformed data would have values of 0 or $\log(2)$
- 4 a linear regression model would give negative predictions for some covariate combinations
- 5 some subjects die or cannot be followed up on for a whole year

Towards a reasonable model

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

- A *multiplicative* model will allow us to make inference on *ratios* of mean emergency room usage
- Modeling *log* of the *mean* emergency usage ensures positive means, and does not suffer from $\log(0)$ problem
- Random component of GLM, or residuals (was $\epsilon_i \stackrel{iid}{\sim} N(0, \sigma_\epsilon^2)$ for linear regression) may still not be normal, but we can choose from other distributions

Proposed model without time

$$\log(E[Y_i]) = \beta_0 + \beta_1 \text{RACE}_i + \beta_2 \text{TRT}_i + \beta_3 \text{ALCH}_i + \beta_4 \text{DRUG}_i$$

Or equivalently:

$$E[Y_i] = \exp(\beta_0 + \beta_1 \text{RACE}_i + \beta_2 \text{TRT}_i + \beta_3 \text{ALCH}_i + \beta_4 \text{DRUG}_i)$$

where $E[Y_i]$ is the expected number of emergency room visits for patient i .

- Important note: Modeling $\log(E[Y_i])$ is *not* equivalent to modeling $E(\log(Y_i))$

Accounting for follow-up time

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

Instead, model mean count per unit time:

$$\log(E[Y_i]/t_i) = \beta_0 + \beta_1 \text{RACE}_i + \beta_2 \text{TRT}_i + \beta_3 \text{ALCH}_i + \beta_4 \text{DRUG}_i$$


Or equivalently:

$$\log(E[Y_i]) = \beta_0 + \beta_1 \text{RACE}_i + \beta_2 \text{TRT}_i + \beta_3 \text{ALCH}_i + \beta_4 \text{DRUG}_i + \log(t_i)$$

- $\log(t_i)$ is not a covariate, it is called an *offset*

The Poisson distribution

- Count data are often modeled as Poisson distributed: + mean λ is greater than 0 + variance is also λ + Probability density $P(k, \lambda) = \frac{\lambda^k}{k!} e^{-\lambda}$



When the Poisson distribution works

- Individual events are low-probability (small p), but many opportunities (large n)
 - e.g. # 911 calls per day
 - e.g. # emergency room visits
- Approximates the binomial distribution when n is large and p is small
 - e.g. $n > 20$, $np < 5$ or $n(1 - p) < 5$
- When mean of residuals is approx. equal to variance

GLM with log-linear link and Poisson error model

- Model the number of counts per unit time as Poisson-distributed + so the expected number of counts per time is λ_i

$$E[Y_i]/t_i = \lambda_i$$

$$\log(E[Y_i]/t_i) = \log(\lambda_i)$$

$$\log(E[Y_i]) = \log(\lambda_i) + \log(t_i)$$

Recalling the log-linear model systematic component:

$$\log(E[Y_i]) = \beta_0 + \beta_1 \text{RACE}_i + \beta_2 \text{TRT}_i + \beta_3 \text{ALCH}_i + \beta_4 \text{DRUG}_i + \log(t_i)$$

GLM with log-linear link and Poisson error model (cont'd)

Then the systematic part of the GLM is:

$$\log(\lambda_i) = \beta_0 + \beta_1 \text{RACE}_i + \beta_2 \text{TRT}_i + \beta_3 \text{ALCH}_i + \beta_4 \text{DRUG}_i$$

Or alternatively:

$$\lambda_i = \exp(\beta_0 + \beta_1 \text{RACE}_i + \beta_2 \text{TRT}_i + \beta_3 \text{ALCH}_i + \beta_4 \text{DRUG}_i)$$

Interpretation of coefficients

- Suppose that $\hat{\beta}_1 = -0.5$ in the fitted model, where $RACE_i = 0$ for white and $RACE_i = 1$ for non-white.
- The mean rate of emergency room visits per unit time for white relative to non-white, all else held equal, is estimated to be:

$$\begin{aligned} & \frac{\exp(\beta_0 + 0 + \beta_2 TRT_i + \beta_3 ALCH_i + \beta_4 DRUG_i)}{\exp(\beta_0 - 0.5 + \beta_2 TRT_i + \beta_3 ALCH_i + \beta_4 DRUG_i)} \\ &= \frac{e^{\beta_0} e^0 e^{\beta_2 TRT_i} e^{\beta_3 ALCH_i} e^{\beta_4 DRUG_i}}{e^{\beta_0} e^{-0.5} e^{\beta_2 TRT_i} e^{\beta_3 ALCH_i} e^{\beta_4 DRUG_i}} \\ &= \frac{e^0}{e^{-0.5}} \\ &= e^{0.5} \approx 1.65 \end{aligned}$$

Interpretation of coefficients (cont'd)

- If $\hat{\beta}_1 = -0.5$ with whites as the reference group: - after adjustment for treatment group, alcohol and drug usage, whites tend to use the emergency room at a rate 1.65 times higher than non-whites. - equivalently, the average rate of usage for whites is 65% higher than that for non-whites
- Multiplicative rules apply for other coefficients as well, because they are exponentiated to estimate the mean rate.

Multi-collinearity

Note on Multicollinearity

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

- 1 *Multicollinearity* exists when two or more of the independent variables in regression are moderately or highly correlated.
- 2 Multicollinearity implies near-linear relationship among the predictors
- 3 The presence of near-linear dependence dramatically impacts the ability to estimate regression coefficients
- 4 High multicollinearity results in larger standard errors for regression coefficients
 - estimates of such regression coefficients will tend to be less stable over repeated sampling

**Session 4:
loglinear
regression
part 1**

Levi Waldron

Learning
objectives and
outline

Review

Motivating
example for
log-linear
models

Multi-
collinearity

Conclusions

Conclusions

Conclusions

- ① Log-linear models are appropriate for non-negative, skewed count data
 - probability of each event is low
- ② The coefficients of log-linear models are *multiplicative*
- ③ An *offset* term can account for varying follow-up time or otherwise varying opportunity to be counted
- ④ Poisson distribution is high number of trials, low probability limit of binomial distribution
- ⑤ Inference from log-linear models is sensitive to the choice of error model (assumption on the distribution of residuals)
- ⑥ We will cover other options next week for when the Poisson error model doesn't fit:
 - Variance proportional to mean, instead of equal
 - Negative Binomial
 - Zero Inflation