

Session 7: Survival Analysis 2

Levi Waldron

CUNY SPH Biostatistics 2

Levi Waldron

**Learning
objectives and
outline**

Review of
 $S(t)$ and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

Learning objectives and outline

Learning objectives

- 1 Define the hazard function and proportional hazards
- 2 Define proportional hazards
- 3 Perform and interpret Cox proportional hazards regression

Outline

- ① Review of Survival function and Kaplan-Meier estimator
 - ② Hazard function
 - ③ Cox regression
 - the Cox proportional hazards model
 - proportional hazards
 - interpretation and inference from the Cox model
- Vittinghoff sections 6.1-6.2, 6.4

Review of $S(t)$ and KM estimator

Recall leukemia Example

Levi Waldron

Learning
objectives and
outline

Review of
 $S(t)$ and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

- Study of 6-mercaptopurine (6-MP) maintenance therapy for children in remission from acute lymphoblastic leukemia (ALL)
- 42 patients achieved remission from induction therapy and were then randomized in equal numbers to 6-MP or placebo.
- Survival time studied was from randomization until relapse.

Leukemia follow-up table

Table 3.13 Follow-up table for placebo patients in the leukemia study


Week of follow-up	No. followed	No. relapsed	No. censored	Conditional prob. of remission	Survival function
1	21	2	0	$19/21 = 0.91$	0.91
2	19	2	0	$17/19 = 0.90$	$0.90 \times 0.91 = 0.81$
3	17	1	0	$16/17 = 0.94$	$0.94 \times 0.81 = 0.76$
4	16	2	0	$14/16 = 0.88$	$0.88 \times 0.76 = 0.67$
5	14	2	0	$12/14 = 0.86$	$0.86 \times 0.67 = 0.57$
6	12	0	0	$12/12 = 1.00$	$1.00 \times 0.57 = 0.57$
7	12	0	0	$12/12 = 1.00$	$1.00 \times 0.57 = 0.57$
8	12	4	0	$8/12 = 0.67$	$0.67 \times 0.57 = 0.38$
9	8	0	0	$8/8 = 1.00$	$1.00 \times 0.38 = 0.38$
10	8	0	0	$8/8 = 1.00$	$1.00 \times 0.38 = 0.38$

Figure 1: leukemia Follow-up Table

This is the **Kaplan-Meier Estimate** $\hat{S}(t)$ of the Survival function $S(t)$.

Leukemia Kaplan-Meier plot

Warning: Vectorized input to 'element_text()' is n
Results may be unexpected or may change in future



The Hazard Function


The hazard function $h(t)$

- *Definition:* (review) The *survival function* at time t , denoted $S(t)$, is the probability of being event-free at t . Equivalently, it is the probability that the survival time is greater than t .
- *Definition:* (review) The *cumulative event function* at time t , denoted $F(t)$, is the probability that the event has occurred by time t , or equivalently, the probability that the survival time is less than or equal to t . $F(t) = 1 - S(t)$.
- *Definition:* The *hazard function* $h(t)$ is the short-term event rate for subjects who have not yet experienced an event.
 - $h(t)$ is the probability of an event in the time interval $[t, t + s]$ (s is small), given that the individual has survived up to time t

$$h(t) = \lim_{s \rightarrow 0} \frac{\Pr(t \leq T < t + s | T \geq t)}{s}$$

Leukemia dataset $h(t)$

Strata + group=Placebo + group=6 MP



The Hazard Ratio (HR)

Levi Waldron

Learning
objectives and
outline

Review of
 $S(t)$ and KM
estimator

**The Hazard
Function**

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

- If we are comparing the hazards of a control and a treatment group, it could in general be a function of time:
 - $HR(t) = h_T(t)/h_C(t)$
- Interpretation: the risk of event for the treatment group compared to the control group, as a function of time

The Proportional Hazards Assumption

- *Definition:* Under the *proportional hazards assumption*, the hazard ratio does not vary with time. That is, $HR(t) \equiv HR$.
- In other words, HR does not vary with time
 - $HR(t)$ is a constant, HR , at *all times* t
 - this assumption is about the population, of course there will be sampling variation

A nice proportional hazards dataset






Fig. 6.3 Hazard functions for 6-, 11-, and 21-year-old transplant recipients

The hazard function $h(t)$ (leukemia dataset)



'summarise()' ungrouping output (override with '.g

Log-minus-log plot



Recall previous regression models

Levi Waldron

Learning
objectives and
outline

Review of
S(t) and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

$$E[y_i|x_i] = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi}$$

- x_p are the predictors or independent variables
- y is the outcome, response, or dependent variable
- $E[y|x]$ is the expected value of y given x
- β_p are the regression coefficients

For logistic regression:

$$\text{Logit}(P(x_i)) = \log\left(\frac{P(x_i)}{1 - P(x_i)}\right) = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi}$$

For log-linear regression:

$$\log(E[y_i|x_i]) = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi}$$

Cox proportional hazards model

Cox proportional hazards model

Levi Waldron

Learning
objectives and
outline

Review of
 $S(t)$ and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

- Cox proportional hazards regression assesses relationship between a right-censored, time-to-event outcome and predictors:
 - categorical variables (e.g., treatment groups)
 - continuous variables

$$\log(HR(x_i)) = \log \frac{h(t|x_i)}{h_0(t)} = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi}$$

- $HR(x_i)$ is the hazard of patient i relative to baseline
- $h(t|x_i)$ is the time-dependent hazard function $h(t)$ for patient i
- $h_0(t)$ is the *baseline hazard function*

Multiplicative or additive model?

Interpretation of coefficients

Levi Waldron

Learning
objectives and
outline

Review of
 $S(t)$ and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

- Coefficients β for a categorical / binary predictor:
 - β is the *log* of the ratio of hazards for the comparison group relative to reference group ($\log(HR)$)
- Coefficients β for a continuous predictor:
 - β is the *log* of the ratio of hazards for someone having a one unit higher value of x (1 year, 1mm Hg, etc)
- If the hazard ratio ($\exp(\beta)$) is close to 1 then the predictor does not affect survival
- If the hazard ratio is less than 1 then the predictor is protective (associated with improved survival)
- If the hazard ratio is greater than 1 then the predictor is associated with increased risk (= decreased survival)

Hypothesis testing and CIs

Levi Waldron

Learning
objectives and
outline

Review of
 $S(t)$ and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

- Wald Test or Likelihood Ratio Test for coefficients
 - $H_0 : \beta = 0, H_a : \beta \neq 0$
 - equivalent to $H_0 : HR = 1, H_a : HR \neq 1$
- CIs typically obtained from Wald Test, reported for HR

CoxPH regression for Leukemia dataset

Levi Waldron

Learning
objectives and
outline

Review of
S(t) and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

```
## Call:
## coxph(formula = Surv(time, cens) ~ group, data = leuk)
##
##      n= 42, number of events= 30
##
##              coef exp(coef) se(coef)      z Pr(>|z|)
## group6 MP -1.5721    0.2076   0.4124 -3.812 0.000138 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
##              exp(coef) exp(-coef) lower .95 upper .95
## group6 MP    0.2076      4.817   0.09251   0.4659
##
## Concordance= 0.69 (se = 0.041 )
## Likelihood ratio test= 16.35  on 1 df,  p=5e-05
## Wald test               = 14.53  on 1 df,  p=1e-04
## Score (logrank) test = 17.25  on 1 df,  p=3e-05
```

Semi-parametric model

Levi Waldron

Learning
objectives and
outline

Review of
 $S(t)$ and KM
estimator

The Hazard
Function

Cox
proportional
hazards model

Parametric
models

Summary:
proportional
hazards
assumption

- Cox proportional hazards model is *semi-parametric*:
 - assumes proportional hazards (PH), but no assumption on $h_0(t)$
 - robust if PH assumption is not violated
 - time-dependent covariates may resolve apparent violations of the PH assumption.

Parametric models

What are “parametric” models?

- “Parametric” models estimate additional *parameters* for the baseline hazard, e.g.:
 - **Weibull**: hazard function is a polynomial
 - **exponential**: hazard function is constant over time, survival function is exponential (special case of Weibull): e.g. healthy population with randomly occurring events
 - many other options for assumption of distributions
- Can be more powerful if assumption is correct
- Allows smooth estimation of baseline hazard, extrapolation
- Still a proportional hazards model
- Fit the same way but using `survival::survreg`

Summary: proportional hazards assumption

Summary: proportional hazards assumption

- Constant hazard *ratio* between groups over time (proportional hazards)
- A linear association between the natural log of the relative hazard and the predictors (log-linearity)
 - A multiplicative relationship between the predictors and the hazard
- Uninformative censoring