

CSER 2207: Numerical Analysis

Lecture-9

Interpolation and Polynomial Approximation

Dr. Mostak Ahmed
Associate Professor
Department of Mathematics, JnU

Lagrange Interpolating Polynomials

The problem of determining a polynomial of degree one that passes through the distinct points (x_0, y_0) and (x_1, y_1) is the same as approximating a function f for which $f(x_0) = y_0$ and $f(x_1) = y_1$ by means of a first-degree polynomial **interpolating**, or agreeing with, the values of f at the given points. Using this polynomial for approximation within the interval given by the endpoints is called **polynomial interpolation**.

Define the functions

$$L_0(x) = \frac{x - x_1}{x_0 - x_1} \quad \text{and} \quad L_1(x) = \frac{x - x_0}{x_1 - x_0}.$$

The linear **Lagrange interpolating polynomial** through (x_0, y_0) and (x_1, y_1) is

$$P(x) = L_0(x)f(x_0) + L_1(x)f(x_1) = \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1).$$

Note that

$$L_0(x_0) = 1, \quad L_0(x_1) = 0, \quad L_1(x_0) = 0, \quad \text{and} \quad L_1(x_1) = 1,$$

which implies that

$$P(x_0) = 1 \cdot f(x_0) + 0 \cdot f(x_1) = f(x_0) = y_0$$

and

$$P(x_1) = 0 \cdot f(x_0) + 1 \cdot f(x_1) = f(x_1) = y_1.$$

So P is the unique polynomial of degree at most one that passes through (x_0, y_0) and (x_1, y_1) .

Example

Example 1 Determine the linear Lagrange interpolating polynomial that passes through the points (2, 4) and (5, 1).

Solution In this case we have


$$L_0(x) = \frac{x-5}{2-5} = -\frac{1}{3}(x-5) \quad \text{and} \quad L_1(x) = \frac{x-2}{5-2} = \frac{1}{3}(x-2),$$

so

$$P(x) = -\frac{1}{3}(x-5) \cdot 4 + \frac{1}{3}(x-2) \cdot 1 = -\frac{4}{3}x + \frac{20}{3} + \frac{1}{3}x - \frac{2}{3} = -x + 6.$$

The graph of $y = P(x)$ is shown in Figure 3.3. ■

Figure 3.3




Generalized Form

To generalize the concept of linear interpolation, consider the construction of a polynomial of degree at most n that passes through the $n + 1$ points

$$(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n)).$$

(See Figure 3.4.)

Figure 3.4



Cont...

In this case we first construct, for each $k = 0, 1, \dots, n$, a function $L_{n,k}(x)$ with the property that $L_{n,k}(x_i) = 0$ when $i \neq k$ and $L_{n,k}(x_k) = 1$. To satisfy $L_{n,k}(x_i) = 0$ for each $i \neq k$ requires that the numerator of $L_{n,k}(x)$ contain the term

$$(x - x_0)(x - x_1) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n).$$


To satisfy $L_{n,k}(x_k) = 1$, the denominator of $L_{n,k}(x)$ must be this same term but evaluated at $x = x_k$. Thus

$$L_{n,k}(x) = \frac{(x - x_0) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n)}{(x_k - x_0) \cdots (x_k - x_{k-1})(x_k - x_{k+1}) \cdots (x_k - x_n)}.$$

A sketch of the graph of a typical $L_{n,k}$ (when n is even) is shown in Figure 3.5.

Cont...

Figure 3.5



The interpolating polynomial is easily described once the form of $L_{n,k}$ is known. This polynomial, called the ***n*th Lagrange interpolating polynomial**, is defined in the following theorem.

Theorem 3.2

If x_0, x_1, \dots, x_n are $n + 1$ distinct numbers and f is a function whose values are given at these numbers, then a unique polynomial $P(x)$ of degree at most n exists with

$$f(x_k) = P(x_k), \quad \text{for each } k = 0, 1, \dots, n.$$

This polynomial is given by

$$P(x) = f(x_0)L_{n,0}(x) + \dots + f(x_n)L_{n,n}(x) = \sum_{k=0}^n f(x_k)L_{n,k}(x), \quad (3.1)$$

where, for each $k = 0, 1, \dots, n$,

$$\begin{aligned} L_{n,k}(x) &= \frac{(x - x_0)(x - x_1) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n)}{(x_k - x_0)(x_k - x_1) \cdots (x_k - x_{k-1})(x_k - x_{k+1}) \cdots (x_k - x_n)} \\ &= \prod_{\substack{i=0 \\ i \neq k}}^n \frac{(x - x_i)}{(x_k - x_i)}. \end{aligned} \quad (3.2)$$

We will write $L_{n,k}(x)$ simply as $L_k(x)$ when there is no confusion as to its degree.

Example 2

- (a) Use the numbers (called *nodes*) $x_0 = 2$, $x_1 = 2.75$, and $x_2 = 4$ to find the second Lagrange interpolating polynomial for $f(x) = 1/x$.
- (b) Use this polynomial to approximate $f(3) = 1/3$.

Solution (a) We first determine the coefficient polynomials $L_0(x)$, $L_1(x)$, and $L_2(x)$. In nested form they are

$$L_0(x) = \frac{(x - 2.75)(x - 4)}{(2 - 2.5)(2 - 4)} = \frac{2}{3}(x - 2.75)(x - 4),$$

$$L_1(x) = \frac{(x - 2)(x - 4)}{(2.75 - 2)(2.75 - 4)} = -\frac{16}{15}(x - 2)(x - 4),$$

and

$$L_2(x) = \frac{(x - 2)(x - 2.75)}{(4 - 2)(4 - 2.5)} = \frac{2}{5}(x - 2)(x - 2.75).$$

Cont...

Also, $f(x_0) = f(2) = 1/2$, $f(x_1) = f(2.75) = 4/11$, and $f(x_2) = f(4) = 1/4$, so

$$\begin{aligned} P(x) &= \sum_{k=0}^2 f(x_k)L_k(x) \\ &= \frac{1}{3}(x - 2.75)(x - 4) - \frac{64}{165}(x - 2)(x - 4) + \frac{1}{10}(x - 2)(x - 2.75) \\ &= \frac{1}{22}x^2 - \frac{35}{88}x + \frac{49}{44}. \end{aligned}$$

(b) An approximation to $f(3) = 1/3$ (see Figure 3.6) is

$$f(3) \approx P(3) = \frac{9}{22} - \frac{105}{88} + \frac{49}{44} = \frac{29}{88} \approx 0.32955.$$

Recall that in the opening section of this chapter (see Table 3.1) we found that no Taylor polynomial expanded about $x_0 = 1$ could be used to reasonably approximate $f(x) = 1/x$ at $x = 3$. ■

More Examples

Example 3.13 Certain corresponding values of x and $\log_{10} x$ are (300, 2.4771), (304, 2.4829), (305, 2.4843) and (307, 2.4871). Find $\log_{10} 301$.

From formula (3.43), we obtain

$$\begin{aligned}\log_{10} 301 &= \frac{(-3)(-4)(-6)}{(-4)(-5)(-7)}(2.4771) + \frac{(1)(-4)(-6)}{(4)(-1)(-3)}(2.4829) \\ &\quad + \frac{(1)(-3)(-6)}{(5)(1)(-2)}(2.4843) + \frac{(1)(-3)(-4)}{(7)(3)(2)}(2.4871) \\ &= 1.2739 + 4.9658 - 4.4717 + 0.7106 \\ &= 2.4786.\end{aligned}$$

Example 3.14 If $y_1 = 4$, $y_3 = 12$, $y_4 = 19$ and $y_x = 7$, find x .

Using formula (3.44), we have

$$\begin{aligned}x &= \frac{(-5)(-12)}{(-8)(-15)}(1) + \frac{(3)(-12)}{(8)(-7)}(3) + \frac{(3)(-5)}{(15)(7)}(4) \\ &= \frac{1}{2} + \frac{27}{14} - \frac{4}{7} \\ &= 1.86.\end{aligned}$$

The actual value is 2.0 since the above values were obtained from the polynomial $y(x) = x^2 + 3$.

Example 3.15 Find the Lagrange interpolating polynomial of degree 2 approximating the function $y = \ln x$ defined by the following table of values. Hence determine the value of $\ln 2.7$.

x	$y = \ln x$
2	0.69315
2.5	0.91629
3.0	1.09861

We have

$$l_0(x) = \frac{(x - 2.5)(x - 3.0)}{(-0.5)(-1.0)} = 2x^2 - 11x + 15.$$

Similarly, we find

$$l_1(x) = -(4x^2 - 20x + 24) \quad \text{and} \quad l_2(x) = 2x^2 - 9x + 10.$$

Hence

$$\begin{aligned} L_2(x) &= (2x^2 - 11x + 15)(0.69315) - (4x^2 - 20x + 24)(0.91629) \\ &\quad + (2x^2 - 9x + 10)(1.09861) \\ &= -0.08164x^2 + 0.81366x - 0.60761, \end{aligned}$$

which is the required quadratic polynomial.

Putting $x = 2.7$, in the above polynomial, we obtain

$$\ln 2.7 \approx L_2(2.7) = -0.08164(2.7)^2 + 0.81366(2.7) - 0.60761 = 0.9941164.$$

Actual value of $\ln 2.7 = 0.9932518$, so that

$$|\text{Error}| = 0.0008646.$$

Last but not least

Example 3.16 The function $y = \sin x$ is tabulated below

x	$y = \sin x$
0	0
$\pi/4$	0.70711
$\pi/2$	1.0

Using Lagrange's interpolation formula, find the value of $\sin(\pi/6)$.

We have

$$\begin{aligned}\sin \frac{\pi}{6} &\approx \frac{(\pi/6 - 0)(\pi/6 - \pi/2)}{(\pi/4 - 0)(\pi/4 - \pi/2)}(0.70711) + \frac{(\pi/6 - 0)(\pi/6 - \pi/4)}{(\pi/2 - 0)(\pi/2 - \pi/4)}(1) \\ &= \frac{8}{9}(0.70711) - \frac{1}{9} \\ &= \frac{4.65688}{9} \\ &= 0.51743.\end{aligned}$$

Thank You