LAKESIDE FAMILY MEDICINE

4400 Lakeshore Blvd, Suite 210, Chicago, IL 60640 | Tel: (773) 555-0344

OFFICE VISIT — SOAP NOTE

Patient:	Maria Elena Vasquez	**DOB:**	07/22/1972
SSN:	318-55-2241	**MRN:**	LFM-20891
Date of Visit:	11/03/2023	**Provider:**	Dr. James Whitfield, MD
Address:	1129 N. Maple Ave, Apt 3B, Chicago, IL 60614	**Phone:**	(773) 555-8821
Email:	mvasquez1972@gmail.com	**Insurance:**	United Healthcare #UH-882341

S — SUBJECTIVE

Ms. Vasquez is a 51-year-old Hispanic female presenting for her 3-month diabetes follow-up. She reports increased fatigue over the past 4 weeks. She has been checking her blood glucose at home — fasting readings typically 145–175 mg/dL. She notes occasional blurred vision in the evenings. She denies polyuria, polydipsia, or lower extremity numbness. Dietary adherence has been difficult due to a new work schedule (night shifts at Cook County Hospital where she works as a nursing aide). She is taking all medications as prescribed but occasionally misses her evening metformin dose.

O — OBJECTIVE

Vitals: BP 138/86 | HR 78 | Wt 172 lbs (↑4 lbs since last visit) | HbA1c today: 8.4% (prev 7.9%)

General: Alert, oriented, in no acute distress. Appears fatigued.

Eyes: PERRL. Fundoscopic deferred — refer ophthalmology.

Extremities: Monofilament sensation intact bilateral feet. No ulcers. Pedal pulses 2+.

Labs (today): FBG 168 mg/dL | BMP: Na 138, K 4.1, Cr 0.9, eGFR 74 | UA: glucose 2+, no protein

A — ASSESSMENT

1. Type 2 Diabetes Mellitus — suboptimally controlled (HbA1c 8.4%, target <7.0%)
2. Hypertension — borderline controlled
3. Weight gain — likely related to night-shift schedule and dietary changes
4. Possible early diabetic retinopathy — visual symptoms require evaluation

P — PLAN

1. Increase Metformin to 1000 mg BID. Counsel on importance of evening dose consistency.
2. Add Empagliflozin 10 mg daily — addresses glucose control and weight.
3. Refer to ophthalmology (Dr. Li Jing, Northwestern) for diabetic eye exam.
4. Dietitian referral for night-shift nutrition counseling.
5. Increase lisinopril from 5 mg to 10 mg for BP management.
6. Return in 3 months. HbA1c, CMP, lipid panel at next visit.

Signed: Dr. James Whitfield, MD | NPI: 9876543210 | Date: 11/03/2023 10:15 AM