

# Decomposition processes

## acetylene and ammonia

W. Dal'Maz Silva

17th May 2016

# Table of Contents

- 1 Introduction
- 2 Residence time distribution
  - Background
  - Measurements
- 3 Acetylene decomposition
  - Atmospheric pressure
  - Pyrolysis mechanism
- 4 Ammonia decomposition
- 5 Ongoing studies
- 6 Conclusion

# Table of Contents

- 1 Introduction
- 2 Residence time distribution
  - Background
  - Measurements
- 3 Acetylene decomposition
  - Atmospheric pressure
  - Pyrolysis mechanism
- 4 Ammonia decomposition
- 5 Ongoing studies
- 6 Conclusion

# Introduction

## Pyrolysis

Thermochemical decomposition of organic material at high temperature in the absence of oxygen.

## Heterogeneous processes

Adsorption, desorption, decomposition and recombination processes of molecules at solid surfaces.

## Context

Decomposition of precursors –  $C_2H_2$  and  $NH_3$  – during carbonitriding treatment of low alloy steels.

## Objective

Study precursors decomposition in order to control interstitial enrichment during steel carbonitriding.

## Methods

Gas chromatography, diffusion profile evaluation, thermogravimetry and numerical simulation.

# Introduction

## Pyrolysis

Thermochemical decomposition of organic material at high temperature in the absence of oxygen.

## Heterogeneous processes

Adsorption, desorption, decomposition and recombination processes of molecules at solid surfaces.

## Context

Decomposition of precursors –  $C_2H_2$  and  $NH_3$  – during carbonitriding treatment of low alloy steels.

## Objective

Study precursors decomposition in order to control interstitial enrichment during steel carbonitriding.

## Methods

Gas chromatography, diffusion profile evaluation, thermogravimetry and numerical simulation.

# Introduction

## Pyrolysis

Thermochemical decomposition of organic material at high temperature in the absence of oxygen.

## Heterogeneous processes

Adsorption, desorption, decomposition and recombination processes of molecules at solid surfaces.

## Context

Decomposition of precursors –  $\text{C}_2\text{H}_2$  and  $\text{NH}_3$  – during carbonitriding treatment of low alloy steels.

## Objective

Study precursors decomposition in order to control interstitial enrichment during steel carbonitriding.

## Methods

Gas chromatography, diffusion profile evaluation, thermogravimetry and numerical simulation.

# Introduction

## Pyrolysis

Thermochemical decomposition of organic material at high temperature in the absence of oxygen.

## Heterogeneous processes

Adsorption, desorption, decomposition and recombination processes of molecules at solid surfaces.

## Context

Decomposition of precursors –  $C_2H_2$  and  $NH_3$  – during carbonitriding treatment of low alloy steels.

## Objective

Study precursors decomposition in order to control interstitial enrichment during steel carbonitriding.

## Methods

Gas chromatography, diffusion profile evaluation, thermogravimetry and numerical simulation.

# Introduction

## Pyrolysis

Thermochemical decomposition of organic material at high temperature in the absence of oxygen.

## Heterogeneous processes

Adsorption, desorption, decomposition and recombination processes of molecules at solid surfaces.

## Context

Decomposition of precursors –  $C_2H_2$  and  $NH_3$  – during carbonitriding treatment of low alloy steels.

## Objective

Study precursors decomposition in order to control interstitial enrichment during steel carbonitriding.

## Methods

Gas chromatography, diffusion profile evaluation, thermogravimetry and numerical simulation.



# Table of Contents

- 1 Introduction
- 2 Residence time distribution
  - Background
  - Measurements
- 3 Acetylene decomposition
  - Atmospheric pressure
  - Pyrolysis mechanism
- 4 Ammonia decomposition
- 5 Ongoing studies
- 6 Conclusion

# Residence time distribution

## Background

**Purpose:** reactor diagnosis/modelling

**Reactor types:** ideal vs. real

**Concepts:**

- residence time distribution (RTD)
- mixing behavior
- reactor model


**Aim:** conversion prediction

**Measuring RTD:**

- pulse of tracer
- step of tracer

**Think!**

Portions of gas injected at a given instant leave the reactor at different moments: the conversion is averaged by these times but also by the mixing of arriving species at each position.



# Residence time distribution

## Background

**Purpose:** reactor diagnosis/modelling

**Reactor types:** ideal vs. real

**Concepts:**

- residence time distribution (RTD)
- mixing behavior
- reactor model


**Aim:** conversion prediction

**Measuring RTD:**

- pulse of tracer
- step of tracer

**Think!**

Portions of gas injected at a given instant leave the reactor at different moments: the conversion is averaged by these times but also by the mixing of arriving species at each position.



# Residence time distribution

## Background

**Purpose:** reactor diagnosis/modelling

**Reactor types:** ideal vs. real

**Concepts:**

- residence time distribution (RTD)
- mixing behavior
- reactor model


**Aim:** conversion prediction

**Measuring RTD:**

- pulse of tracer
- step of tracer

**Think!**

Portions of gas injected at a given instant leave the reactor at different moments: the conversion is averaged by these times but also by the mixing of arriving species at each position.



# Residence time distribution

## Background

**Purpose:** reactor diagnosis/modelling

**Reactor types:** ideal vs. real

**Concepts:**

- residence time distribution (RTD)
- mixing behavior
- reactor model


**Aim:** conversion prediction

**Measuring RTD:**

- pulse of tracer
- step of tracer

**Think!**

Portions of gas injected at a given instant leave the reactor at different moments: the conversion is averaged by these times but also by **the mixing of arriving species at each position.**



# Residence time distribution

## Background

**Purpose:** reactor diagnosis/modelling

**Reactor types:** ideal vs. real

**Concepts:**

- residence time distribution (RTD)
- mixing behavior
- reactor model


**Aim:** conversion prediction

**Measuring RTD:**

- pulse of tracer
- step of tracer

**Think!**

Portions of gas injected at a given instant leave the reactor at different moments: the conversion is averaged by these times but also by the mixing of arriving species at each position.



# Residence time distribution

## Background

**Purpose:** reactor diagnosis/modelling

**Reactor types:** ideal vs. real

**Concepts:**

- residence time distribution (RTD)
- mixing behavior
- reactor model


**Aim:** conversion prediction

**Measuring RTD:**

- pulse of tracer
- step of tracer

**Think!**

Portions of gas injected at a given instant leave the reactor at different moments: the conversion is averaged by these times but also by the mixing of arriving species at each position.



# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$



# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$

# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$

# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$

# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$

# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$

# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$

# Residence time distribution

## Measurements

**Method:** tracer ( $\text{CH}_4$ ) pulse in  $\text{N}_2$

**Detection:** FID (by-passing the column)


**Study parameters:**

- total flow rate
- heated zone temperature
- loading (with or without sample)

**Bodenstein number:** 5 – 10

**Mixing:** complete micromixing

**Model:** maximum mixedness



↑ flow rate  $\Rightarrow \downarrow \tau \downarrow \sigma$

↑ temperature  $\Rightarrow \downarrow \tau \sim \sigma$

# Table of Contents

- 1 Introduction
- 2 Residence time distribution
  - Background
  - Measurements
- 3 Acetylene decomposition
  - Atmospheric pressure
  - Pyrolysis mechanism
- 4 Ammonia decomposition
- 5 Ongoing studies
- 6 Conclusion



# Acetylene decomposition

Atmospheric pressure

**Input atmosphere:**  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



**Temperature range:** 823 K to 1223 K

**Flow rates:**  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

**Measured species:**  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

**Observations:**

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere



# Acetylene decomposition

Atmospheric pressure

**Input atmosphere:**  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



**Temperature range:** 823 K to 1223 K

**Flow rates:**  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

**Measured species:**  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

**Observations:**

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere



# Acetylene decomposition

Atmospheric pressure

**Input atmosphere:**  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



**Temperature range:** 823 K to 1223 K

**Flow rates:**  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

**Measured species:**  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

**Observations:**

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere



# Acetylene decomposition

Atmospheric pressure

**Input atmosphere:**  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



**Temperature range:** 823 K to 1223 K

**Flow rates:**  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

**Measured species:**  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

**Observations:**

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere



# Acetylene decomposition

Atmospheric pressure

Input atmosphere:  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



Temperature range: 823 K to 1223 K

Flow rates:  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

Measured species:  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

## Observations:

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere



# Acetylene decomposition

Atmospheric pressure

**Input atmosphere:**  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



**Temperature range:** 823 K to 1223 K

**Flow rates:**  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

**Measured species:**  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

## Observations:

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere



# Acetylene decomposition

Atmospheric pressure

**Input atmosphere:**  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



**Temperature range:** 823 K to 1223 K

**Flow rates:**  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

**Measured species:**  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

## Observations:

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere



# Acetylene decomposition

Atmospheric pressure

**Input atmosphere:**  $\text{N}_2 - 0.02 \text{ C}_2\text{H}_2$



**Temperature range:** 823 K to 1223 K

**Flow rates:**  $500 \text{ cm}^3 \text{ min}^{-1}$  and  $1000 \text{ cm}^3 \text{ min}^{-1}$

**Measured species:**  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$

## Observations:

- Decomposition detectable above 873 K
- Most important light-weight hydrocarbon formed below 1100 K:  $\text{C}_2\text{H}_4$
- At 1173 K:  $x(\text{C}_2\text{H}_2)$  is  $5\times$  other hydrocarbons and only 50% of the released H atoms are on  $\text{H}_2$  molecules
- $\text{C}_2\text{H}_6$  is not detected unless excess  $\text{H}_2$  is added to the input atmosphere





# Acetylene decomposition



Atmospheric pressure: mass balance

From the previous results one may establish a mole balance for carbon and hydrogen.

Since both TCD and FID detectors were used, total and hydrocarbon H-atoms can be quantified.

An increasing ratio C/H with respect to the temperature is observed for the missing atoms: C—C bonds are prevalent over C—H bonds in the undetected species.

Given the boundary and initial conditions, C/H ratio seems independent of the flow rate: the quantity is governed by the high temperature zone in the reactor.



# Acetylene decomposition



Atmospheric pressure: mass balance

From the previous results one may establish a mole balance for carbon and hydrogen.

Since both TCD and FID detectors were used, total and hydrocarbon H-atoms can be quantified.

An increasing ratio C/H with respect to the temperature is observed for the missing atoms: C—C bonds are prevalent over C—H bonds in the undetected species.

Given the boundary and initial conditions, C/H ratio seems independent of the flow rate: the quantity is governed by the high temperature zone in the reactor.



# Acetylene decomposition



Atmospheric pressure: mass balance

From the previous results one may establish a mole balance for carbon and hydrogen.

Since both TCD and FID detectors were used, total and hydrocarbon H-atoms can be quantified.

An increasing ratio C/H with respect to the temperature is observed for the missing atoms: C—C bonds are prevalent over C—H bonds in the undetected species.

Given the boundary and initial conditions, C/H ratio seems independent of the flow rate: the quantity is governed by the high temperature zone in the reactor.



# Acetylene decomposition



Atmospheric pressure: mass balance

From the previous results one may establish a mole balance for carbon and hydrogen.

Since both TCD and FID detectors were used, total and hydrocarbon H-atoms can be quantified.

An increasing ratio C/H with respect to the temperature is observed for the missing atoms: **C—C bonds are prevalent over C—H bonds in the undetected species.**

Given the boundary and initial conditions, C/H ratio seems independent of the flow rate: the quantity is governed by the high temperature zone in the reactor.



# Acetylene decomposition



Atmospheric pressure: mass balance

From the previous results one may establish a mole balance for carbon and hydrogen.

Since both TCD and FID detectors were used, total and hydrocarbon H-atoms can be quantified.

An increasing ratio C/H with respect to the temperature is observed for the missing atoms: C—C bonds are prevalent over C—H bonds in the undetected species.

Given the boundary and initial conditions, C/H ratio seems independent of the flow rate: the quantity is governed by the high temperature zone in the reactor.



# Acetylene decomposition



Atmospheric pressure: mass balance

From the previous results one may establish a mole balance for carbon and hydrogen.

Since both TCD and FID detectors were used, total and hydrocarbon H-atoms can be quantified.

An increasing ratio C/H with respect to the temperature is observed for the missing atoms: C—C bonds are prevalent over C—H bonds in the undetected species.

Given the boundary and initial conditions, C/H ratio seems independent of the flow rate: **the quantity is governed by the high temperature zone in the reactor.**



# Acetylene decomposition

Atmospheric pressure: carburizing


**Input atmosphere:**  $\text{N}_2 - 0.01 \text{ C}_2\text{H}_2$

**Temperature:** 1173 K

**Flow rate:**  $500 \text{ cm}^3 \text{ min}^{-1}$

## Observations:

- Metallic sample does not interfere in the order of magnitude of the observed hydrocarbons
- Good agreement between simulation and mass intake for a constant concentration boundary condition
- Even partial pressures below 5 mbar are enough for surface saturation



# Acetylene decomposition

Atmospheric pressure: carburizing


Input atmosphere:  $\text{N}_2 - 0.01 \text{ C}_2\text{H}_2$

Temperature: 1173 K

Flow rate:  $500 \text{ cm}^3 \text{ min}^{-1}$

## Observations:

- Metallic sample does not interfere in the order of magnitude of the observed hydrocarbons
- Good agreement between simulation and mass intake for a constant concentration boundary condition
- Even partial pressures below 5 mbar are enough for surface saturation





# Acetylene decomposition

Atmospheric pressure: carburizing


Input atmosphere:  $\text{N}_2 - 0.01 \text{ C}_2\text{H}_2$

Temperature: 1173 K

Flow rate:  $500 \text{ cm}^3 \text{ min}^{-1}$

## Observations:

- Metallic sample does not interfere in the order of magnitude of the observed hydrocarbons
- Good agreement between simulation and mass intake for a constant concentration boundary condition
- Even partial pressures below 5 mbar are enough for surface saturation



# Acetylene decomposition

Atmospheric pressure: carburizing


Input atmosphere:  $\text{N}_2 - 0.01 \text{ C}_2\text{H}_2$

Temperature: 1173 K

Flow rate:  $500 \text{ cm}^3 \text{ min}^{-1}$

## Observations:

- Metallic sample does not interfere in the order of magnitude of the observed hydrocarbons
- Good agreement between simulation and mass intake for a constant concentration boundary condition
- Even partial pressures below 5 mbar are enough for surface saturation



# Acetylene decomposition

Atmospheric pressure: carburizing


Input atmosphere:  $\text{N}_2 - 0.01 \text{ C}_2\text{H}_2$

Temperature: 1173 K

Flow rate:  $500 \text{ cm}^3 \text{ min}^{-1}$

## Observations:

- Metallic sample does not interfere in the order of magnitude of the observed hydrocarbons
- Good agreement between simulation and mass intake for a constant concentration boundary condition
- Even partial pressures below 5 mbar are enough for surface saturation



# Acetylene decomposition

Atmospheric pressure: carburizing


Input atmosphere:  $\text{N}_2 - 0.01 \text{ C}_2\text{H}_2$

Temperature: 1173 K

Flow rate:  $500 \text{ cm}^3 \text{ min}^{-1}$

## Observations:

- Metallic sample does not interfere in the order of magnitude of the observed hydrocarbons
- Good agreement between simulation and mass intake for a constant concentration boundary condition
- Even partial pressures below 5 mbar are enough for surface saturation



# Acetylene decomposition

Atmospheric pressure: RTD application

Using global rate parameters  $k = 1.5$  and  $n = 2.7$  for acetylene pyrolysis<sup>1</sup> coupled to experimental RTD functions with maximum mixedness model

$$\frac{dc_i}{dt} = - \left[ kc_i^n + (c_i - c_{i,0}) \frac{E(t_{max} - t)}{1 - F(t_{max} - t)} \right]$$

the following predictions are made (given in mole fractions):

Flow (cm <sup>3</sup> min <sup>-1</sup> )	Measured ×10 <sup>3</sup>	Simulated ×10 <sup>3</sup>	Ratio
250	3.67	3.01	0.82
500	4.25	4.18	0.98
1000	6.96	5.54	0.80

<sup>1</sup>Norinaga et al. Carbon 44 (2006) 1790–1800.

# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46

# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46

# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46



# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46

# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46

# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46

# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46

# Acetylene decomposition


## Pyrolysis mechanism

### Kinetic mechanism:

- Norinaga et al. J. Anal. Appl. Pyrolysis 86 (2009) 148-160
- 241 species (nodes)
- 902 reactions
- 1933 undirected edges

### Analyses:

- Simplification using DRG method
- System conditioning through interaction ordering
- Maximum mixedness solution of complete and simplified systems



Species	Degree
H	229
H <sub>2</sub>	210
C <sub>2</sub> H <sub>2</sub>	153
CH <sub>3</sub>	112
CH <sub>4</sub>	95
C <sub>2</sub> H <sub>3</sub>	87
C <sub>2</sub> H <sub>4</sub>	82
C <sub>6</sub> H <sub>6</sub>	56
C <sub>6</sub> H <sub>5</sub>	51
C <sub>4</sub> H <sub>4</sub>	46

# Table of Contents

- 1 Introduction
- 2 Residence time distribution
  - Background
  - Measurements
- 3 Acetylene decomposition
  - Atmospheric pressure
  - Pyrolysis mechanism
- 4 Ammonia decomposition
- 5 Ongoing studies
- 6 Conclusion

# Ammonia decomposition

**Pressure:** 1 bar


**Temperature range:** 773 K to 1223 K

**Flow rate:**  $415 \text{ cm}^3 \text{ min}^{-1}$

**Atmosphere:**  $0.24 \text{ N}_2 - 0.72 \text{ H}_2 - 0.04 \text{ NH}_3$

**Observations:**

- Decomposition break-through above 950 K
- Less than 10% of initial ammonia is reminiscent at carbonitriding temperatures



# Ammonia decomposition

Pressure: 1 bar


Temperature range: 773 K to 1223 K

Flow rate:  $415 \text{ cm}^3 \text{ min}^{-1}$

Atmosphere:  $0.24 \text{ N}_2 - 0.72 \text{ H}_2 - 0.04 \text{ NH}_3$

Observations:

- Decomposition break-through above 950 K
- Less than 10% of initial ammonia is reminiscent at carbonitriding temperatures





# Ammonia decomposition

**Pressure:** 1 bar


**Temperature range:** 773 K to 1223 K

**Flow rate:**  $415 \text{ cm}^3 \text{ min}^{-1}$

**Atmosphere:**  $0.24 \text{ N}_2 - 0.72 \text{ H}_2 - 0.04 \text{ NH}_3$

**Observations:**

- Decomposition break-through above 950 K
- Less than 10% of initial ammonia is reminiscent at carbonitriding temperatures



# Ammonia decomposition

**Pressure:** 1 bar


**Temperature range:** 773 K to 1223 K

**Flow rate:**  $415 \text{ cm}^3 \text{ min}^{-1}$

**Atmosphere:**  $0.24 \text{ N}_2 - 0.72 \text{ H}_2 - 0.04 \text{ NH}_3$

**Observations:**

- Decomposition break-through above 950 K
- Less than 10% of initial ammonia is reminiscent at carbonitriding temperatures



# Ammonia decomposition

**Pressure:** 1 bar


**Temperature range:** 773 K to 1223 K

**Flow rate:**  $415 \text{ cm}^3 \text{ min}^{-1}$

**Atmosphere:**  $0.24 \text{ N}_2 - 0.72 \text{ H}_2 - 0.04 \text{ NH}_3$

## Observations:

- Decomposition break-through above 950 K
- Less than 10% of initial ammonia is reminiscent at carbonitriding temperatures



# Ammonia decomposition

**Pressure:** 1 bar


**Temperature range:** 773 K to 1223 K

**Flow rate:**  $415 \text{ cm}^3 \text{ min}^{-1}$

**Atmosphere:**  $0.24 \text{ N}_2 - 0.72 \text{ H}_2 - 0.04 \text{ NH}_3$

## Observations:

- Decomposition break-through above 950 K
- Less than 10% of initial ammonia is reminiscent at carbonitriding temperatures



# Table of Contents

- 1 Introduction
- 2 Residence time distribution
  - Background
  - Measurements
- 3 Acetylene decomposition
  - Atmospheric pressure
  - Pyrolysis mechanism
- 4 Ammonia decomposition
- 5 Ongoing studies**
- 6 Conclusion

## Ongoing studies

Low pressure admission control for gas chromatography currently operating above 30 mbar.

Reliable measurements of  $C_2H_2$  and  $NH_3$  decomposition are currently being held.

Semi-automate system simplification routine for CFD simulations is under improvement.

Development of graph approach (pre-conditioning) to reduce stiffness of kinetic Jacobian.

Estimation of kinetic data from carburizing studies is being conducted.

## Ongoing studies

Low pressure admission control for gas chromatography currently operating above 30 mbar.

Reliable measurements of  $\text{C}_2\text{H}_2$  and  $\text{NH}_3$  decomposition are currently being held.

Semi-automate system simplification routine for CFD simulations is under improvement.

Development of graph approach (pre-conditioning) to reduce stiffness of kinetic Jacobian.

Estimation of kinetic data from carburizing studies is being conducted.

## Ongoing studies

Low pressure admission control for gas chromatography currently operating above 30 mbar.

Reliable measurements of  $\text{C}_2\text{H}_2$  and  $\text{NH}_3$  decomposition are currently being held.

Semi-automate system simplification routine for CFD simulations is under improvement.

Development of graph approach (pre-conditioning) to reduce stiffness of kinetic Jacobian.

Estimation of kinetic data from carburizing studies is being conducted.



## Ongoing studies

Low pressure admission control for gas chromatography currently operating above 30 mbar.

Reliable measurements of  $\text{C}_2\text{H}_2$  and  $\text{NH}_3$  decomposition are currently being held.

Semi-automate system simplification routine for CFD simulations is under improvement.

Development of graph approach (pre-conditioning) to reduce stiffness of kinetic Jacobian.

Estimation of kinetic data from carburizing studies is being conducted.

## Ongoing studies

Low pressure admission control for gas chromatography currently operating above 30 mbar.

Reliable measurements of  $\text{C}_2\text{H}_2$  and  $\text{NH}_3$  decomposition are currently being held.

Semi-automate system simplification routine for CFD simulations is under improvement.

Development of graph approach (pre-conditioning) to reduce stiffness of kinetic Jacobian.

Estimation of kinetic data from carburizing studies is being conducted.

# Table of Contents

- 1 Introduction
- 2 Residence time distribution
  - Background
  - Measurements
- 3 Acetylene decomposition
  - Atmospheric pressure
  - Pyrolysis mechanism
- 4 Ammonia decomposition
- 5 Ongoing studies
- 6 Conclusion

# Conclusion

Decomposition of  $C_2H_2$  leads to the formation of  $CH_4$ ,  $C_2H_4$  and  $H_2$ ;  $C_2H_6$  is observed under excess  $H_2$ .

Undetected species: C/H-ratio larger than unity, i.e. unsaturated character at intermediate levels (1-3) and the formation of soot (above 3).

System characterization (RTD) allowed the estimation of acetylene decomposition by using global rate parameters.

Graph analyses seem coherent with literature results<sup>2</sup> and allow the simplification of system kinetics for CFD simulations.

At atmospheric pressure austenitic nitriding conditions less than 10% of the injected ammonia is available to provide nitrogen to the material.

---

<sup>2</sup>F. Graf, PhD Thesis

# Conclusion

Decomposition of  $C_2H_2$  leads to the formation of  $CH_4$ ,  $C_2H_4$  and  $H_2$ ;  $C_2H_6$  is observed under excess  $H_2$ .

Undetected species: C/H-ratio larger than unity, i.e. unsaturated character at intermediate levels (1-3) and the formation of soot (above 3).

System characterization (RTD) allowed the estimation of acetylene decomposition by using global rate parameters.

Graph analyses seem coherent with literature results<sup>2</sup> and allow the simplification of system kinetics for CFD simulations.

At atmospheric pressure austenitic nitriding conditions less than 10% of the injected ammonia is available to provide nitrogen to the material.

---

<sup>2</sup>F. Graf, PhD Thesis

# Conclusion

Decomposition of  $C_2H_2$  leads to the formation of  $CH_4$ ,  $C_2H_4$  and  $H_2$ ;  $C_2H_6$  is observed under excess  $H_2$ .

Undetected species: C/H-ratio larger than unity, i.e. unsaturated character at intermediate levels (1-3) and the formation of soot (above 3).

System characterization (RTD) allowed the estimation of acetylene decomposition by using global rate parameters.

Graph analyses seem coherent with literature results<sup>2</sup> and allow the simplification of system kinetics for CFD simulations.

At atmospheric pressure austenitic nitriding conditions less than 10% of the injected ammonia is available to provide nitrogen to the material.

---

<sup>2</sup>F. Graf, PhD Thesis

# Conclusion

Decomposition of  $C_2H_2$  leads to the formation of  $CH_4$ ,  $C_2H_4$  and  $H_2$ ;  $C_2H_6$  is observed under excess  $H_2$ .

Undetected species: C/H-ratio larger than unity, i.e. unsaturated character at intermediate levels (1-3) and the formation of soot (above 3).

System characterization (RTD) allowed the estimation of acetylene decomposition by using global rate parameters.

Graph analyses seem coherent with literature results<sup>2</sup> and allow the simplification of system kinetics for CFD simulations.

At atmospheric pressure austenitic nitriding conditions less than 10% of the injected ammonia is available to provide nitrogen to the material.

---

<sup>2</sup>F. Graf, PhD Thesis

# Conclusion

Decomposition of  $\text{C}_2\text{H}_2$  leads to the formation of  $\text{CH}_4$ ,  $\text{C}_2\text{H}_4$  and  $\text{H}_2$ ;  $\text{C}_2\text{H}_6$  is observed under excess  $\text{H}_2$ .

Undetected species: C/H-ratio larger than unity, i.e. unsaturated character at intermediate levels (1-3) and the formation of soot (above 3).

System characterization (RTD) allowed the estimation of acetylene decomposition by using global rate parameters.

Graph analyses seem coherent with literature results<sup>2</sup> and allow the simplification of system kinetics for CFD simulations.

At atmospheric pressure austenitic nitriding conditions less than 10% of the injected ammonia is available to provide nitrogen to the material.

---

<sup>2</sup>F. Graf, PhD Thesis



Thanks for your attention!