

Recall that an **operator** is just a (continuous) linear transformation between two vectors spaces. A **Banach space** is a complete normed vector space X ; being complete means every Cauchy sequence in X converges to some vector in X . There are several notions of expansivity that have been recently studied:

Definition 0.1. Let X be a Banach space, let $S_X = \{x \in X : \|x\| = 1\}$ (meaning S_X is the unit sphere of X), and let $T : X \rightarrow X$ be an operator.

- We say T is **positively expansive** (PE) if for all $x \in S_X$, $\sup_{n \in \mathbb{N}} \|T^n x\| = +\infty$.
- We say T is **average positively expansive** (APE) if for all non-zero $x \in X$, $\sup_{n \in \mathbb{N}} \left(\frac{1}{n} \sum_{j=1}^n \|T^j x\| \right) = +\infty$.
- We say T is **uniformly positively expansive** (UPE) if $\lim_{n \rightarrow \infty} \|T^n x\| = +\infty$ uniformly on S_X , meaning for all $M > 0$, there exists $N \in \mathbb{N}$ such that for all $x \in S_X$ and for all $n \geq N$, $\|T^n x\| \geq M$.

There are reasons why operator theorists care about these definitions, but we'll ignore those reasons for now; what we are interested in is characterizing which forward and backward shifts are PE, APE, and UPE. It may look odd that APE doesn't restrict the x 's to the unit sphere S_X , but we leave it as an exercise to prove the following:

$$T \text{ is APE} \iff \forall x \in S_X, \sup_n \left(\frac{1}{n} \sum_{j=1}^n \|T^j x\| \right) = +\infty.$$

The exercise follows from the linearity of T , and at least one of these useful basic facts about suprema:

- If $c > 0$ and $A \subseteq \mathbb{R}$, then $\sup(cA) = c \sup(A)$.
- $\sup\{a_n : n \in \mathbb{N}\} = +\infty \iff$ for each $k \in \mathbb{N}$, $\sup\{a_n : n \geq k\} = +\infty$.

There are some equivalent characterizations of PE and UPE that were proved in other papers:

- T is PE $\iff \forall x \in S_X, \exists n \in \mathbb{N}$ s.t. $\|T^n x\| \geq 2$.
- T is UPE $\iff \exists n \in \mathbb{N}$ s.t. $\forall x \in S_X, \|T^n x\| \geq 2$.

Those characterizations highlight the difference between PE and UPE. We are not sure if there is an analogous characterization of APE, something like:

- T is APE $\iff \forall x \in S_X, \exists n \in \mathbb{N}$ s.t. $\sup_{n \in \mathbb{N}} \left(\frac{1}{n} \sum_{j=1}^n \|T^j x\| \right) \geq 2$.

Such a characterization could be worth exploring...

The last thing we'll mention is that it is straightforward to prove

$$\text{UPE} \implies \text{APE} \implies \text{PE},$$

and the reverse implications do not hold in general.

1 Positive Expansivity

From now on and for the rest of this document, let X be $c_0(\mathbb{N})$ or one of the spaces $\ell^p(\mathbb{N})$, $1 \leq p < \infty$. A backward shift on X defined by $B_w(x_1, x_2, x_3, \dots) = (w_2 x_2, w_3 x_3, w_4 x_4, \dots)$ is never PE. In fact, any operator T with a non-trivial kernel cannot be PE. If $Tx = 0$ for some non-zero vector x , then $\frac{x}{\|x\|}$ is a unit vector, i.e. $\frac{x}{\|x\|} \in S_X$, and $\|T^n(x/\|x\|)\| = 0$ for all n .

Let's see how forward shifts on X , defined by $F_w(x_1, x_2, x_3, \dots) = (0, w_1 x_1, w_2 x_2, \dots)$, can be PE. For a given basis vector e_v , note that $F_w^n e_v = w_v w_{v+1} \cdots w_{v+n-1} e_{v+n}$.

Proposition 1.1. F_w is PE on X if and only if $\sup_n |w_1 w_2 \cdots w_n| = +\infty$.

Proof. (\implies) Suppose F_w is PE. Then $e_1 \in S_X$, and since $F_w^n e_1 = w_1 w_2 \cdots w_n e_{n+1}$, by positive expansivity we have $\sup_n \|F_w^n e_1\| = \sup_n |w_1 w_2 \cdots w_n| = +\infty$.

(\impliedby) Suppose $\sup_n |w_1 w_2 \cdots w_n| = +\infty$, from which we can use basic facts about suprema to also conclude that for any $v \in \mathbb{N}$, $\sup_{n \geq v} |w_v \cdots w_{v+n-1}| = +\infty$. Let $x \in S_X$. Since $x \neq 0$, there exists some non-zero term x_v in the sequence x . Then

$$\|F_w^n x\| \geq \|F_w^n(x_v e_v)\| = |x_v| |w_v w_{v+1} \cdots w_{v+n-1}| = \frac{|x_v|}{|w_1 \cdots w_{v-1}|} |w_1 \cdots w_{v+n-1}|.$$

Whence $\sup_n \|F_w^n x\| = +\infty$. □

2 Average Positive Expansivity

Proposition 2.1. F_w is APE on X if and only if $\sup_n \left(\frac{1}{n} \sum_{j=1}^n |w_1 \cdots w_j| \right) = +\infty$.

Proof. (\implies) Suppose F is APE. Then $e_1 \in S_X$, and $F^j(e_1) = w_1 \cdots w_j$, so we must have

$$+\infty = \sup_n \left(\frac{1}{n} \sum_{j=1}^n \|F^j e_1\| \right) = \sup_n \left(\frac{1}{n} \sum_{j=1}^n |w_1 \cdots w_j| \right).$$

(\impliedby) Suppose $\sup_n \left(\frac{1}{n} \sum_{j=1}^n |w_1 \cdots w_j| \right) = +\infty$. Then by basic suprema facts, for any $k \in \mathbb{N}$, we know $\sup_{n \geq k} \left(\frac{1}{n+k} \sum_{j=1}^{n+k} |w_1 \cdots w_j| \right) = +\infty$. Let $x \in S_X$. Since $x \neq 0$, there exists some non-zero term x_v in the sequence x . Since

$$\begin{aligned} \frac{1}{n} \sum_{j=1}^n \|F^j x\| &\geq \frac{1}{n} \sum_{j=1}^n \|F^j(x_v e_v)\| \\ &= \frac{|x_v|}{n} \sum_{j=1}^n |w_v \cdots w_{v+j-1}| \\ &= \frac{1}{n} \frac{|x_v|}{|w_1 \cdots w_{v-1}|} \left(\frac{n+v-1}{n+v-1} \sum_{j=1}^{n+v-1} |w_1 \cdots w_j| - \sum_{j=1}^{v-1} |w_1 \cdots w_j| \right), \end{aligned}$$

and since $\lim_{n \rightarrow \infty} \frac{n+v-1}{n} = 1$ and $\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{j=1}^{v-1} |w_1 \cdots w_j| = 0$, we conclude that $\sup_n \left(\frac{1}{n} \sum_{j=1}^n \|F^j x\| \right) = +\infty$, which had to be shown. □

3 Uniform Positive Expansivity

Proposition 3.1. F_w is UPE on X if and only if $\lim_{n \rightarrow \infty} \left(\inf_{v \in \mathbb{N}} |w_v \cdots w_{v+n-1}| \right) = +\infty$.

Proof. (\implies) Suppose F_w is UPE. Let $M > 0$ be given. By UPE, there exists some $N \in \mathbb{N}$ such that for all $x \in S_X$ and all $n \geq N$, $\|F_w^n x\| > M$. In particular, for each unit vector e_v ,

we have $\|F_w^n e_v\| = |w_v \cdots w_{v+n-1}| > M$ whenever $n \geq N$. Hence for $n \geq N$, we can conclude

$$\inf_{v \in \mathbb{N}} |w_v \cdots w_{v+n-1}| \geq M,$$

which had to be shown.

(\Leftarrow) Suppose $\lim_{n \rightarrow \infty} (\inf_{v \in \mathbb{N}} |w_v \cdots w_{v+n-1}|) = +\infty$. Then there exists some $N \in \mathbb{N}$ such that for all $n > N$, $\inf_{v \in \mathbb{N}} |w_v \cdots w_{v+n-1}| \geq 2$. Thus for any $x \in S_X$, we have

$$\|F_w^n x\| = \sum_{v=1}^{\infty} |w_v \cdots w_{v+n-1}| \cdot |x_v| \geq 2 \sum_{v=1}^{\infty} |x_v| = 2\|x\|,$$

which shows F_w is UPE. □

4 Expansivity on Trees

We already have a characterization of which forward shifts are APE on directed trees. The next steps are to complete the following quests. First, recall a definition.

Definition 4.1. A linear transformation $T : X \rightarrow X$ is **frequently positively expansive** (FPE) if for each $x \in S_X$ and each $M > 0$, the set $\{n \in \mathbb{N}_0 : \|T^n x\| > M\}$ has positive lower density.

1. Let $X = \ell^1(V)$. Prove F_λ is UE on X iff $\lim_{n \rightarrow \infty} \left(\inf_{v \in V} \sum_{u \in \text{Chi}^n(v)} |\lambda(v \rightarrow u)| \right) = +\infty$. Does the theorem change if $X = \ell^p(V)$?
2. Let $X = \ell^1(V)$. Prove F_λ is PE on X iff $\forall v \in V, \sup_{n \in \mathbb{N}} \left(\sum_{u \in \text{Chi}^n(v)} |\lambda(v \rightarrow u)| \right) = +\infty$.
3. Can we characterize the UPE backward shifts? APE? PE? You can look up the definition of a backward shift in the directed trees paper. It might help to know that for $f \in \ell^1(V)$ (or any ℓ^p space or c_0), the n th iterate of the backward shift acting on f satisfies

$$(B_\lambda^n f)(v) = \sum_{u \in \text{Chi}^n(v)} \lambda(v \rightarrow u) f(u).$$

4. When is a backward shift on a tree conjugate to a backward shift on \mathbb{N} ?
5. Find an example of a PE forward shift on \mathbb{N} which is not FPE.
6. Find an example of an FPE forward shift on \mathbb{N} which is not UE. (I think we have done this).
7. Find an example an FPE forward shift on \mathbb{N} which is not UE but for each $x \in S_X$ and each $M > 0$, the set $\{n \in \mathbb{N}_0 : \|T^n x\| > M\}$ has lower density equal to 1.
8. Prove that UE implies FPE.
9. Does FPE imply APE?
10. Does APE imply FPE?