

Problem solving: kinematics (II)

Physics 211
Syracuse University, Physics 211 Spring 2015
Walter Freeman

February 1, 2016

Announcements

- Homework 2 due date is **this Friday**
- Exam 1 is next Tuesday
 - No homework due next week
 - Sample exam is posted; solutions posted Friday
 - Extended office hours and review sessions this week
 - Wednesday, 5-7 PM
 - Thursday 1:30-5:30
 - Friday, 10AM-4PM, exam review; location TBA
 - Weekend: Exam review in Stolkin, time TBA – vote in the Facebook poll

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- The exam will be substantially easier than the homework.

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- **The exam will be substantially easier than the homework.**
- You may use any calculator on the exam, but no cellphones or computers
- Students who do not speak English well: I will try to use only simple English on the exam, but if you like you may bring a dictionary
- Bring: your calculator, pencils, and your physics smarts (frog optional)

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- **The exam will be substantially easier than the homework.**
- You may use any calculator on the exam, but no cellphones or computers
- Students who do not speak English well: I will try to use only simple English on the exam, but if you like you may bring a dictionary
- Bring: your calculator, pencils, and your physics smarts (frog optional)
- Formal review session in class on Thursday
 - At this review you will create your reference sheet, which I will post in final form that day
- Other review times TBA (poll)

Exam 1, promises

- There will be one problem where you need the quadratic formula
 - ... this means interpreting the two values it spits out
- There will be at least one instance where you need to interpret or sketch position, velocity, and acceleration graphs
- You will *not* need to compute derivatives or integrals algebraically
- The exam will be four or five problems

- Vectors: objects with direction and magnitude.
 - Two representations:

- Vectors: objects with direction and magnitude.
 - Two representations:
 - Magnitude and direction (easiest to state, hardest to work with)
 - Components (easiest to work with)
 - Use trigonometry to go back and forth
- One more piece of notation about vectors...

Unit vectors

In the “ordered pair” notation for vectors’ components, you might write:

$$\vec{v} = (5, 3)$$

But this is clunky, if you’re trying to write it as part of an algebraic statement.

Instead we introduce “unit vectors”, vectors with length 1, in the x, y, and z directions.

$$\hat{i} = (1, 0, 0)$$

$$\hat{j} = (0, 1, 0)$$

$$\hat{k} = (0, 0, 1)$$

- $\vec{v} = (5, 3)$: Ordered pair
- $\vec{v} = 5\hat{i} + 3\hat{j}$: Unit vectors

Unit vectors

In the “ordered pair” notation for vectors’ components, you might write:

$$\vec{v} = (5, 3)$$

But this is clunky, if you’re trying to write it as part of an algebraic statement.

Instead we introduce “unit vectors”, vectors with length 1, in the x, y, and z directions.

$$\hat{i} = (1, 0, 0)$$

$$\hat{j} = (0, 1, 0)$$

$$\hat{k} = (0, 0, 1)$$

- $\vec{v} = (5, 3)$: Ordered pair
- $\vec{v} = 5\hat{i} + 3\hat{j}$: Unit vectors
- Both give you the same information, but unit vectors can be easier algebraically
- They won’t be essential for this class, but you should know the notation

Last time

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{r}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{r}_0$$

Last time

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{r}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{r}_0$$

$$\vec{v}_x(t) = a_xt + v_{x,0}$$

$$\vec{v}_y(t) = a_yt + v_{y,0}$$

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{r}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{r}_0$$

$$\vec{v}_x(t) = a_xt + v_{x,0}$$

$$\vec{v}_y(t) = a_yt + v_{y,0}$$

$$\vec{x}(t) = \frac{1}{2}a_xt^2 + v_{x,0}t + x_0$$

$$\vec{y}(t) = \frac{1}{2}a_yt^2 + v_{y,0}t + y_0$$

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

Example from cannon problem:

$$x(t) = \frac{1}{2}a_x t^2 + v_{x,0}t + x_0$$
$$y(t) = \frac{1}{2}a_y t^2 + v_{y,0}t + y_0$$

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

Example from cannon problem:

$$x(t) = v_{x,0}t$$

$$y(t) = -\frac{1}{2}gt^2 + v_{y,0}t$$

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

Example from cannon problem:

$$x(t) = v_0 \cos 45^\circ t$$

$$y(t) = -\frac{1}{2}gt^2 + v_0 \sin 45^\circ t$$

(I leave the rest to you for now...)

Problem solving: 2D kinematics, constant acceleration

1. If you have vectors in the “angle and magnitude” form $(\vec{a}, \vec{v}, \vec{r})$, convert them to components
2. Write down the kinematics relations, separately for x and y
 - Many terms will usually be zero
 - Freefall: $a_x = 0$, $a_y = -g$ (with conventional choice of axes)
3. Understand what instant in time you want to know about
4. Put in what you know; solve for what you don't (using substitution, if necessary)
5. Think about the physical meaning of your solution

Problem solving: 2D kinematics, constant acceleration

- ➊ 1. If you have vectors in the “angle and magnitude” form $(\vec{a}, \vec{v}, \vec{r})$, convert them to components
- ➋ 2. Write down the kinematics relations, separately for x and y
 - Many terms will usually be zero
 - Freefall: $a_x = 0$, $a_y = -g$ (with conventional choice of axes)
- ➌ 3. Understand what instant in time you know about (or want to know about)
- ➍ 4. Put in what you know; solve for what you don't (using substitution, if necessary)
- ➎ 5. Think about the physical meaning of your solution

“What instant in time do you know about?”

This is often the most difficult part of problems: it requires thought, not just math.

You throw a ball upward off of a cliff of height h . The top of the cliff is the origin, and up is positive.

What condition means “the ball has hit the ground”?

- A: $y = 0$
- B: $y = h$
- C: $y = -h$
- D: $v_y = 0$

“What instant in time do you know about?”

You throw a ball upward off of a cliff of height h . The top of the cliff is the origin, and up is positive.

What condition means “the ball is at its highest point?”?

- A: $y = 0$
- B: $v_y = 0$
- C: $y = h$
- D: y is a maximum

A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.

How can we frame the question “How far does the cannonball go?” in terms of our variables?

- A: What is x at the same time that v_x is zero?
- B: What is y at the same time that x is zero?
- C: What is x at the same time that y is zero?
- D: What is x at the same time that v_y is zero?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?
- How fast is it traveling when it strikes the ground?

A cannon

- A cannon shoots a cannonball at initial velocity v_0 at an angle θ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?

A cannon

- A cannon shoots a cannonball at initial velocity v_0 at an angle θ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?

A cannon

- A cannon shoots a cannonball at initial velocity v_0 at an angle θ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?

- A cannon shoots a cannonball at initial velocity v_0 at an angle θ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?
- How fast is it traveling when it strikes the ground?

Note that this algebraic solution can be used to do other things rather simply!

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red):

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red):
$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$
$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$
- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?

- Write down the basic position relations (terms that are zero in red):

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$
- We can find the time it takes to fall from the y equation:

$$0 = \frac{1}{2}(-g)t^2 + h$$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?

- Write down the basic position relations (terms that are zero in red):

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$

- We can find the time it takes to fall from the y equation:

$$0 = \frac{1}{2}(-g)t^2 + h$$

- Now we just do some algebra:

$$t = \sqrt{\frac{2h}{g}}$$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?

- Write down the basic position relations (terms that are zero in red):

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$
- We can find the time it takes to fall from the y equation:

$$0 = \frac{1}{2}(-g)t^2 + h$$

- Now we just do some algebra:

$$t = \sqrt{\frac{2h}{g}}$$

$$d = v_0 \sqrt{\frac{2h}{g}} \rightarrow v_0 = (30\text{m})\sqrt{\frac{(9.8\text{m/s}^2)}{(100\text{m}) \cdot 2}} = 6.64 \text{ m/s}$$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?

- Write down the basic position relations (terms that are zero in red):

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$
- We can find the time it takes to fall from the y equation:

$$0 = \frac{1}{2}(-g)t^2 + h$$

- Now we just do some algebra:

$$t = \sqrt{\frac{2h}{g}}$$

$$d = v_0 \sqrt{\frac{2h}{g}} \rightarrow v_0 = (30\text{m})\sqrt{\frac{(9.8\text{m/s}^2)}{(100\text{m}) \cdot 2}} = 6.64 \text{ m/s}$$

- To get the speed when it hits, we just use the velocity relations:

- $v_x = v_{0,x}$ and $v_y = -gt$

- $v_x = 6.64 \text{ m/s}$, $v_y = \sqrt{2gh} = -44.2 \text{ m/s}$

- $|v| = \sqrt{v_x^2 + v_y^2} = 44.7 \text{ m/s}$

The roadrunner problem

The position of the car is given by the ordinary 1D kinematics relation:


$$x(t) = \frac{1}{2}at^2 + v_0t = \frac{1}{2}(-9\frac{\text{m}}{\text{s}^2})t^2 + (30.6\frac{\text{m}}{\text{s}})t$$

The roadrunner problem

The position of the car is given by the ordinary 1D kinematics relation:

$$x(t) = \frac{1}{2}at^2 + v_0t = \frac{1}{2}\left(-9\frac{\text{m}}{\text{s}^2}\right)t^2 + \left(30.6\frac{\text{m}}{\text{s}}\right)t$$

We care about the time when it meets up with the position of the roadrunner, which is 30m. So we set $x(t) = 30$ and solve.




This seems easy enough, but the quadratic formula gives us two solutions! What happened?

The roadrunner problem

The position of the car is given by the ordinary 1D kinematics relation:

$$x(t) = \frac{1}{2}at^2 + v_0t = \frac{1}{2}(-9m/s^2)t^2 + (30.6m/s)t$$

We care about the time when it meets up with the position of the roadrunner, which is 30m. So we set $x(t) = 30$ and solve.




The roadrunner problem

The position of the car is given by the ordinary 1D kinematics relation:

$$x(t) = \frac{1}{2}at^2 + v_0t = \frac{1}{2}(-9m/s^2)t^2 + (30.6m/s)t$$

We care about the time when it meets up with the position of the roadrunner, which is 30m. So we set $x(t) = 30$ and solve.



Moral of the story: mathematics is a very blunt tool!

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- Now the condition for “rock hits ground” requires some thought:

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- Now the condition for “rock hits ground” requires some thought:
- The rock hits the ground when $x(t) = -y(t)$

$$v_{0,x}t = \frac{1}{2}gt^2 \rightarrow t = \frac{2v_{0,x}}{g}$$

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- Now the condition for “rock hits ground” requires some thought:

- The rock hits the ground when $x(t) = -y(t)$

$$v_{0,x}t = \frac{1}{2}gt^2 \rightarrow t = \frac{2v_{0,x}}{g}$$

- This gives us $x(t) = \frac{2v_{0,x}^2}{g}$

- $y(t)$ will have the same magnitude: the Pythagorean theorem gives $|r| = 2\sqrt{2}\frac{v_{0,x}^2}{g}$

A rocket

A rocket is launched from rest on level ground. While its motor burns, it accelerates at 10 m/s^2 at an angle 30° below the vertical. After ten seconds its motor burns out and it follows a ballistic trajectory until it hits the ground.

How far does it go?