

Problem solving: kinematics

Physics 211
Syracuse University, Physics 211 Spring 2015
Walter Freeman

January 22, 2015

- Homework 2 is due next Wednesday
- The first *Mastering Physics* tutorial is due ASAP
- No new *MP* assignments are mandatory until after exam 1, but I will post some optional ones for review/study
- I will be testing clickers Friday at 2PM in Stolkin; we will use them next week
- The Facebook page is set up: see <https://www.facebook.com/groups/384100861768360/>
- The wiki pages now have discussion options available to you (you don't need an account)
 - Up to 2% extra credit for participation here and on the wiki in helping your peers understand things
- The course schedule is available on the wiki

Last time

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{r}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{r}_0$$

Last time

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{r}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{r}_0$$

$$\vec{v}_x(t) = a_x t + v_{x,0}$$

$$\vec{v}_y(t) = a_y t + v_{y,0}$$

Last time

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{r}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{r}_0$$

$$\vec{v}_x(t) = a_x t + v_{x,0}$$

$$\vec{v}_y(t) = a_y t + v_{y,0}$$

$$\vec{x}(t) = \frac{1}{2}a_x t^2 + v_{x,0}t + x_0$$

$$\vec{y}(t) = \frac{1}{2}a_y t^2 + v_{y,0}t + y_0$$

One more piece of notation

There's one piece of notation I didn't mention last time about vectors.

“Unit vectors”: vectors of length 1 along the x and y axes

- Vectors are written with an arrow: \vec{r} , \vec{F} , \vec{a}
- Unit vectors are written with a hat: \hat{i} , \hat{j} , \hat{k}
- $\hat{i} = (1, 0, 0)$, $\hat{j} = (0, 1, 0)$, $\hat{k} = (0, 0, 1)$

So we might write: $\vec{V} = -3\hat{i} + 4\hat{j} + 2\hat{k}$. This describes the vector $\vec{V} = (-3, 4, 2)$.

Problem solving: 2D kinematics, constant acceleration

- ➊ 1. If you have vectors in the “angle and magnitude” form, convert them to components
- ➋ 2. Write down the kinematics relations, separately for x and y
 - Many terms will usually be zero
 - Freefall: $a_x = 0$, $a_y = -g$ (with conventional choice of axes)
- ➌ 3. Understand what instant in time you want to know about
- ➍ 4. Put in what you know; solve for what you don't (using substitution, if necessary)
- ➎ 5. Convert vectors into whatever format the problem asks for

- Today: solve problems involving 2D kinematics and projectile motion
- Next Tuesday: introduce rotational kinematics (it's easy); more problems
- Next Thursday: review for Exam 1
- *Tuesday, 3 Feb: Exam 1*

A reminder: there are lots of resources available for you.
(I was in the Clinic all day Tuesday helping folks, for instance.)

A simple problem: the cart and ball

- If the cart launcher propels the ball straight up at 2 m/s , how long does it take to come back down?

A simple problem: the cart and ball

- If the cart launcher propels the ball straight up at 2 m/s , how long does it take to come back down?
- How high does it go?

A simple problem: the cart and ball

- If the cart launcher propels the ball straight up at 2 m/s , how long does it take to come back down?
- How high does it go?
- How fast is it going when it lands?

A simple problem: the cart and ball

- If the cart launcher propels the ball straight up at 2 m/s , how long does it take to come back down?
- How high does it go?
- How fast is it going when it lands?
- Now let's say the cart is moving at 1.2 m/s when it shoots the ball out. Where does it land?

A cannon

- A cannon shoots a cannonball at 80 m/s at angles of 30 and 45 degrees above the horizontal.
- If the cannon is on level ground, how far do the cannonballs go?

A cannon

- A cannon shoots a cannonball at 80 m/s at angles of 30 and 45 degrees above the horizontal.
- If the cannon is on level ground, how far do the cannonballs go?
- How high does the cannonball go?

A cannon

- A cannon shoots a cannonball at initial velocity v_0 at an angle θ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?

A cannon

- A cannon shoots a cannonball at initial velocity v_0 at an angle θ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red): $x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$
 $y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red): $x(t) = \frac{1}{2}a_x t^2 + v_{0,x} t + x_0$
 $y(t) = \frac{1}{2}a_y t^2 + v_{0,y} t + y_0$
- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red): $x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$
 $y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$
- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$
- We can find the time it takes to fall from the y equation:
 $0 = \frac{1}{2} - gt^2 + h$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red): $x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$
 $y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$
- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$
- We can find the time it takes to fall from the y equation:
 $0 = \frac{1}{2} - g t^2 + h$
- Now we just do some algebra:
 $t = \sqrt{\frac{2h}{g}}$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red): $x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$
 $y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$
- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$
- We can find the time it takes to fall from the y equation:
 $0 = \frac{1}{2} - g t^2 + h$
- Now we just do some algebra:
 $t = \sqrt{\frac{2h}{g}}$
 $d = v_0 \sqrt{\frac{2h}{g}} \rightarrow v_0 = 30 \sqrt{\frac{9.8}{100 \cdot 2}} = 6.64 \text{ m/s}$

Throwing a rock off a cliff

- A hiker throws a rock horizontally off of a 100m tall cliff. If the rock strikes the ground 30m away, how hard did she throw it? How fast was it going when it hit the ground?
- Write down the basic position relations (terms that are zero in red): $x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$
 $y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$
- The first one gives us a very simple relationship between the horizontal distance d travelled and the time the rock takes to fall: $d = v_0 t$
- We can find the time it takes to fall from the y equation:
 $0 = \frac{1}{2} - g t^2 + h$
- Now we just do some algebra:
 $t = \sqrt{\frac{2h}{g}}$
 $d = v_0 \sqrt{\frac{2h}{g}} \rightarrow v_0 = 30 \sqrt{\frac{9.8}{100 \cdot 2}} = 6.64 \text{ m/s}$
- To get the speed when it hits, we just use the velocity relations:
- $v_x = v_{0,x}$ and $v_y = -gt$
- $v_x = 6.64 \text{ m/s}$, $v_y = \sqrt{2gh} = -44.2 \text{ m/s}$
- $|v| = \sqrt{v_x^2 + v_y^2} = 44.7 \text{ m/s}$

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- Now the condition for “rock hits ground” requires some thought:

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- Now the condition for “rock hits ground” requires some thought:
- The rock hits the ground when $x(t) = -y(t)$

$$v_{0,x}t = \frac{1}{2}gt^2 \rightarrow t = \frac{2v_{0,x}}{g}$$

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land?

- Same kinematic relations as before (this is always a good starting point!)

$$x(t) = \frac{1}{2}a_x t^2 + v_{0,x}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{0,y}t + y_0$$

- Now the condition for “rock hits ground” requires some thought:

- The rock hits the ground when $x(t) = -y(t)$

$$v_{0,x}t = \frac{1}{2}gt^2 \rightarrow t = \frac{2v_{0,x}}{g}$$

- This gives us $x(t) = \frac{2v_{0,x}^2}{g}$

- $y(t)$ will have the same magnitude: the Pythagorean theorem gives $|r| = 2\sqrt{2}\frac{v_{0,x}^2}{g}$

A rocket

A rocket is launched from rest on level ground. While its motor burns, it accelerates at 10 m/s^2 at an angle 30° below the vertical. After ten seconds its motor burns out and it follows a ballistic trajectory until it hits the ground.

How far does it go?