

Newton's Law of Motion

Physics 211
Syracuse University, Physics 211 Spring 2017
Walter Freeman

February 21, 2022

- Homework 3 due Friday
- You will get new groups in recitation tomorrow
- Exam misgrades/grade appeals: talk to your TA's
- I got a migraine on Friday and didn't get the check-in done :(
 - If you have any comments for me, please send them to me by email and I'll answer
 - I still owe you answers to the things you sent in last week; I'll do that today

Clinic hours this week

My availability in the Physics Clinic is a bit limited this week because of other commitments.

You can find me in the Clinic:

- Today (Tuesday), 1:30-3:30 pm
- Today (Tuesday), 5:15-6:15 (if possible)
- Wednesday, 2-4 pm
- Thursday, 9:45-10:30 am

Many of you asked for evening help hours. I am working to get coaches available for this and I'll let you know their availability by mail tonight.

Rational mechanics must be the science of the motions which result from any forces, and of the forces which are required for any motions, accurately propounded and demonstrated. For many things induce me to suspect, that all natural phenomena may depend upon some forces by which the particles of bodies are either drawn towards each other, and cohere, or repel and recede from each other: and these forces being hitherto unknown, philosophers have pursued their researches in vain. And I hope that the principles expounded in this work will afford some light, either to this mode of philosophizing, or to some mode which is more true.

-Isaac Newton, *Philosophiae Naturalis Principia Mathematica* (1687), translated from the Latin by Whewell (1837)

*Mechanics involves figuring out how things move from knowing the forces that act on them, and figuring out what forces act on them if we know how they move. I suspect that all physical things involve things exerting forces on each other, and since we don't know what forces these are, nobody's been able to figure much out. Hopefully someone will read this book and figure this stuff out, either following my suspicion that it's all forces under the hood (**classical physics!**), or with some deeper understanding of nature (**quantum physics!**)*

-Isaac Newton, *Philosophiae Naturalis Principia Mathematica*, in modern English

Summary from last time

- Forces: anything that pushes or pulls
- Forces cause accelerations: $\sum \vec{F} = m\vec{a}$
 - If $\sum \vec{F} = 0$, $\vec{a} = 0$: motion at a constant velocity
- Forces come in pairs: if A pushes on B, B pushes back on A
- It's the vector sum $\sum \vec{F}$ that matters
- Draw force diagrams to keep all of this straight

An important note

- Only *real physical things* are forces

An important note

- Only *real physical things* are forces
 - Acceleration is not a force
 - “Net force” is not a force (it’s the sum of them)
 - Velocity certainly isn’t a force
- Message phy211staff Choose FilesNo file chosen

An important note

- Only *real physical things* are forces
- Acceleration is not a force
- “Net force” is not a force (it’s the sum of them)
- Velocity certainly isn’t a force
- If two things don’t touch, or interact by gravity, electricity, etc., they don’t exchange forces

An important note

- Only *real physical things* are forces
- Acceleration is not a force
- “Net force” is not a force (it’s the sum of them)
- Velocity certainly isn’t a force
- If two things don’t touch, or interact by gravity, electricity, etc., they don’t exchange forces
- “A force is something that can send you to the doctor”

Questions from homework or last week's recitation?

Force is a vector; handle it like any other

One copy of Newton's second law in each direction (per object)

$$\vec{F} = m\vec{a} \rightarrow \begin{pmatrix} F_x = ma_x \\ F_y = ma_y \end{pmatrix}$$

Forces in 2D (and 3D)

Force is a vector; handle it like any other

One copy of Newton's second law in each direction (per object)

$$\vec{F} = m\vec{a} \rightarrow \begin{pmatrix} F_x = ma_x \\ F_y = ma_y \end{pmatrix}$$

Important: When dealing with inclines, choose your axes to align with the incline! (F_N is easy that way)

Force is a vector; handle it like any other

One copy of Newton's second law in each direction (per object)

$$\vec{F} = m\vec{a} \rightarrow \begin{pmatrix} F_x = ma_x \\ F_y = ma_y \end{pmatrix}$$

Forces in 2D (and 3D)

Force is a vector; handle it like any other

One copy of Newton's second law in each direction (per object)

$$\vec{F} = m\vec{a} \rightarrow \begin{pmatrix} F_x = ma_x \\ F_y = ma_y \end{pmatrix}$$

Important: When dealing with inclines, choose your axes to align with the incline! (F_N is easy that way)

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't
- **Kinematics:** Connect acceleration to motion

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't
- **Kinematics:** Connect acceleration to motion

It really is this easy; I promise!

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't
- **Kinematics:** Connect acceleration to motion

It really is this easy; I promise!
“Ask physics the question, don't tell it the answer”

Sample questions: dealing with two dimensions

A stone hangs from the roof of a car by a string; the car accelerates forward at 3 m/s^2 .

- What happens to the string?

Sample questions: dealing with two dimensions

A stone hangs from the roof of a car by a string; the car accelerates forward at 3 m/s^2 .

- What happens to the string?
- What angle does the string make with the vertical?

Sample questions: dealing with two dimensions

A stone hangs from the roof of a car by a string; the car accelerates forward at 3 m/s^2 .

- What happens to the string?
- What angle does the string make with the vertical?
- What is the tension in the string?

Sample questions: dealing with two dimensions

A cart slides down a frictionless track elevated at angle θ ; what is its acceleration?

Sample questions: dealing with multiple objects

Two masses of m_1 and m_2 kg hang from a massless pulley on either side. How do they move?

Sample questions: dealing with multiple objects

What if we put one object on a ramp and another object suspended from the table?

Nothing changes here – just follow the same procedure as always!

- Forces: anything that pushes or pulls
- Forces cause accelerations: $\sum \vec{F} = m\vec{a}$
 - If $\sum \vec{F} = 0$, $\vec{a} = 0$: motion at a constant velocity
- Forces come in pairs: if A pushes on B, B pushes back on A
- It's the vector sum $\sum \vec{F}$ that matters
- Draw force diagrams to keep all of this straight