

Friction (and sundry)

Physics 211
Syracuse University, Physics 211 Spring 2015
Walter Freeman

February 17, 2015

- Homework 4 extended until Friday
- *Mastering Physics* assignment due Thursday
- Read Chapter 8

- Homework 4 extended until Friday
- *Mastering Physics* assignment due Thursday
- Read Chapter 8
- Exam date change: *let me know* if there's a problem with moving it
 - Feb 26 → March 3

Ask a Physicist: How does an airplane work?

Homework review: Two objects moving

- Demo: Atwood's machine
- Same as always: Force diagram \rightarrow Newton's law for each object \rightarrow solve
- Two different objects \rightarrow two different accelerations!

$$T_1 - m_1 g = m_1 a_1$$

$$T_2 - m_2 g = m_2 a_2$$

Homework review: Two objects moving

- Demo: Atwood's machine
- Same as always: Force diagram \rightarrow Newton's law for each object \rightarrow solve
- Two different objects \rightarrow two different accelerations!

$$T_1 - m_1 g = m_1 a_1$$

$$T_2 - m_2 g = m_2 a_2$$

$$T_1 = T_2$$

Homework review: Two objects moving

- Demo: Atwood's machine
- Same as always: Force diagram \rightarrow Newton's law for each object \rightarrow solve
- Two different objects \rightarrow two different accelerations!

$$T_1 - m_1 g = m_1 a_1$$

$$T_2 - m_2 g = m_2 a_2$$

$$T_1 = T_2$$

$$a_1 = -a_2 \text{ ("standard coordinate system")}$$

$$a_1 = a_2 \text{ ("reverse coordinate system")}$$

Often things in nature are constrained to go in circles:

- Planets orbiting stars; moons orbiting planets (close enough to circles)
- Wheels; things on strings; many others

We'll study “uniform circular motion” here:

- Something moves at a constant distance from a fixed point
- ... at a constant speed.

Rotational motion: description

- Object moves in a circle of radius r , with its angle changing at a constant rate
- “Position = rate \times time” \rightarrow “Angle = rate \times time”

Rotational motion: description

- Object moves in a circle of radius r , with its angle changing at a constant rate
- “Position = rate \times time” \rightarrow “Angle = rate \times time”

$$\theta = \omega t$$

ω (omega) called the “angular velocity”; measured in radians per second

- Angular velocity tells you how fast something spins: RPM's are another unit
- A larger radius does *not* mean something has a higher angular velocity

Some new terms:

- “Radial”: directed in and out of the circle
- “Tangential”: directed around the circle
- The radial velocity is 0 (r doesn't change)
- The tangential velocity depends on r and ω , as you'd expect



Radians

The radian: new unit of angle. 2π radians = 360 degrees.

1 complete circle is 2π radians; 1 complete circumference is 2π radiuses ($C = 2\pi r$).

1 radian thus has an arc length of 1 radius.

θ radians therefore have an arc length of $r\theta$.



→ Tangential movement (in meters) = angular movement (in radians) times the radius

Some new terms:



- “Radial”: directed in and out of the circle
- “Tangential”: directed around the circle
- The radial velocity is 0 (r doesn't change)
- The tangential velocity depends on r and ω , as you'd expect

Some new terms:

- “Radial”: directed in and out of the circle
- “Tangential”: directed around the circle
- The radial velocity is 0 (r doesn’t change)
- The tangential velocity depends on r and ω , as you’d expect
- $v_T = \omega r$: “meters per second = radians per second times meters per radian”



Kinematic challenge: what's the acceleration

Clearly an object moving in a circle is accelerating. What's the acceleration?



Kinematic challenge: what's the acceleration

Clearly an object moving in a circle is accelerating. What's the acceleration?



Near the top of the circle, the y -component of the velocity decreases; we expect then that \vec{a} points downward.

Can we make this rigorous?

Some math

$$x(t) = r \cos(\omega t) \quad y(t) = r \sin(\omega t)$$

Some math

$$x(t) = r \cos(\omega t) \quad y(t) = r \sin(\omega t)$$

Differentiate to get v_x and v_y :

$$v_x(t) = -\omega r \sin(\omega t)$$

$$v_y(t) = \omega r \cos(\omega t)$$

Some math

$$x(t) = r \cos(\omega t) \quad y(t) = r \sin(\omega t)$$

Differentiate to get v_x and v_y :

$$v_x(t) = -\omega r \sin(\omega t)$$

$$v_y(t) = \omega r \cos(\omega t)$$

Differentiate again to get a_x and a_y :

$$a_x(t) = -\omega^2 r \cos(\omega t) = -\omega^2 x(t)$$

$$a_y(t) = -\omega^2 r \sin(\omega t) = -\omega^2 y(t)$$

Some math

$$x(t) = r \cos(\omega t) \quad y(t) = r \sin(\omega t)$$

Differentiate to get v_x and v_y :

$$v_x(t) = -\omega r \sin(\omega t)$$

$$v_y(t) = \omega r \cos(\omega t)$$

Differentiate again to get a_x and a_y :

$$a_x(t) = -\omega^2 r \cos(\omega t) = -\omega^2 x(t)$$

$$a_y(t) = -\omega^2 r \sin(\omega t) = -\omega^2 y(t)$$

$$\rightarrow \vec{a} = -\omega^2 \vec{r}$$

Some math

$$x(t) = r \cos(\omega t) \quad y(t) = r \sin(\omega t)$$

Differentiate to get v_x and v_y :

$$v_x(t) = -\omega r \sin(\omega t)$$

$$v_y(t) = \omega r \cos(\omega t)$$

Differentiate again to get a_x and a_y :

$$a_x(t) = -\omega^2 r \cos(\omega t) = -\omega^2 x(t)$$

$$a_y(t) = -\omega^2 r \sin(\omega t) = -\omega^2 y(t)$$

$$\rightarrow \vec{a} = -\omega^2 \vec{r}$$

An object in uniform circular motion accelerates toward the center of the circle with

$$\rightarrow \vec{a} = \omega^2 r = v^2/r \leftarrow$$

Uniform circular motion, consequences

If you know an object is undergoing uniform circular motion, you know something about the acceleration:

$$a = \omega^2 r \text{ or } a = v^2/r \text{ toward the center of the circle.}$$

Uniform circular motion, consequences

If you know an object is undergoing uniform circular motion, you know something about the acceleration:

$$a = \omega^2 r \text{ or } a = v^2/r \text{ toward the center of the circle.}$$

Circular motion problems aren't scary; they are just like equilibrium problems.

- Equilibrium problem: $\sum F_x = ma_x = 0$ and $\sum F_y = ma_y = 0$
- Circular motion problem: $\sum F_T = ma_T = 0$ and $\sum F_r = ma_r = v^2/r$

→ If we tell you that a thing is in uniform circular motion, we're just telling you something about its acceleration.

Centripetal force

“Centripetal” means “toward the center” in Latin.

- If something is going to accelerate toward the center, a force must do that.
- Centripetal force is **not** a “new” force. No arrows labeled “centripetal force”!
- “Centripetal” is a word that describes a force you already know about.
- Centripetal force: describes a force that holds something in a circle
- It can be lots of things:
 - Tension (stuffed animal on a string demo)
 - Normal force (platform, bucket demos)
 - Friction (Ferris wheel)
 - Gravity (the moon!)

Sample problems

(from demos)