

Momentum

Physics 211
Syracuse University, Physics 211 Spring 2022
Walter Freeman

April 5, 2022

- A reminder: The final exam is on **May 11**. You cannot take it early by University rules.
- Group Exam 3 on Friday; Exam 3 next Tuesday
- Review notes posted on the website
- Extra homework help hours today: 9:45-10:45, 2:30-3:30 PM, 5-7 PM (Physics Clinic – combination of me and others)
- HW7 due Friday

Recitation or homework questions?

Recitation or homework questions?

A request: “the one with the penguin” from last week

Sample problems: an excited dog

A person of mass m is sitting in a tire swing with a string of length L when their dog (mass M) runs and jumps horizontally into their lap.

If they swing up to an angle θ above the horizontal, how fast was their dog running?

Sample problems: “springs and momentum”

This was on 2020's practice exam (on a separate PDF)...

What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)

What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)

What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)
- Objects bounce “fully”: no kinetic energy lost (“elastic collision”)

What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)
- Objects bounce “fully”: no kinetic energy lost (“elastic collision”)
- Kinetic energy *increases*: only possible with some energy source!

What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)
- Objects bounce “fully”: no kinetic energy lost (“elastic collision”)
- Kinetic energy *increases*: only possible with some energy source!

The Newton's cradle

Remember:

- **All** collisions conserve momentum
- **Elastic** collisions conserve kinetic energy too
- **These** collisions are mostly elastic

The Newton's cradle

Remember:

- **All** collisions conserve momentum
- **Elastic** collisions conserve kinetic energy too
- **These** collisions are mostly elastic

Let's look at some examples from 2020's practice exam on this. For each one, tell me:

- A: Conserves momentum, but loses energy (inelastic collision)
- B: Conserves momentum and conserves energy (elastic collision)
- C: Conserves momentum, but gains energy (not possible)
- D: Does not conserve momentum or energy (not possible)
- E: Does not conserve momentum, but does conserve energy (not possible)

An application: neutron moderators

The only truly elastic collisions in nature are between particles. If we want *totally* elastic collisions, we should look to nuclear physics!

A note: this calculation we are going to do here demonstrates two things:

- How elastic collisions work
- ... and how *the art of approximation* is used in physics and engineering!

Recall how a nuclear reactor works:

- ^{235}U fissions when struck by neutrons with low energy (600 times more likely at low energy, less than 0.1 eV)
- When ^{235}U fissions, it produces neutrons with 2 MeV of kinetic energy ($v \approx 20$ million m/s)

How do we make these neutrons go from 2 million eV to 0.1 eV of kinetic energy so they can produce more fissions?

Bounce it off of other atoms!

Compared to a neutron moving at 2×10^6 m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

Bounce it off of other atoms!

Compared to a neutron moving at 2×10^6 m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass m and kinetic energy E_0 strikes another atom of mass M . They collide elastically. What is the neutron's energy after the collision?

Bounce it off of other atoms!

Compared to a neutron moving at 2×10^6 m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass m and kinetic energy E_0 strikes another atom of mass M . They collide elastically. What is the neutron's energy after the collision?

$$v_f = -v_0 \frac{M - m}{m + M}$$

Bounce it off of other atoms!

Compared to a neutron moving at 2×10^6 m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass m and kinetic energy E_0 strikes another atom of mass M . They collide elastically. What is the neutron's energy after the collision?

$$v_f = -v_0 \frac{M - m}{m + M}$$

How much kinetic energy is lost?

Bounce it off of other atoms!

Compared to a neutron moving at 2×10^6 m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass m and kinetic energy E_0 strikes another atom of mass M . They collide elastically. What is the neutron's energy after the collision?

$$v_f = -v_0 \frac{M - m}{m + M}$$

How much kinetic energy is lost?

The fraction of kinetic energy lost is

$$4 \frac{m}{M}.$$

So what atoms can we use as moderators?

They have to scatter neutrons more readily than they absorb them (hydrogen so-so, oxygen/carbon/heavy hydrogen great)

They have to be lightweight (that's what we just found)

They have to not be chemically grouchy (no hydrogen or oxygen by themselves!)

- H_2O (light water: most of the world, not the best moderator)
- D_2O (heavy water: Canadians)
- CO_2 (carbon dioxide: British)
- Pure carbon (Graphite: Soviets)