

Problem solving: kinematics (II)

Physics 211
Syracuse University, Physics 211 Spring 2015
Walter Freeman

January 30, 2017

- Homework 2 due date is **this Friday**
- Exam 1 is next Tuesday
 - No homework due next week
 - HW2 problems are similar to those on Exam 1
 - Physics practice: tomorrow night, 7:30-9:30, here
 - Topic: “setting up problems”, i.e. the actual physics
 - Recitation tomorrow has an extra credit question
 - Recitation Friday is your group practice exam
 - If you must miss Friday, notify your TA in advance
 - Weekend: Exam review in Stolkin, time TBA – vote in the Facebook poll

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- The exam will be somewhat easier than the homework.

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- **The exam will be somewhat easier than the homework.**
- You may use any ordinary calculator or graphing calculator on the exam, but no cellphones or computers, or Ti N-spire level devices
- Students who do not speak English well: I will try to use only simple English on the exam, but if you like you may bring a dictionary
- Bring: your calculator, pencils, and your physics smarts (frog optional)

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- **The exam will be somewhat easier than the homework.**
- You may use any ordinary calculator or graphing calculator on the exam, but no cellphones or computers, or Ti N-spire level devices
- Students who do not speak English well: I will try to use only simple English on the exam, but if you like you may bring a dictionary
- Bring: your calculator, pencils, and your physics smarts (frog optional)
- Review session in class on Thursday
- You are allowed to bring one page of notes that *you* *handwrite yourself*
 - No typed notes unless you have a disability that prevents you from writing
 - Your friend can't write it
 - You can't photocopy stuff from the book

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- **The exam will be somewhat easier than the homework.**
- You may use any ordinary calculator or graphing calculator on the exam, but no cellphones or computers, or Ti N-spire level devices
- Students who do not speak English well: I will try to use only simple English on the exam, but if you like you may bring a dictionary
- Bring: your calculator, pencils, and your physics smarts (frog optional)
- Review session in class on Thursday
- You are allowed to bring one page of notes that *you* *handwrite yourself*
 - No typed notes unless you have a disability that prevents you from writing
 - Your friend can't write it
 - You can't photocopy stuff from the book
 - It won't help you as much anyway

Exam 1, promises

- There will be one problem where you need the quadratic formula
 - ... this means interpreting the two values it spits out
- There will be at least one instance where you need to interpret or sketch position, velocity, and acceleration graphs
- You will *not* need to compute derivatives or integrals algebraically
- The exam will be four or five problems

- Vectors: objects with direction and magnitude.
 - Two representations:

- Vectors: objects with direction and magnitude.
 - Two representations:
 - Magnitude and direction (easiest to state, hardest to work with)
 - Components (easiest to work with)
 - Use trigonometry to go back and forth
- One more piece of notation about vectors...

Unit vectors

In the “ordered pair” notation for vectors’ components, you might write:

$$\vec{v} = (5, 3)$$

But this is clunky, if you’re trying to write it as part of an algebraic statement.

Instead we introduce “unit vectors”, vectors with length 1, in the x, y, and z directions.

$$\hat{i} = (1, 0, 0)$$

$$\hat{j} = (0, 1, 0)$$

$$\hat{k} = (0, 0, 1)$$

- $\vec{v} = (5, 3)$: Ordered pair
- $\vec{v} = 5\hat{i} + 3\hat{j}$: Unit vectors

Unit vectors

In the “ordered pair” notation for vectors’ components, you might write:

$$\vec{v} = (5, 3)$$

But this is clunky, if you’re trying to write it as part of an algebraic statement.

Instead we introduce “unit vectors”, vectors with length 1, in the x, y, and z directions.

$$\hat{i} = (1, 0, 0)$$

$$\hat{j} = (0, 1, 0)$$

$$\hat{k} = (0, 0, 1)$$

- $\vec{v} = (5, 3)$: Ordered pair
- $\vec{v} = 5\hat{i} + 3\hat{j}$: Unit vectors
- Both give you the same information, but unit vectors can be easier algebraically
- They won’t be essential for this class, but you should know the notation – other people use it quite a bit, and I use it once on your homework

A word on positive and negative acceleration, velocity, “speed”, and displacement:

When you choose your origin, you choose one direction to be positive, and the other to be negative. (Here: right = positive.)

- An object with $x < 0$ just means it's left of the origin.
- An object with $v < 0$ means it's moving to the left.
- An object with $a < 0$ means:
 - A: it is moving to the left and gaining speed
 - B: it is moving to the right and slowing down
 - C: it is moving to the left and slowing down
 - D: it is moving to the right and gaining speed

A word on positive and negative acceleration, velocity, “speed”, and displacement:

When you choose your origin, you choose one direction to be positive, and the other to be negative. (Here: right = positive.)

- An object with $x < 0$ just means it's left of the origin.
- An object with $v < 0$ means it's moving to the left.
- An object with $a < 0$ means:
 - A: it is moving to the left and gaining speed
 - B: it is moving to the right and slowing down
 - C: it is moving to the left and slowing down
 - D: it is moving to the right and gaining speed

Do not confuse the sign of something with the sign of its derivative!

Last time

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{s}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{s}_0$$

Last time

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{s}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{s}_0$$

$$v_x(t) = a_x t + v_{x,0}$$

$$v_y(t) = a_y t + v_{y,0}$$

Acceleration, velocity, and position relationships are the same in 2D; they just apply **independently** for each component.

$$\vec{v}(t) = \vec{a}t + \vec{v}_0$$

$$\vec{s}(t) = \frac{1}{2}\vec{a}t^2 + \vec{v}_0t + \vec{s}_0$$

$$v_x(t) = a_x t + v_{x,0}$$

$$v_y(t) = a_y t + v_{y,0}$$

$$x(t) = \frac{1}{2}a_x t^2 + v_{x,0}t + x_0$$

$$y(t) = \frac{1}{2}a_y t^2 + v_{y,0}t + y_0$$

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

Example from cannon problem:

$$x(t) = \frac{1}{2}a_x t^2 + v_{x,0}t + x_0$$
$$y(t) = \frac{1}{2}a_y t^2 + v_{y,0}t + y_0$$

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

Example from cannon problem:

$$x(t) = v_{x,0}t$$

$$y(t) = -\frac{1}{2}gt^2 + v_{y,0}t$$

If you don't know the numerical value of a quantity yet,
it's fine to leave it as a variable!

This is essential for solving many problems.

Example from cannon problem:

$$x(t) = v_0 \cos 45^\circ t$$

$$y(t) = -\frac{1}{2}gt^2 + v_0 \sin 45^\circ t$$

(I leave the rest to you for now...)

Problem solving: 2D kinematics, constant acceleration

1. If you have vectors in the “angle and magnitude” form $(\vec{a}, \vec{v}, \vec{s})$, convert them to components
2. Write down the kinematics relations, separately for x and y
 - Many terms will usually be zero
 - Freefall: $a_x = 0$, $a_y = -g$ (with conventional choice of axes)
3. Understand what instant in time you want to know about: ask the right question
4. Put in what you know; solve for what you don't (using substitution, if necessary)
5. Think about the physical meaning of your solution

Problem solving: 2D kinematics, constant acceleration

1. If you have vectors in the “angle and magnitude” form $(\vec{a}, \vec{v}, \vec{s})$, convert them to components
2. Write down the kinematics relations, separately for x and y
 - Many terms will usually be zero
 - Freefall: $a_x = 0$, $a_y = -g$ (with conventional choice of axes)
3. Understand what instant in time you want to know about: ask the right question
4. Put in what you know; solve for what you don't (using substitution, if necessary)
5. Think about the physical meaning of your solution

“What instant in time do you know about?”

This is often the most difficult part of problems: it requires thought, not just math.

You throw a ball upward over a hole of height h . Your position is the origin, and up is positive.

What condition means “the ball has hit the ground”?

- A: $y = 0$
- B: $y = h$
- C: $y = -h$
- D: $v_y = 0$

“What instant in time do you know about?”

You throw a ball upward off of a cliff of height h . The top of the cliff is the origin, and up is positive.

What condition means “the ball is at its highest point?”?

- A: $y = 0$
- B: $v_y = 0$
- C: $y = h$
- D: y is a maximum

A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.

How can we frame the question “How far does the cannonball go?” in terms of our variables?

- A: What is x at the same time that v_x is zero?
- B: What is y at the same time that x is zero?
- C: What is x at the same time that y is zero?
- D: What is x at the same time that v_y is zero?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?

A cannon

- A cannon shoots a cannonball at 80 m/s at an angle of 30 degrees above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?
- How fast is it traveling when it strikes the ground?

A cannon

- A cannon shoots a cannonball at initial velocity $v_0 = 80\text{m/s}$ at an angle $\theta = 30^\circ$ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?

A cannon

- A cannon shoots a cannonball at initial velocity $v_0 = 80\text{m/s}$ at an angle $\theta = 30^\circ$ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?

A cannon

- A cannon shoots a cannonball at initial velocity $v_0 = 80\text{m/s}$ at an angle $\theta = 30^\circ$ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?

A cannon

- A cannon shoots a cannonball at initial velocity $v_0 = 80\text{m/s}$ at an angle $\theta = 30^\circ$ above the horizontal.
- If the cannon is on level ground, how far does the cannonball go?
- How high does the cannonball go?
- How fast is it traveling at its highest point?
- How fast is it traveling when it strikes the ground?

What is $v_{0,x}$?

A: $v_0 \cos \theta$

B: $v_0 \sin \theta$

C: $v_0 \tan \theta$

D: v_0

A cannon

- What changes if I put the cannon up on a hill?

A cannon

- What changes if I put the cannon up on a hill?
- What changes if I'm trying to hit a target up on a plateau?

A cannon

- What changes if I put the cannon up on a hill?
- What changes if I'm trying to hit a target up on a plateau?
- What changes if I want to know what velocity I need to hit a target?

A cannon

- What changes if I put the cannon up on a hill?
- What changes if I'm trying to hit a target up on a plateau?
- What changes if I want to know what velocity I need to hit a target?
- What changes if I have air resistance?

Throwing a rock off a cliff

A hiker throws a rock horizontally off of a $h = 100$ m tall cliff. If the rock strikes the ground $d = 30$ m away, how hard did she throw it? How fast was it going when it hit the ground? (Choose the origin at the base of the cliff, up/direction of throw as positive)

What is $v_{0,x}$ here?

A: 0

B: $10/3$ m/s

C: You don't know *a priori*

What is $v_{0,y}$ here?

A: 0

B: 9.81 m/s

C: You don't know *a priori*

What is a_x here?

A: 0

B: -g

C: +g

D: You don't know *a priori*

What is a_y here?

A: 0

B: $-g$

C: $+g$

D: You don't know *a priori*

What is x_0 here?

A: 0

B: h

C: d

D: You don't know *a priori*

What is y_0 here?

A: 0

B: h

C: d

D: You don't know *a priori*

What question do you ask to find “how hard did she throw it?”

A: What value of $v_{x,0}$ makes it such that $x = d$ when $y = 0$?

B: What value of $v_{y,0}$ makes it such that $x = d$ when $y = h$?

C: What is the value of v_x when $y = 0$?

D: What is the magnitude of \vec{v} when $y = 0$?

E: What is the magnitude of \vec{v}_x when $y = h$?

What question do you ask to find “how fast is it going when it hits the ground?”

A: What is v_x at the time when $v_y = 0$?

B: What is v_x at the time when $y = 0$?

C: What is v_y at the time when $y = h$?

D: What is the magnitude of \vec{v} when $y = 0$?

E: What is the magnitude of \vec{v} when $y = h$?

What's the magnitude of \vec{v} ?

A: $v \cos \theta$

B: $v \sin \theta$

C: $\tan^{-1} \frac{v_x}{v_y}$

A: $\sqrt{v_x^2 + v_y^2}$

Throwing a stone onto a slope

A hiker kicks a stone off of a mountain slope with an initial velocity of v_0 3 m/s horizontally. If the mountain has a slope of 45 degrees, how far down the slope does it land? (Choose the origin as the starting point.)

A: What is the magnitude of \vec{s} when $x = y$?

B: What is the magnitude of \vec{s} when $x = -y$?

C: What is the magnitude of \vec{s} when $y = 0$?

D: What is y when $x = -y$?

E: What is y when $x = 0$?

A rocket is launched from rest on level ground. While its motor burns, it accelerates at 10 m/s^2 at an angle 30° below the vertical. After $\tau = 10 \text{ s}$ its motor burns out and it follows a ballistic trajectory until it hits the ground.

How far does it go?