

Problem solving: kinematics (II)

Physics 211
Syracuse University, Physics 211 Spring 2015
Walter Freeman

February 10, 2022

- Homework 2 due date is **tomorrow**
- Exam 1 is next Tuesday
 - No homework due next week
 - HW2 problems are similar to those on Exam 1
 - Recitation Friday is your group practice exam
 - If you must miss Friday, notify your TA in advance and explain to them why you will be absent; be prepared to provide documentation
 - If your absence is for a justified reason (see the syllabus), we will give you your grade on Exam 1 as a replacement for Group Exam 1
 - Saturday: Exam review in Stolkin, 5-8 PM

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- The exam will be somewhat easier than the homework.

Exam 1

- The exam covers kinematics in one and two dimensions
- Kinematics: how are an object's position, velocity, and acceleration related?
- The exam will be somewhat easier than the homework.
- There will be one problem where you need the quadratic formula
 - ... this means interpreting the two values it spits out
- There will be at least one instance where you need to interpret or sketch position, velocity, and acceleration graphs
- You will *not* need to compute derivatives or integrals algebraically
- The exam will be four or five problems

Exam 1, protocol

- You will have assigned seats; we will post a seating chart before the exam
- Some seats are left-handed; we'll ask you which hand you write with on Friday's checkin
- You are allowed to bring one page of notes that *you handwrite yourself*
 - No typed notes unless you have a disability that prevents you from writing
 - Your friend can't write it
 - You can't photocopy stuff from the book
 - It won't help you as much anyway

Problem solving: 2D kinematics, constant acceleration

1. If you have vectors in the “angle and magnitude” form $(\vec{a}, \vec{v}, \vec{s})$, convert them to components
2. Write down the kinematics relations, separately for x and y
 - Many terms will usually be zero
 - Freefall: $a_x = 0$, $a_y = -g$ (with conventional choice of axes)
3. Understand what instant in time you want to know about: ask the right question
4. Put in what you know; solve for what you don't (using substitution, if necessary)
5. Think about the physical meaning of your solution

Problem solving: 2D kinematics, constant acceleration

- ➊ 1. If you have vectors in the “angle and magnitude” form $(\vec{a}, \vec{v}, \vec{s})$, convert them to components
- ➋ 2. Write down the kinematics relations, separately for x and y
 - Many terms will usually be zero
 - Freefall: $a_x = 0$, $a_y = -g$ (with conventional choice of axes)
- ➌ 3. Understand what instant in time you want to know about: ask the right question
- ➍ 4. Put in what you know; solve for what you don't (using substitution, if necessary)
- ➎ 5. Think about the physical meaning of your solution

Do you have any questions from homework or recitation this week?

Throwing a rock off a cliff

A hiker throws a rock horizontally off of a $h = 100$ m tall cliff. If the rock strikes the ground $d = 30$ m away, how hard did she throw it? How fast was it going when it hit the ground? (Choose the origin at the base of the cliff, up/direction of throw as positive)

What is $v_{0,x}$ here?

A: 0

B: $10/3$ m/s

C: You don't know *a priori*

What is $v_{0,y}$ here?

A: 0

B: 9.8 m/s

C: You don't know *a priori*

What is a_x here?

A: 0

B: -g

C: +g

D: You don't know *a priori*

What is a_y here?

A: 0

B: $-g$

C: $+g$

D: You don't know *a priori*

What is x_0 here?

A: 0

B: h

C: d

D: You don't know *a priori*

What is y_0 here?

A: 0

B: h

C: d

D: You don't know *a priori*

What question do you ask to find “how hard did she throw it?”

A: What value of $v_{x,0}$ makes it such that $x = d$ when $y = 0$?

B: What value of $v_{y,0}$ makes it such that $x = d$ when $y = h$?

C: What is the value of v_x when $y = 0$?

D: What is the magnitude of \vec{v} when $y = 0$?

E: What is the magnitude of \vec{v}_x when $y = h$?

What question do you ask to find “how fast is it going when it hits the ground?”

A: What is v_x at the time when $v_y = 0$?

B: What is v_x at the time when $y = 0$?

C: What is v_y at the time when $y = h$?

D: What is the magnitude of \vec{v} when $y = 0$?

E: What is the magnitude of \vec{v} when $y = h$?

What's the magnitude of \vec{v} ?

A: $v \cos \theta$

B: $v \sin \theta$

C: $\tan^{-1} \frac{v_x}{v_y}$

A: $\sqrt{v_x^2 + v_y^2}$

Monkey and coconut (modified from “monkey and hunter”)

A monkey is hanging from a tree branch and is carrying a coconut. A human walks by, and the monkey throws it at them.

Monkey and coconut (modified from “monkey and hunter”)

A monkey is hanging from a tree branch and is carrying a coconut. A human walks by, and the monkey throws it at them.

They pick up the coconut and throw it back at the monkey. Not understanding gravity, they aim the coconut directly at the monkey. (This means that \vec{v}_0 is pointed directly at the monkey.)

Monkey and coconut (modified from “monkey and hunter”)

A monkey is hanging from a tree branch and is carrying a coconut. A human walks by, and the monkey throws it at them.

They pick up the coconut and throw it back at the monkey. Not understanding gravity, they aim the coconut directly at the monkey. (This means that \vec{v}_0 is pointed directly at the monkey.)

The monkey, being a clever sort and not wanting to get hit by the coconut, lets go of the tree branch as soon as the human releases the coconut; it hopes to fall to the ground and let the coconut fly over its head.

Monkey and coconut (modified from “monkey and hunter”)

A monkey is hanging from a tree branch and is carrying a coconut. A human walks by, and the monkey throws it at them.

They pick up the coconut and throw it back at the monkey. Not understanding gravity, they aim the coconut directly at the monkey. (This means that \vec{v}_0 is pointed directly at the monkey.)

The monkey, being a clever sort and not wanting to get hit by the coconut, lets go of the tree branch as soon as the human releases the coconut; it hopes to fall to the ground and let the coconut fly over its head.

Will the monkey dodge the coconut by doing this, or will the coconut return to monkey?

Kicking a football from a cliff

We return to our example football player from last class. They kick a football from the top of a 10-meter-tall cliff at 20 m/s at an angle 30° above the horizontal. Where does it land?

A rocket is launched from rest on level ground. While its motor burns, it accelerates at 10 m/s^2 at an angle 30° below the vertical. After $\tau = 10 \text{ s}$ its motor burns out and it follows a ballistic trajectory until it hits the ground.

How far does it go?